

On the solution of the massless Thirring model with fermion fields quantized in the chiral symmetric phase

M. Faber^{*} and A. N. Ivanov^{†‡}

February 1, 2008

Atominstytut der Österreichischen Universitäten, Arbeitsbereich Kernphysik und Nukleare Astrophysik, Technische Universität Wien, Wiedner Hauptstr. 8-10, A-1040 Wien, Österreich

Abstract

Correlation functions of fermionic fields described by the massless Thirring model are analysed within the operator formalism developed by Klaiber and the path-integral approach with massless fermions quantized in the chiral symmetric phase. We notice that Klaiber's composite fermion operators possess non-standard properties under parity transformations and construct operators with standard parity properties. We find that Klaiber's parameterization of a one-parameter family of solutions of the massless Thirring model is not well defined, since it is not consistent with the requirement of chiral symmetry. We show that the dynamical dimensions of correlation functions depend on an arbitrary parameter induced by ambiguities of the evaluation of the chiral Jacobian. A non-perturbative renormalization of the massless Thirring model is discussed. We demonstrate that the infrared divergences of Klaiber's correlation functions can be transferred into ultra-violet divergences by renormalization of the wave function of fermionic fields. This makes Klaiber's correlation functions non-singular in the infrared limit. We show that the requirement of non-perturbative renormalizability of the massless Thirring model fixes a free parameter of the path-integral approach. In turn, the operator formalism is inconsistent with non-perturbative renormalizability of the massless Thirring model. We carry out a non-perturbative renormalization of the sine-Gordon model and show that it is not an asymptotically free theory as well as the massless Thirring model. We calculate the fermion condensate by using the Fourier transform of the two-point Green function of massless Thirring fermion fields quantized in the chiral symmetric phase.

^{*}E-mail: faber@kph.tuwien.ac.at, Tel.: +43-1-58801-14261, Fax: +43-1-5864203

[†]E-mail: ivanov@kph.tuwien.ac.at, Tel.: +43-1-58801-14261, Fax: +43-1-5864203

[‡]Permanent Address: State Technical University, Department of Nuclear Physics, 195251 St. Petersburg, Russian Federation

1 Introduction

The massless Thirring model [1] is a theory of a self-coupled Dirac field $\psi(x)$

$$\mathcal{L}_{\text{Th}}(x) = \bar{\psi}(x)i\gamma^\mu\partial_\mu\psi(x) - \frac{1}{2}g\bar{\psi}(x)\gamma^\mu\psi(x)\bar{\psi}(x)\gamma_\mu\psi(x), \quad (1.1)$$

where g is a dimensionless coupling constant that can be both positive and negative as well. The field $\psi(x)$ is a spinor field with two components $\psi_1(x)$ and $\psi_2(x)$, x is a 2-vector $x^\mu = (x^0, x^1)$, where x^0 and x^1 are time and spatial components. The γ -matrices are defined in terms of the well-known 2×2 Pauli matrices σ_1 , σ_2 and σ_3

$$\gamma^0 = \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^1 = -i\sigma_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^5 = \gamma^0\gamma^1 = \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.2)$$

These γ -matrices obey the standard relations

$$\begin{aligned} \gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu &= 2g^{\mu\nu}, \\ \gamma^\mu\gamma^5 + \gamma^5\gamma^\mu &= 0. \end{aligned} \quad (1.3)$$

We use the metric tensor $g^{\mu\nu}$ defined by $g^{00} = -g^{11} = 1$ and $g^{01} = g^{10} = 0$. The axial-vector product $\gamma^\mu\gamma^5$ can be expressed in terms of γ^ν

$$\gamma^\mu\gamma^5 = -\epsilon^{\mu\nu}\gamma_\nu, \quad (1.4)$$

where $\epsilon^{\mu\nu}$ is the anti-symmetric tensor defined by $\epsilon^{01} = -\epsilon^{10} = 1$. Further, we also use the relation $\gamma^\mu\gamma^\nu = g^{\mu\nu} + \epsilon^{\mu\nu}\gamma^5$.

The Lagrangian (1.1) is obviously invariant under the chiral group $U_V(1) \times U_A(1)$

$$\begin{aligned} \psi(x) &\xrightarrow{V} \psi'(x) = e^{i\alpha_V}\psi(x), \\ \psi(x) &\xrightarrow{A} \psi'(x) = e^{i\alpha_A\gamma^5}\psi(x), \end{aligned} \quad (1.5)$$

where α_V and α_A are real parameters defining global rotations.

Due to invariance under chiral group $U_V(1) \times U_A(1)$ the vector and axial-vector current $j^\mu(x)$ and $j_5^\mu(x)$, induced by vector (V) and axial-vector (A) rotations and defined by

$$\begin{aligned} j^\mu(x) &= \bar{\psi}(x)\gamma^\mu\psi(x), \\ j_5^\mu(x) &= \bar{\psi}(x)\gamma^\mu\gamma^5\psi(x), \end{aligned} \quad (1.6)$$

are conserved

$$\partial_\mu j^\mu(x) = \partial_\mu j_5^\mu(x) = 0. \quad (1.7)$$

Recall, that in 1+1-dimensional field theories the vector and axial-vector currents are related by $j_5^\mu(x) = -\epsilon^{\mu\nu}j_\nu(x)$ due to the properties of Dirac matrices.

The massless Thirring model being a local quantum field theoretic model is invariant under C , P and T transformations separately. Since below we are concerned with the properties of fermion fields under parity transformations, we remind that invariance of the

massless Thirring model under parity transformation imposes the constraints [2,3] (see also Appendix A of this paper)

$$\begin{aligned}\mathcal{P}\mathcal{L}(x^0, x^1)\mathcal{P}^\dagger &= +\mathcal{L}(x^0, -x^1), \\ \mathcal{P}j^\mu(x^0, x^1)\mathcal{P}^\dagger &= +j_\mu(x^0, -x^1), \\ \mathcal{P}j_5^\mu(x^0, x^1)\mathcal{P}^\dagger &= -j_{5\mu}(x^0, -x^1).\end{aligned}\tag{1.8}$$

These constraints fix the transformation of the fermionic field [2,3]

$$\mathcal{P}\psi(x^0, x^1)\mathcal{P}^\dagger = \gamma^0\psi(x^0, -x^1)\tag{1.9}$$

up to an insignificant phase factor [2,3] which we have dropped.

The relation (1.9) will be important for the discussion of solutions of the Dirac equation for a free massless fermion field. We will show below that for a free massless fermionic field in 1+1-dimensional space-time the Dirac equation has two solutions. One of them possesses non-standard properties under parity transformations and does not obey the constraint (1.9). For the first time this solution has been obtained by Thirring in his seminal paper [1], though, without reference to non-standard properties under parity transformation.

As has been shown by Thirring [1] the model described by the Lagrangian (1.1) can be solved exactly. Thirring constructed the eigenstates of the Hamiltonian and calculated some observable quantities and found them to be finite after renormalization.

Of course, the exact solution of the massless Thirring model means also the possibility of an explicit evaluation of any correlation function of fermions. For the first time an explicit evaluation of any correlation function of the massless Thirring model has been carried out by Klaiber in the traditional quantum field theoretic way within the operator technique with fermion fields quantized in the chiral symmetric phase relative to the perturbative chiral symmetric vacuum [4]¹. According to Klaiber's statement the massless Thirring model can be reduced to a quantum field theory of a massless free fermionic field Ψ by a corresponding canonical transformation of the self-coupled fermionic field $\psi \rightarrow \Psi$. The two-point Green function $S_F(x - y)$ of the free massless fermionic field Ψ is defined by

$$S_F(x - y) = i\langle 0 | T(\Psi(x)\bar{\Psi}(y)) | 0 \rangle = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x - y)^2 - i0},\tag{1.10}$$

where $\hat{x} - \hat{y} = \gamma^\mu(x - y)_\mu$.

The main implicit point of Klaiber's approach for the evaluation of correlation functions in the massless Thirring model is the quantization of the self-coupled fermionic field ψ and the free field Ψ in the chiral symmetric phase [6]. As we have shown in Ref.[6] the massless Thirring model with an attractive coupling $g > 0$ is unstable under spontaneous breaking of chiral symmetry and energetic the chirally broken phase is preferable with respect to the chiral symmetric one. This implies that for $g > 0$ the self-coupled fermion fields ψ should be quantized in the chirally broken phase. The wave function of

¹The same technique has been used then by Lowenstein [5] who studied the short-distance behaviour of products of fermion fields in the massless Thirring model and the validity of Wilson's expansion hypothesis.

the non-perturbative vacuum coincides with the wave function of the ground state of the superconducting phase in the Bardeen, Cooper and Schrieffer theory of superconductivity (the BCS-theory) [6].

The problem of an exact solution of the massless Thirring model has been revised within the path-integral approach by Furuya, Gamboa Saravi and Schaposnik [7] and Naón [8]. These authors used the technique of auxiliary vector fields where the four-fermion interaction in the Lagrangian (1.1) is linearized by the inclusion of a local vector field V_μ . The main object of the path-integral approach is the generating functional of Green functions

$$Z_{\text{Th}}[J, \bar{J}] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp i \int d^2x \left[\bar{\psi}(x) i\gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) + \bar{\psi}(x) J(x) + \bar{J}(x) \psi(x) \right], \quad (1.11)$$

where $\bar{J}(x)$ and $J(x)$ are external sources of the $\psi(x)$ and $\bar{\psi}(x)$ fields.

The exponential of the four-fermion interaction can be replaced by the Gaussian integral over a vector field V_μ :

$$\begin{aligned} & \exp i \int d^2x \left[-\frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \right] = \\ & = \int \mathcal{D}V^\mu \exp i \int d^2x \left[\frac{1}{2g} V_\mu(x) V^\mu(x) + \bar{\psi}(x) \gamma^\mu \psi(x) V_\mu(x) \right]. \end{aligned} \quad (1.12)$$

Using the path-integral representation (1.12) the generating functional (1.11) can be recast into the form

$$Z_{\text{Th}}[J, \bar{J}] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}V^\mu \exp i \int d^2x \left[\bar{\psi}(x) i\gamma^\mu (\partial_\mu - iV_\mu(x)) \psi(x) + \frac{1}{2g} V_\mu(x) V^\mu(x) + \bar{\psi}(x) J(x) + \bar{J}(x) \psi(x) \right]. \quad (1.13)$$

The first step of the path-integral approach to the solution of the massless Thirring model suggested in Refs.[7,8] is in the use of the Hodge decomposition of the vector field $V_\mu(x)$

$$V_\mu(x) = \partial_\mu \eta(x) + \varepsilon_{\mu\nu} \partial^\nu \xi(x), \quad (1.14)$$

where $\eta(x)$ and $\xi(x)$ are a scalar and a pseudoscalar field, respectively.

Substituting the decomposition (1.14) in (1.13) one reduces the generating functional to the form

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] = & \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}\eta \mathcal{D}\xi \exp i \int d^2x \left[\bar{\psi}(x) i\gamma^\mu (\partial_\mu - i\partial_\mu \eta(x) - i\varepsilon_{\mu\nu} \partial^\nu \xi(x)) \psi(x) \right. \\ & \left. + \frac{1}{2g} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2g} \partial_\mu \xi(x) \partial^\mu \xi(x) + \bar{\psi}(x) J(x) + \bar{J}(x) \psi(x) \right], \end{aligned} \quad (1.15)$$

where we have used the relation $\varepsilon_{\mu\nu} \varepsilon^{\mu\alpha} = -g_\nu^\alpha$.

The second step of the path-integral approach to the solution of the massless Thirring model suggested in Refs.[7,8] is in the transformation of the fermion field

$$\begin{aligned} \psi(x) &= e^{i\eta(x) + i\gamma^5 \xi(x)} \Psi(x), \\ \bar{\psi}(x) &= \bar{\Psi}(x) e^{-i\eta(x) + i\gamma^5 \xi(x)}. \end{aligned} \quad (1.16)$$

This is an analogy of the canonical transformation suggested by Klaiber reducing the massless Thirring model of the self-coupled fermionic field ψ to a quantum field theory of a free massless fermionic field Ψ [4].

As a result the generating functional of the Green functions reads

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi J[\xi] \\ &\times \exp i \int d^2x \left[\bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2g} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2g} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\ &\left. + \bar{\Psi}(x) e^{-i\eta(x) + i\gamma^5 \xi(x)} J(x) + \bar{J}(x) e^{i\eta(x) + i\gamma^5 \xi(x)} \Psi(x) \right], \end{aligned} \quad (1.17)$$

where $J[\xi]$ is the Jacobian caused by the chiral part of the transformation (1.16).

In Appendix B we show that following the procedure developed in [9–14] the chiral Jacobian $J[\xi]$ can be given by the expression

$$J[\xi] = \exp \left[-i \int d^2x \frac{\alpha_J}{2\pi} \partial_\mu \xi(x) \partial^\mu \xi(x) \right], \quad (1.18)$$

where an arbitrary real parameter α_J reflects ambiguities of the evaluation of chiral Jacobians induced by different regularization procedures [11–13]².

Our expression for the chiral Jacobian (1.18) taken at $\alpha_J = 1$ coincides with the chiral Jacobian calculated by Damgaard, Nielsen and Sollacher [15]. Unlike Refs.[7,8] Damgaard, Nielsen and Sollacher [15] analysed quantum field theories in 1+1-dimensional Minkowski space-time as we are. The agreement can be reached setting $V^\mu(x) = \varepsilon^{\mu\nu} \partial_\nu \theta(x)$ in Eq.(14) of Ref.[15]. This change corresponds to a decoupling of the vector field $V^\mu(x)$ from massless fermion fields as suggested in Refs.[7,8]³.

Due to (1.18) the generating functional (1.17) can be rewritten as follows

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi \\ &\times \exp i \int d^2x \left[\bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2g} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2g} \left(1 + \alpha_J \frac{g}{\pi} \right) \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\ &\left. + \bar{\Psi}(x) e^{-i\eta(x) + i\gamma^5 \xi(x)} J(x) + \bar{J}(x) e^{i\eta(x) + i\gamma^5 \xi(x)} \Psi(x) \right]. \end{aligned} \quad (1.19)$$

Using the identity

$$e^{i\gamma^5 \xi(x)} = \left(\frac{1 + \gamma^5}{2} \right) e^{i\xi(x)} + \left(\frac{1 - \gamma^5}{2} \right) e^{-i\xi(x)} \quad (1.20)$$

one can rewrite the generating functional (1.19) in the more convenient form

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi \\ &\times \exp i \int d^2x \left\{ \bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2g} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2g} \left(1 + \alpha_J \frac{g}{\pi} \right) \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \end{aligned}$$

²In Refs.[13] ambiguities of fermionic determinants are investigated in the massless Schwinger model.

³Nice reviews of quantum field theories in 1+1-dimensional space-time one can find in books [16] and [17].

$$\begin{aligned}
& + \bar{\Psi}(x) \left[\left(\frac{1+\gamma^5}{2} \right) e^{-i\eta(x) + i\xi(x)} + \left(\frac{1-\gamma^5}{2} \right) e^{-i\eta(x) - i\xi(x)} \right] J(x) \\
& + \bar{J}(x) \left[\left(\frac{1+\gamma^5}{2} \right) e^{i\eta(x) + i\xi(x)} + \left(\frac{1-\gamma^5}{2} \right) e^{i\eta(x) - i\xi(x)} \right] \Psi(x) \}.
\end{aligned} \tag{1.21}$$

The last step in the transformation of the generating functional $Z_{\text{Th}}[J, \bar{J}]$ is to rescale the fields η and ξ : $\eta \rightarrow \sqrt{g}\eta$ and $\xi \rightarrow \lambda(\alpha_J)\xi$, where $\lambda(\alpha_J)$ is determined by

$$\lambda(\alpha_J) = \sqrt{\frac{g}{1 + \alpha_J \frac{g}{\pi}}} \tag{1.22}$$

As a result the generating functional of Green functions of the massless Thirring model $Z_{\text{Th}}[J, \bar{J}]$ is defined by the path integral

$$\begin{aligned}
Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi \\
&\times \exp i \int d^2x \left\{ \bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\
&+ \bar{\Psi}(x) \left[\left(\frac{1+\gamma^5}{2} \right) e^{-i\sqrt{g}\eta(x) + i\lambda(\alpha_J)\xi(x)} + \left(\frac{1-\gamma^5}{2} \right) e^{-i\sqrt{g}\eta(x) - i\lambda(\alpha_J)\xi(x)} \right] J(x) \\
&+ \bar{J}(x) \left[\left(\frac{1+\gamma^5}{2} \right) e^{i\sqrt{g}\eta(x) + i\lambda(\alpha_J)\xi(x)} + \left(\frac{1-\gamma^5}{2} \right) e^{i\sqrt{g}\eta(x) - i\lambda(\alpha_J)\xi(x)} \right] \Psi(x) \}.
\end{aligned} \tag{1.23}$$

It is important to emphasize that the kinetic term of the pseudoscalar field ξ enters with opposite sign. The free Lagrangian of the scalar and pseudoscalar fields $\eta(x)$ and $\xi(x)$ is invariant under global $O(1, 1)$ rotations

$$\begin{aligned}
\eta(x) \rightarrow \eta'(x) &= \cosh \gamma \eta(x) + \sinh \gamma \xi(x), \\
\xi(x) \rightarrow \xi'(x) &= \sinh \gamma \eta(x) + \cosh \gamma \xi(x).
\end{aligned} \tag{1.24}$$

In parallel to the generating functional of Green functions (1.11) one can consider the generating functional of correlation functions of the scalar fermionic field density $\bar{\psi}(x)\psi(x)$ determined by

$$\begin{aligned}
Z_{\text{Th}}[\sigma] &= \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \\
&\times \exp i \int d^2x \left[\bar{\psi}(x) i\gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) + \bar{\psi}(x) \psi(x) \sigma(x) \right],
\end{aligned} \tag{1.25}$$

where $\sigma(x)$ is an external source of $\bar{\psi}(x)\psi(x)$. Due to the transformations expounded above the generating functional $Z_{\text{Th}}[\sigma]$ can be reduced to the form of the following path integral

$$\begin{aligned}
Z_{\text{Th}}[\sigma] &= \\
&= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\xi \exp i \int d^2x \left[\bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right.
\end{aligned}$$

$$\begin{aligned}
& + \bar{\Psi}(x) e^{2i\lambda(\alpha_J) \gamma^5 \xi(x)} \Psi(x) \sigma(x) \Big] = \\
& = \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\xi \exp i \int d^2x \left\{ \bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\
& \left. + \bar{\Psi}(x) \left[\left(\frac{1+\gamma^5}{2} \right) e^{2i\lambda(\alpha_J) \xi(x)} + \left(\frac{1-\gamma^5}{2} \right) e^{-2i\lambda(\alpha_J) \xi(x)} \right] \Psi(x) \sigma(x) \right\}. \quad (1.26)
\end{aligned}$$

We emphasize that the η -field does not contribute to the correlation functions of the scalar fermion density $\bar{\psi}(x)\psi(x)$.

In order to consider the correlation functions, treated by Coleman [18] for a proof of equivalence between the massive Thirring model and the sine-Gordon, one can introduce the following generating functional

$$\begin{aligned}
Z_{\text{Th}}[\sigma, \varphi] = & \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp i \int d^2x \left[\bar{\psi}(x) i \gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \right. \\
& \left. + \bar{\psi}(x) \psi(x) \sigma(x) + \bar{\psi}(x) i \gamma^5 \psi(x) \varphi(x) \right], \quad (1.27)
\end{aligned}$$

where $\sigma(x)$ and $\varphi(x)$ are external sources of scalar $\bar{\psi}(x)\psi(x)$ and pseudoscalar $\bar{\psi}(x)i\gamma^5\psi(x)$ fermion densities, respectively.

Recall that Klaiber's expressions for $2n$ -point correlation functions of massless fermion fields have been used by Coleman [18] for his proof of equivalence between the massive Thirring model and the sine-Gordon model which are described by the Lagrangians [18]

$$\mathcal{L}_{\text{Th}}(x) = \bar{\psi}(x) (i \gamma^\mu \partial_\mu - m) \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \quad (1.28)$$

and

$$\mathcal{L}_{\text{SG}}(x) = \frac{1}{2} \partial_\mu \vartheta(x) \partial^\mu \vartheta(x) + \frac{\bar{\alpha}}{\beta^2} (\cos \bar{\beta} \vartheta(x) - 1), \quad (1.29)$$

where m is the mass of the Thirring fermion fields, $\vartheta(x)$ is a sine-Gordon model pseudoscalar (or scalar) field, and $\bar{\alpha}$ and $\bar{\beta}$ are real positive parameters [18].

The most interesting property of the sine-Gordon model is the existence of classical, stable solutions of the equations of motion – solitons and anti-solitons. Many-soliton solutions obey Pauli's exclusion principle and can be interpreted as a fermion-like behaviour.

The equivalence between the massive Thirring model and the sine-Gordon model Coleman proved perturbatively developing a perturbation theory with respect to the parameters m and α and quantizing fermion and boson field relative to the perturbative chiral symmetric vacuum. He showed that n -point correlation functions of the products of massless Thirring fermion field densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ and massless pseudoscalar boson sine-Gordon fields coincide if (i) the coupling constants g and β are related by

$$\frac{4\pi^2}{\beta^2} = 1 + \frac{g}{\pi} \quad (1.30)$$

and (ii) there exist the bosonization rules (so-called Abelian bosonization rules)

$$\begin{aligned}\bar{\psi}(x)\gamma^\mu\psi(x) &= \frac{\beta}{2\pi}\varepsilon^{\mu\nu}\partial_\nu\vartheta(x), \\ Z\bar{\psi}(x)\left(\frac{1\pm\gamma^5}{2}\right)\psi(x) &= -\frac{\alpha}{\beta^2}e^{\mp i\beta\vartheta(x)},\end{aligned}\tag{1.31}$$

where Z is a constant depending on the regularization procedure [18].

The generating functional (1.27) reduces itself to the path integral

$$\begin{aligned}Z_{\text{Th}}[\sigma, \varphi] &= Z_{\text{Th}}[\sigma_-, \sigma_+] = \\ &= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\xi \exp i \int d^2x \left\{ \bar{\Psi}(x) i\gamma^\mu \partial_\mu \Psi(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\ &\quad + \bar{\Psi}(x) \left(\frac{1+\gamma^5}{2} \right) e^{+2i\lambda(\alpha_J)\xi(x)} \Psi(x) \sigma_-(x) \\ &\quad \left. + \bar{\Psi}(x) \left(\frac{1-\gamma^5}{2} \right) e^{-2i\lambda(\alpha_J)\xi(x)} \Psi(x) \sigma_+(x) \right\},\end{aligned}\tag{1.32}$$

where $\sigma_\mp(x) = \sigma(x) \mp i\varphi(x)$ are external sources of the lefthanded and righthanded fermionic field density, respectively.

The evaluation of correlation functions of Thirring fermions by means of the generating functionals evidences an exact solution of the massless Thirring model.

In this paper we discuss the problems of the uniqueness of the solution of the massless Thirring model with fermion fields quantized in the chiral symmetric phase in terms of correlation functions. We analyse the evaluation of n -point correlation functions of left(right)handed fermion densities $\bar{\psi}(x)(1\pm\gamma^5)\psi(x)$ and the $2n$ -point Green functions and the dependence of these quantities on arbitrary parameters. In section 2 we discuss the operator formalism suggested by Klaiber for the solution of the massless Thirring model. We notice that Klaiber's operator approach to the solution of the massless Thirring model possesses some shortcomings. One of them concerns the non-standard parity properties. In order to show that the fermion vector current is proportional to the gradient of a scalar density Klaiber used the solution for a free massless fermionic field contradicting to the standard properties under parity transformation. This has led to composite field operators $c(k^1)$ and $c^\dagger(k^1)$ which do not have definite parity when compared with the standard definition [2,3]. In section 3 we give a representation of the vector current of the massless Thirring model using the solution of a free massless fermion field having standard parity properties. In section 4 we discuss canonical properties of the composite operators $c(k^1)$ and $c^\dagger(k^1)$ and construct new canonical composite operators $C(k^1)$ and $C^\dagger(k^1)$ behaving as scalars under parity transformations. We show that the transition to these new operators $C(k^1)$ and $C^\dagger(k^1)$ does not affect Klaiber's result for the evaluation of correlation functions and the statement on the existence of a one-parameter family of solutions of the massless Thirring model. Canonical properties of the operators $c(k^1)$ and $c^\dagger(k^1)$ and $C(k^1)$ and $C^\dagger(k^1)$ are discussed in connection with the derivation of the Schwinger terms. In section 5 within Klaiber's operator formalism we consider quantization of the massless Thirring model in the chiral symmetric phase. We derive the quantum equation of motion for the massless Thirring model and confirm the existence of a one-parameter

family of solutions for quantized fermionic fields. We show that Klaiber's parameterization of a one-parameter family of solutions is incorrect. We demonstrate that Klaiber's parameterization is based on the definition of the quantum vector current of the massless Thirring model breaking chiral symmetry explicitly. In fact, the divergence of the quantum axial-vector current $j_5^\mu(x)$ determined according to Klaiber's definition of the quantum vector current $j^\mu(x)$ does not vanish due to the contribution of non-covariant terms. We show that if one considers simultaneously a definition of $j_5^\mu(x)$ and requires the fulfillment of the standard relation $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$ all parameters become fixed and there is no one-parameter family of solutions of the massless Thirring model within Klaiber's operator formalism. We give (i) a definition of $j^\mu(x)$ and $j_5^\mu(x)$ consistent with chiral symmetry and the standard relation, $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$, definitions of quantum vector and axial-vector currents, (ii) a new definition of Klaiber's parameters and discuss (iii) a possibility to retain a one-parameter family of solutions of the massless Thirring model with quantized fermionic fields. In section 6 we replicate Naón's results for the n -point correlation functions of the left(right)handed fermion densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ which have been treated by Coleman [18] in connection with his proof of equivalence between the massive Thirring model and the sine-Gordon model quantizing the Thirring fermion fields in the chiral symmetric phase. We show that the dynamical dimensions of $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ depend on an arbitrary parameter α_J reflecting the ambiguities of the evaluation of chiral Jacobian by different regularization procedures. In section 7 we perform a complete bosonization of the generating functionals (1.32) and (1.26). We show that the generating functional (1.26) in the bosonized form coincides with the partition function of the sine-Gordon model at $\sigma(x) = -m$, where m can be treated as a Thirring fermion mass. In section 8 we evaluate the two-point fermion Green function by using the generating functional (1.23). We find again a dependence of the dynamical dimension of the massless fermionic field on the parameter α_J . The dependence of dynamical dimensions of correlation function on α_J agrees with the existence of a one-parameter family of solutions of the massless Thirring model. In section 9 we evaluate $2n$ -point Green functions of massless fermion fields and compare the obtained results with Klaiber's [4]. We show that Klaiber's correlation functions are singular in the infrared limit. This implies the necessity of an infrared renormalization of Klaiber's correlation functions. In section 10 we consider two-point correlation functions of left(right)handed fermion densities. We treat this correlation function as the two-point limit of the four-point correlation function. The former is evaluated explicitly both with Klaiber's operator formalism and the path-integral approach. We argue the necessity of the non-perturbative infrared renormalization of Klaiber's expression to get the result depending on the ultra-violet cut-off as we have in the path-integral approach. In section 11 we investigate the non-perturbative renormalization of the massless Thirring model. We find that (i) the coupling constant g is unrenormalized giving a vanishing Gell-Mann-Low function $\beta(g) = 0$ and (ii) the requirement of renormalizability fixes the parameter α_J to the value $\alpha_J = -2\pi/g$. We notice that our result concerning the vanishing of the Gell-Mann-Low function or, correspondingly, the absence of a coupling constant renormalization is supported by Mueller and Trueman [19] and Gomes and Lowenstein [20] who observed the same result within the massive Thirring model. In the Conclusion we discuss the obtained results. In Appendix A we discuss the properties of fermion fields under parity transformations. In Appendix B we evaluate the chiral Jacobian induced by local fermionic field transformations (1.16)

and demonstrate, following Christos [11], the dependence of the chiral Jacobian on an arbitrary parameter α_J . In Appendix C we calculate the two-point correlation functions defining commutators of scalar and pseudoscalar fermionic field densities. In Appendix D we make a justification of Klaiber's ansatz for the definition of the quantum fermionic vector current $K^\mu(x)$ and define in analogous way the quantum fermionic axial-vector current $K_5^\mu(x)$. In Appendix E we calculate a Fourier transform of the two-point Green function of the massless Thirring fermion fields. In Appendix F we analyse the non-perturbative renormalizability of the sine-Gordon model and show that it is not an asymptotically free quantum field theory as well as the massless Thirring model.

2 Proof of Klaiber's basic assumption and unusual parity properties of fermion fields

In this section we discuss Klaiber's operator approach [4] to the solution of the massless Thirring model. We focus our discussion on parity properties of fermion fields. Klaiber's operator formalism is based on the assumption of a proportionality of the vector fermion current to the gradient of a *scalar* fermion density, the proof of which has been carried out by the employment of the solution of the Dirac equation for a free massless fermionic field with non-standard parity properties.

Klaiber's approach to the solution of the massless Thirring model starts with the analysis of the equation of motion for the massless Thirring fermion field. Therefore, following Klaiber we have to start with the classical equation of motion and conservation equations. Using the Lagrangian (1.1) the equation of motion reads

$$i\gamma^\mu \partial_\mu \psi(x) = g j^\mu(x) \gamma_\mu \psi(x), \quad (2.1)$$

where the vector current $j^\mu(x)$ as well as the axial vector current $j_5^\mu(x)$ are defined by (1.6) and (1.7).

As has been stated by Klaiber [4] the vector and axial-vector currents can be written in the form of gradients

$$j_\mu(x) = \frac{1}{\sqrt{\pi}} \partial_\mu j(x) \quad , \quad j_{5\mu}(x) = \frac{1}{\sqrt{\pi}} \partial_\mu j_5(x). \quad (2.2)$$

Due to the properties of the vector and axial-vector currents, $j_\mu(x)$ and $j_{5\mu}(x)$, under the parity transformation (1.8) the quantities $j(x)$ and $j_5(x)$ should be scalar and pseudoscalar densities [2,3]

$$\begin{aligned} \mathcal{P} j(x^0, x^1) \mathcal{P}^\dagger &= +j(x^0, -x^1), \\ \mathcal{P} j_5(x^0, x^1) \mathcal{P}^\dagger &= -j_5(x^0, -x^1). \end{aligned} \quad (2.3)$$

Assuming the relations (2.2) Klaiber suggested to introduce a new fermionic field $\Psi(x)$

$$\Psi(x) = e^{i g j(x)/\sqrt{\pi}} \psi(x) \quad (2.4)$$

and its currents

$$\begin{aligned} J^\mu(x) &= \bar{\Psi}(x) \gamma^\mu \Psi(x), \\ J_5^\mu(x) &= \bar{\Psi}(x) \gamma^\mu \gamma^5 \Psi(x), \end{aligned} \quad (2.5)$$

In the case of the validity of (2.2) one can easily show that the fermion field (2.4) and the currents (2.5)⁴ satisfy the relations [4]

$$\begin{aligned} i\gamma^\mu \partial_\mu \Psi(x) &= 0, \\ J^\mu(x) &= j^\mu(x), \\ J_5^\mu(x) &= j_5^\mu(x). \end{aligned} \quad (2.6)$$

This implies that the fermion field $\Psi(x)$ has the same properties under parity transformations as the field $\psi(x)$

$$\begin{aligned} \mathcal{P} \Psi(x^0, x^1) \mathcal{P}^\dagger &= \gamma^0 \Psi(x^0, -x^1), \\ \mathcal{P} J^\mu(x^0, x^1) \mathcal{P}^\dagger &= +J_\mu(x^0, -x^1), \\ \mathcal{P} J_5^\mu(x^0, x^1) \mathcal{P}^\dagger &= -J_{5\mu}(x^0, -x^1). \end{aligned} \quad (2.7)$$

Thus, the general solution of the equation of motion (2.1) can be written in the form

$$\psi(x) = e^{-i g J(x)/\sqrt{\pi}} \Psi(x), \quad (2.8)$$

where $\Psi(x)$ obeys the free field equation of motion and $J(x)$ is *its integrated current* [4].

Due to the relation $J_5^\mu(x) = -\varepsilon^{\mu\nu} J_\nu(x)$ Klaiber claimed that the solution (2.8) can be extended and represented in the more general form of a one-parameter family

$$\begin{aligned} \psi(x) &= e^{-i\alpha J(x) - i\beta \gamma^5 J_5(x)} \Psi(x), \\ \bar{\psi}(x) &= \bar{\Psi}(x) e^{i\alpha J(x) - i\beta \gamma^5 J_5(x)}, \end{aligned} \quad (2.9)$$

where the parameters α and β obey the constraint

$$\alpha - \beta = \frac{g}{\sqrt{\pi}}, \quad (2.10)$$

and the scalar and pseudoscalar densities $J(x)$ and $J_5(x)$ are related to the vector and axial-vector currents $J_\mu(x)$ and $J_{5\mu}(x)$ as

$$J_\mu(x) = \bar{\Psi}(x) \gamma_\mu \Psi(x) = \frac{1}{\sqrt{\pi}} \partial_\mu J(x) \quad , \quad J_{5\mu}(x) = \bar{\Psi}(x) \gamma_\mu \gamma^5 \Psi(x) = \frac{1}{\sqrt{\pi}} \partial_\mu J_5(x). \quad (2.11)$$

Under parity transformations the densities $J(x)$ and $J_5(x)$ behave like $j(x)$ and $j_5(x)$, respectively,

$$\begin{aligned} \mathcal{P} J(x^0, x^1) \mathcal{P}^\dagger &= +J(x^0, -x^1), \\ \mathcal{P} J_5(x^0, x^1) \mathcal{P}^\dagger &= -J_5(x^0, -x^1). \end{aligned} \quad (2.12)$$

In quantizing the model Klaiber assumed that $\Psi(x)$ is a canonical free fermion field. As a free canonical field a quantized $\Psi(x)$ should have the following expansion into plane

⁴In the case of quantized fermionic fields the vector and axial-vector currents should be taken in the normal-ordered form $J^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \Psi(x) :$ and $J_5^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \gamma^5 \Psi(x) :$ as well as the equation of motion (2.1): $i\gamma^\mu \partial_\mu \psi(x) = g : j^\mu(x) \gamma_\mu \psi(x) :$.

waves [6]

$$\begin{aligned}\Psi(x) &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} \left[u(p^0, p^1) a(p^1) e^{-ip \cdot x} + v(p^0, p^1) b^\dagger(p^1) e^{ip \cdot x} \right], \\ \bar{\Psi}(x) &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} \left[\bar{u}(p^0, p^1) a^\dagger(p^1) e^{ip \cdot x} + \bar{v}(p^0, p^1) b(p^1) e^{-ip \cdot x} \right],\end{aligned}\quad (2.13)$$

where $p \cdot x = p^0 x^0 - p^1 x^1$. The creation $a^\dagger(p^1)$ ($b^\dagger(p^1)$) and annihilation $a(p^1)$ ($b(p^1)$) operators of fermions (antifermions) with momentum p^1 and energy $p^0 = |p^1|$ obey the anticommutation relations

$$\begin{aligned}\{a(p^1), a^\dagger(q^1)\} &= \{b(p^1), b^\dagger(q^1)\} = \delta(p^1 - q^1), \\ \{a(p^1), a(q^1)\} = \{a^\dagger(p^1), a^\dagger(q^1)\} &= \{b(p^1), b(q^1)\} = \{b^\dagger(p^1), b^\dagger(q^1)\} = 0.\end{aligned}\quad (2.14)$$

The spinorial wave functions $u(p^0, p^1)$ and $v(p^0, p^1) = u(-p^0, -p^1)$ are the solutions of the Dirac equation in the momentum representation for positive and negative energies, respectively. They are defined by [6]

$$\begin{aligned}u(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ \sqrt{p^0 - p^1} \end{pmatrix} = \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ \theta(-p^1) \end{pmatrix}, \\ \bar{u}(p^0, p^1) &= (\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) = \sqrt{2p^0} (\theta(-p^1), \theta(+p^1)), \\ v(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ -\sqrt{p^0 - p^1} \end{pmatrix} = \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ -\theta(-p^1) \end{pmatrix}, \\ \bar{v}(p^0, p^1) &= (-\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) = \sqrt{2p^0} (-\theta(-p^1), \theta(+p^1))\end{aligned}\quad (2.15)$$

at $p^0 = |p^1|$ and normalized to

$$\begin{aligned}u^\dagger(p^0, p^1) u(p^0, p^1) &= v^\dagger(p^0, p^1) v(p^0, p^1) = 2p^0, \\ \bar{u}(p^0, p^1) u(p^0, p^1) &= \bar{v}(p^0, p^1) v(p^0, p^1) = 0, \\ \bar{u}(p^0, p^1) v(p^0, p^1) &= \bar{v}(p^0, p^1) u(p^0, p^1) = 0.\end{aligned}\quad (2.16)$$

The spinorial wave functions $u(p^0, p^1)$ and $v(p^0, p^1)$ satisfy the following matrix relations [6]

$$u(p^0, p^1) \bar{u}(p^0, p^1) = v(p^0, p^1) \bar{v}(p^0, p^1) = \gamma^0 p^0 - \gamma^1 p^1 = \hat{p} \quad (2.17)$$

and the identities [2,3] (see Appendix A)

$$u(p^0, p^1) = \gamma^0 u(p^0, -p^1) \quad , \quad v(p^0, p^1) = -\gamma^0 v(p^0, -p^1). \quad (2.18)$$

These identities are important for the correct behaviour of the operators of creation and annihilation of fermions and antifermions under parity transformation (see Appendix A)

$$\begin{aligned}\mathcal{P} a(k^1) \mathcal{P}^\dagger &= +a(-k^1) \quad , \quad \mathcal{P} a^\dagger(k^1) \mathcal{P}^\dagger = +a^\dagger(-k^1), \\ \mathcal{P} b(k^1) \mathcal{P}^\dagger &= -b(-k^1) \quad , \quad \mathcal{P} b^\dagger(k^1) \mathcal{P}^\dagger = -b^\dagger(-k^1).\end{aligned}\quad (2.19)$$

We would like to emphasize that Klaiber [4] following Thirring used the expansion of the Ψ -field into plane wave, where

$$v(p^0, p^1) = u(p^0, p^1) = \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ \theta(-p^1) \end{pmatrix}. \quad (2.20)$$

For such a solution all relations (2.16) and (2.17) are retained. However, as has been shown in [6] this solution cannot be obtained from the solution for a free massive fermionic field in the zero-mass limit. As a result the spinorial wave function $v(p^0, p^1)$ breaks the identity (2.18) that entails the violation of the relations (2.19). In fact, according to the solution (2.20) they read (see Appendix A)

$$\begin{aligned} \mathcal{P} a(k^1) \mathcal{P}^\dagger &= +a(-k^1) \quad , \quad \mathcal{P} a^\dagger(k^1) \mathcal{P}^\dagger = +a^\dagger(-k^1) \\ \mathcal{P} b(k^1) \mathcal{P}^\dagger &= \overset{?}{+} b(-k^1) \quad , \quad \mathcal{P} b^\dagger(k^1) \mathcal{P}^\dagger = \overset{?}{+} b^\dagger(-k^1). \end{aligned} \quad (2.21)$$

This contradicts to the well-known properties of the fermion creation and annihilation operators under parity transformations.

Now let us replicate the derivation of the representation of the vector current $J^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \Psi(x)$: suggested by Klaiber [4]

$$J^\mu(x) = -\frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} k^\mu \left[c(k^1) e^{-ik \cdot x} - c^\dagger(k^1) e^{ik \cdot x} \right], \quad (2.22)$$

and confirming Klaiber's assumption (2.11).

Following Klaiber and using the expansion of the Ψ -field into plane waves with the spinorial functions related by (2.20) we obtain

$$\begin{aligned} J^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \Psi(x) &:= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dp^1 dq^1}{2\pi} \frac{1}{\sqrt{2p^0 2q^0}} \bar{u}(p^0, p^1) \gamma^\mu u(q^0, q^1) \\ &\times \left\{ a^\dagger(p^1) a(q^1) e^{i(p-q) \cdot x} - b^\dagger(q^1) b(p^1) e^{-i(p-q) \cdot x} \right. \\ &\left. + b(p^1) a(q^1) e^{-i(p+q) \cdot x} + a^\dagger(p^1) b^\dagger(q^1) e^{i(p+q) \cdot x} \right\}. \end{aligned} \quad (2.23)$$

The analytical expression of the quantity $[\bar{u}(p^0, p^1) \gamma^\mu u(q^0, q^1)]$ suitable for the derivation of the momentum representation is rather ambiguous and the result depends fully on the skill and intuition of the investigator. Following Klaiber's intuition we represent this quantity as follows

$$\begin{aligned} \bar{u}(p^0, p^1) \gamma^\mu u(q^0, q^1) &= \\ &= \frac{p^\mu - q^\mu}{p^0 - q^0} \sqrt{2p^0 2q^0} \theta(q^1(p^1 - q^1)) + \frac{q^\mu - p^\mu}{q^0 - p^0} \sqrt{2p^0 2q^0} \theta(p^1(q^1 - p^1)) \end{aligned} \quad (2.24)$$

for the first and second term in (2.23) and

$$\bar{u}(p^0, p^1) \gamma^\mu u(q^0, q^1) = \sqrt{2p^0 2q^0} \frac{p^\mu + q^\mu}{p^0 + q^0} \theta(p^1 q^1) \quad (2.25)$$

for the third and fourth term, respectively.

Using (2.24) and (2.25) we represent (2.23) in the form

$$\begin{aligned}
J^\mu(x) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dp^1 dq^1}{2\pi} \\
& \times \left\{ \left[\frac{p^\mu - q^\mu}{p^0 - q^0} \theta(q^1(p^1 - q^1)) + \frac{q^\mu - p^\mu}{q^0 - p^0} \theta(p^1(q^1 - p^1)) \right] a^\dagger(p^1) a(q^1) e^{+i(p-q) \cdot x} \right. \\
& - \left[\frac{p^\mu - q^\mu}{p^0 - q^0} \theta(q^1(p^1 - q^1)) + \frac{q^\mu - p^\mu}{q^0 - p^0} \theta(p^1(q^1 - p^1)) \right] b^\dagger(q^1) b(p^1) e^{-i(p-q) \cdot x} \\
& + \frac{p^\mu + q^\mu}{p^0 + q^0} \theta(p^1 q^1) b(p^1) a(q^1) e^{-i(p+q) \cdot x} \\
& \left. + \frac{p^\mu + q^\mu}{p^0 + q^0} \theta(p^1 q^1) a^\dagger(p^1) b^\dagger(q^1) e^{+i(p+q) \cdot x} \right\}. \tag{2.26}
\end{aligned}$$

Then, let us rewrite (2.26) in an equivalent form and consider the components of the vector current separately

$$\begin{aligned}
J^0(x) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dp^1 dq^1}{2\pi} \\
& \times \left\{ \theta(q^1(p^1 - q^1)) [a^\dagger(q^1) a(p^1) - b^\dagger(q^1) b(p^1)] e^{-i(p-q) \cdot x} \right. \\
& + \theta(q^1(p^1 - q^1)) [a^\dagger(p^1) a(q^1) - b^\dagger(p^1) b(q^1)] e^{+i(p-q) \cdot x} \\
& + \theta(p^1 q^1) b(p^1) a(q^1) e^{-i(p+q) \cdot x} \\
& \left. + \theta(p^1 q^1) a^\dagger(p^1) b^\dagger(q^1) e^{+i(p+q) \cdot x} \right\}, \\
J^1(x) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dp^1 dq^1}{2\pi} \\
& \times \left\{ \varepsilon(p^1 - q^1) \theta(q^1(p^1 - q^1)) [a^\dagger(q^1) a(p^1) - b^\dagger(q^1) b(p^1)] e^{-i(p-q) \cdot x} \right. \\
& + \varepsilon(p^1 - q^1) \theta(q^1(p^1 - q^1)) [a^\dagger(p^1) a(q^1) - b^\dagger(p^1) b(q^1)] e^{+i(p-q) \cdot x} \\
& + \varepsilon(p^1 + q^1) \theta(p^1 q^1) b(p^1) a(q^1) e^{-i(p+q) \cdot x} \\
& \left. + \varepsilon(p^1 + q^1) \theta(p^1 q^1) a^\dagger(p^1) b^\dagger(q^1) e^{+i(p+q) \cdot x} \right\}, \tag{2.27}
\end{aligned}$$

where $\varepsilon(p^1 - q^1)$ is a sign function.

Making a change of variables $p^1 - q^1 \rightarrow k^1$ in the first and second terms and $p^1 + q^1 \rightarrow k^1$ in the third and fourth ones we arrive at the expression

$$\begin{aligned}
J^0(x) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk^1 dq^1}{2\pi} \\
& \times \left\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)] e^{-i(|k^1 + q^1| - |q^1|)x^0 + ik^1 x^1} \right. \\
& \left. + \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)] e^{+i(|k^1 + q^1| - |q^1|)x^0 - ik^1 x^1} \right\}
\end{aligned}$$

$$\begin{aligned}
& + \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) e^{-i(|k^1 - q^1| + |q^1|)x^0 + ik^1x^1} \\
& + \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) e^{+i(|k^1 - q^1| + |q^1|)x^0 - ik^1x^1} \Big\}, \\
J^1(x) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk^1 dq^1}{2\pi} \\
& \times \Big\{ \varepsilon(k^1) \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)] e^{-i(|k^1 + q^1| - |q^1|)x^0 + ik^1x^1} \\
& + \varepsilon(k^1) \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)] e^{+i(|k^1 + q^1| - |q^1|)x^0 - ik^1x^1} \\
& + \varepsilon(k^1) \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) e^{-i(|k^1 - q^1| + |q^1|)x^0 + ik^1x^1} \\
& + \varepsilon(k^1) \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) e^{+i(|k^1 - q^1| + |q^1|)x^0 - ik^1x^1} \Big\}. \tag{2.28}
\end{aligned}$$

The relations (2.28) can be represented in the compact form

$$\begin{aligned}
J^\mu(x) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk^1 dq^1}{2\pi} \frac{k^\mu}{k^0} \\
& \times \Big\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)] e^{-i(|k^1 + q^1| - |q^1|)x^0 + ik^1x^1} \\
& + \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)] e^{+i(|k^1 + q^1| - |q^1|)x^0 - ik^1x^1} \\
& + \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) e^{-i(|k^1 - q^1| + |q^1|)x^0 + ik^1x^1} \\
& + \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) e^{+i(|k^1 - q^1| + |q^1|)x^0 - ik^1x^1} \Big\}. \tag{2.29}
\end{aligned}$$

Due to the Heaviside functions in the meaningful region of momentum integrals we have $|k^1 + q^1| - |q^1| = |k^1| = k^0$ and $|k^1 - q^1| - |q^1| = |k^1| = k^0$. This yields

$$\begin{aligned}
J^\mu(x) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk^1 dq^1}{2\pi} \frac{k^\mu}{k^0} \\
& \times \Big\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)] e^{-ik \cdot x} \\
& + \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)] e^{+ik \cdot x} \\
& + \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) e^{-ik \cdot x} \\
& + \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) e^{+ik \cdot x} \Big\} = \\
& = -\frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} k^\mu \left[c(k^1) e^{-ik \cdot x} - c^\dagger(k^1) e^{ik \cdot x} \right]. \tag{2.30}
\end{aligned}$$

Thus, the vector current can be represented in the form (2.22), where the operators $c(k^1)$ and $c^\dagger(k^1)$ are defined by

$$c(k^1) = \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)] \right.$$

$$\begin{aligned}
& +\theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) \Big\}, \\
c^\dagger(k^1) &= -\frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \Big\{ \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)] \\
& +\theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) \Big\}. \tag{2.31}
\end{aligned}$$

These operators agree fully with the expressions adduced by Klaiber (see (II.15) of Ref.[4]).

The vector current given by (2.28) satisfies the relation (2.11) with the fermion density $J(x)$ equal to

$$J(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[c(k^1) e^{-ik \cdot x} + c^\dagger(k^1) e^{ik \cdot x} \right]. \tag{2.32}$$

The axial vector current $J_5^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \gamma^5 \Psi(x) :$ and the fermion density $J_5(x)$ related by $J_5^\mu(x) = \partial^\mu J_5(x) / \sqrt{\pi}$ have the following momentum representations

$$\begin{aligned}
J_5^\mu(x) &= -\frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \varepsilon(k^1) k^\mu \left[c(k^1) e^{-ik \cdot x} - c^\dagger(k^1) e^{ik \cdot x} \right], \\
J_5(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \varepsilon(k^1) \left[c(k^1) e^{-ik \cdot x} + c^\dagger(k^1) e^{ik \cdot x} \right], \tag{2.33}
\end{aligned}$$

where $\varepsilon(k^1)$ is a sign function.

According to Klaiber's analysis the operators $c(k^1)$ and $c^\dagger(k^1)$ should be scalar operators. This implies that under parity transformations these operators should transform like

$$\begin{aligned}
\mathcal{P} c(k^1) \mathcal{P}^\dagger &= c(-k^1), \\
\mathcal{P} c^\dagger(k^1) \mathcal{P}^\dagger &= c^\dagger(-k^1). \tag{2.34}
\end{aligned}$$

Using the standard properties (2.19) of the creation and annihilation operators of fermions and antifermions under parity transformation one finds

$$\begin{aligned}
\mathcal{P} c(k^1) \mathcal{P}^\dagger &= \\
&= \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \Big\{ \theta(k^1 q^1) [a^\dagger(-q^1) a(-k^1 - q^1) - b^\dagger(-q^1) b(-k^1 - q^1)] \\
&\quad - \theta(q^1(k^1 - q^1)) b(-k^1 + q^1) a(-q^1) \Big\} = \\
&= \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \Big\{ \theta(-k^1 q^1) [a^\dagger(q^1) a(-k^1 + q^1) - b^\dagger(q^1) b(-k^1 + q^1)] \\
&\quad - \theta(q^1(-k^1 - q^1)) b(-k^1 - q^1) a(q^1) \Big\} \neq c(-k^1), \\
\mathcal{P} c^\dagger(k^1) \mathcal{P}^\dagger &\neq c^\dagger(-k^1). \tag{2.35}
\end{aligned}$$

Hence, within the standard definition of the properties of fermion and antifermion operators of creation and annihilation, the operators $J^\mu(x)$ and $J_5^\mu(x)$ given by (2.22) and (2.33) do not possess a definite parity as well as the fermion densities $J(x)$ and $J_5(x)$.

3 Fermionic fields with standard properties under parity transformation and vector current decomposition

In this section we derive the decomposition of the vector current operator $J^\mu(x) =: \bar{\Psi}(x)\gamma^\mu\Psi(x)$ into scalar and pseudoscalar fermion densities. We use the solution for a free massless fermionic field (2.15) with standard properties of creation and annihilation operators under parity transformations [2,3] (see Appendix A). In this case (2.23) reads

$$\begin{aligned} J^\mu(x) =: \bar{\Psi}(x)\gamma^\mu\Psi(x) := & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dp^1 dq^1}{2\pi} \frac{1}{\sqrt{2p^0 2q^0}} \left\{ [\bar{u}(p^0, p^1)\gamma^\mu u(q^0, q^1)] \right. \\ & \times a^\dagger(p^1)a(q^1)e^{i(p-q)\cdot x} - [\bar{v}(p^0, p^1)\gamma^\mu v(q^0, q^1)] b^\dagger(q^1)b(p^1)e^{-i(p-q)\cdot x} \\ & + [\bar{v}(p^0, p^1)\gamma^\mu u(q^0, q^1)] b(p^1)a(q^1)e^{-i(p+q)\cdot x} \\ & \left. + [\bar{u}(p^0, p^1)\gamma^\mu v(q^0, q^1)] a^\dagger(p^1)b^\dagger(q^1)e^{i(p+q)\cdot x} \right\}. \end{aligned} \quad (3.1)$$

The products of spinorial wave functions are equal to

$$\begin{aligned} \bar{u}(p^0, p^1)\gamma^\mu u(q^0, q^1) &= [\bar{v}(p^0, p^1)\gamma^\mu v(q^0, q^1)] = \\ &= \frac{p^\mu - q^\mu}{p^0 - q^0} \sqrt{2p^0 2q^0} \theta(q^1(p^1 - q^1)) + \frac{q^\mu - p^\mu}{q^0 - p^0} \sqrt{2p^0 2q^0} \theta(p^1(q^1 - p^1)), \\ [\bar{v}(p^0, p^1)\gamma^\mu u(q^0, q^1)] &= [\bar{v}(p^0, p^1)\gamma^\mu u(q^0, q^1)] = -\sqrt{2p^0 2q^0} \varepsilon^{\mu\nu} \frac{p_\nu + q_\nu}{p^0 + q^0} \theta(p^1 q^1). \end{aligned} \quad (3.2)$$

Using these expressions we recast the r.h.s. of (3.1) into the form

$$\begin{aligned} J^\mu(x) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk^1 dq^1}{2\pi} \frac{k_\nu}{k^0} \\ &\times \left\{ g^{\mu\nu} \theta(k^1 q^1) [a^\dagger(q^1)a(k^1 + q^1) - b^\dagger(q^1)b(k^1 + q^1)] e^{-ik\cdot x} \right. \\ &- \varepsilon^{\mu\nu} \theta(q^1(k^1 - q^1)) b(k^1 - q^1)a(q^1) e^{-ik\cdot x} \\ &+ g^{\mu\nu} \theta(k^1 q^1) [a^\dagger(k^1 + q^1)a(q^1) - b^\dagger(k^1 + q^1)b(q^1)] e^{ik\cdot x} \\ &- \varepsilon^{\mu\nu} \theta(q^1(k^1 - q^1)) a^\dagger(q^1)b^\dagger(k^1 - q^1) e^{ik\cdot x} \left. \right\} = \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[-ik^\mu d(k^1) e^{-ik\cdot x} - ik_\nu \varepsilon^{\mu\nu} d_5(k^1) e^{-ik\cdot x} \right. \\ &\left. + ik^\mu d^\dagger(k^1) e^{ik\cdot x} + ik_\nu \varepsilon^{\mu\nu} d_5^\dagger(k^1) e^{ik\cdot x} \right]. \end{aligned} \quad (3.3)$$

This implies that the vector-current $J^\mu(x)$ has the following decomposition

$$J^\mu(x) = \frac{1}{\sqrt{\pi}} \partial^\mu J(x) + \frac{1}{\sqrt{\pi}} \varepsilon^{\mu\nu} \partial_\nu J_5(x) \quad (3.4)$$

in terms of gradients of the scalar and pseudoscalar fermion densities $J(x)$ and $J_5(x)$ defined by

$$J(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[d(k^1) e^{-ik\cdot x} + d^\dagger(k^1) e^{ik\cdot x} \right],$$

$$J_5(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[d_5(k^1) e^{-ik \cdot x} + d_5^\dagger(k^1) e^{ik \cdot x} \right]. \quad (3.5)$$

The operators $d(k^1)$, $d^\dagger(k^1)$, $d_5(k^1)$ and $d_5^\dagger(k^1)$ read

$$\begin{aligned} d(k^1) &= \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1)], \\ d^\dagger(k^1) &= -\frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - b^\dagger(k^1 + q^1) b(q^1)], \\ d_5(k^1) &= -\frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1), \\ d_5^\dagger(k^1) &= \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1). \end{aligned} \quad (3.6)$$

They possess correct properties under parity transformation

$$\begin{aligned} \mathcal{P} d(k^1) \mathcal{P}^\dagger &= +d(-k^1), \\ \mathcal{P} d_5(k^1) \mathcal{P}^\dagger &= -d_5(-k^1). \end{aligned} \quad (3.7)$$

Since $J^\mu(x) = j^\mu(x)$, the same decomposition is valid for $j^\mu(x)$, i.e.

$$j^\mu(x) = \frac{1}{\sqrt{\pi}} \partial^\mu j(x) + \frac{1}{\sqrt{\pi}} \varepsilon^{\mu\nu} \partial_\nu j_5(x). \quad (3.8)$$

Thus, the vector currents $J^\mu(x)$ and $j^\mu(x)$ have standard Hodge decompositions.

4 Canonical commutation relation for $c(k^1)$ and $c^\dagger(k^1)$, the Schwinger term and a one-parameter family of solutions

As has been shown by Klaiber the operators $c(k^1)$ and $c^\dagger(p^1)$ obey the canonical commutation relation

$$[c(k^1), c^\dagger(p^1)] = \delta(k^1 - p^1). \quad (4.1)$$

This is very important and amazing result of Klaiber's operator formalism, if to take into account that the commutation relation (4.1) can be obtained by using canonical anti-commutation relations for operators of creation and annihilation of fermions and antifermions (2.14).

Recall, as has been proved by Callan, Dashen, and Sharp [21] the commutation relation between time and spatial components of the vector current $j^\mu(x) = \bar{\psi}(x) \gamma^\mu \psi(x)$ vanishes

$$[j^0(x^0, x^1), j^1(x^0, y^1)] = 0 \quad (4.2)$$

using canonical equal-time anticommutation relations for the massless Thirring fermion fields

$$\{\psi(x^0, x^1), \psi^\dagger(x^0, y^1)\} = \delta(x^1 - y^1). \quad (4.3)$$

However, according to Schwinger [22] the equal-time commutator (4.2) should not vanish and reads

$$[j^0(x^0, x^1), j^1(x^0, y^1)] = -i c \frac{\partial}{\partial x^1} \delta(x^1 - y^1), \quad (4.4)$$

where c is the Schwinger term. In the massless Thirring model for fermion fields quantized in the chiral symmetric phase the Schwinger term has been calculated by Sommerfield [23]: $c = 1/\pi$. We would like to emphasize, that the result $c = 1/\pi$ is not a consequence of canonical anti-commutativity of fermionic field operators but the result of an evaluation of vacuum expectation values (see also [6]).

The calculation of the equal-time commutator $[j^0(x^0, x^1), j^1(x^0, y^1)]$ within Klaiber's formalism runs in the way

$$\begin{aligned} [j^0(x^0, x^1), j^1(x^0, y^1)] &= [J^0(x^0, x^1), J^1(x^0, y^1)] = -\frac{1}{\sqrt{\pi}} \frac{\partial}{\partial y^1} [J^0(x^0, x^1), J(x^0, y^1)] = \\ &= \frac{\partial}{\partial y^1} \frac{i}{2\pi^2} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \int_{-\infty}^{\infty} \frac{dq^1}{\sqrt{2q^0}} k^0 \left\{ [c(k^1), c^\dagger(q^1)] e^{-i(k^0 - q^0)x^0 + ik^1x^1 - iq^1y^1} \right. \\ &\times \left. + [c(q^1), c^\dagger(k^1)] e^{i(k^0 - q^0)x^0 - ik^1x^1 + iq^1y^1} \right\} = i \frac{1}{\pi} \frac{\partial}{\partial y^1} \int_{-\infty}^{\infty} \frac{dk^1}{2\pi} e^{ik^1(x^1 - y^1)} = \\ &= -i \frac{1}{\pi} \frac{\partial}{\partial x^1} \delta(x^1 - y^1). \end{aligned} \quad (4.5)$$

The obtained result agrees completely with (4.4) but contradicts to (4.2). This should be much more astonishing due to the fact that $J^0(x^0, x^1)$ and $J^1(x^0, y^1)$ are the components of the vector current of a free massless fermion field. The derivation of the equal-time commutator (4.5) in agreement with Schwinger's ansatz (4.4) testifies that the operators $c(k^1)$ and $c^\dagger(k^1)$ are not real canonical quantum field operators. We prove this statement in section 5.

Now let us calculate the commutator $[c(k^1), c^\dagger(p^1)]$. Substituting (2.31) and using the relation

$$[AB, CD] = A\{B, C\}D + CA\{B, D\} - C\{A, D\}B - \{A, C\}BD, \quad (4.6)$$

where A, B, C and D are any operators, we get

$$\begin{aligned} [c(k^1), c^\dagger(p^1)] &= \frac{1}{\sqrt{k^0 p^0}} \int_{-\infty}^{\infty} dq^1 \int_{-\infty}^{\infty} d\ell^1 \left\{ \theta(k^1 q^1) \theta(p^1 \ell^1) \right. \\ &\times \left([a^\dagger(q^1) a(k^1 + q^1), a^\dagger(p^1 + \ell^1) a(\ell^1)] + [b^\dagger(q^1) b(k^1 + q^1), b^\dagger(p^1 + \ell^1) b(\ell^1)] \right) \\ &+ \theta(q^1(k^1 - q^1)) \theta(\ell^1(p^1 - \ell^1)) [b(k^1 - q^1) a(q^1), a^\dagger(\ell^1) b^\dagger(p^1 - \ell^1)] \\ &+ \theta(k^1 q^1) \theta(\ell^1(p^1 - \ell^1)) [a^\dagger(q^1) a(k^1 + q^1) - b^\dagger(q^1) b(k^1 + q^1), a^\dagger(\ell^1) b^\dagger(p^1 - \ell^1)] \\ &\left. + \theta(q^1(k^1 - q^1)) \theta(p^1 \ell^1) [b(k^1 - q^1) a(q^1), a^\dagger(p^1 + \ell^1) a(\ell^1) - b^\dagger(p^1 + \ell^1) b(\ell^1)] \right\} = \\ &= \frac{1}{\sqrt{k^0 p^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \delta(k^1 - p^1) \theta(q^1(k^1 - q^1)) \theta(q^1(p^1 - q^1)) + [\theta(k^1 q^1) \theta(p^1(k^1 - p^1 + q^1)) \right. \\ &\left. - \theta(k^1(q^1 - p^1)) \theta(p^1(q^1 - p^1)) - \theta(q^1(p^1 - q^1)) \theta((p^1 - q^1)(k^1 - p^1 + q^1))] \right\} \end{aligned}$$

$$\begin{aligned} & \times [a^\dagger(q^1)a(k^1 - p^1 + q^1) + b^\dagger(q^1)b(k^1 - p^1 + q^1)] + [\theta(k^1 q^1)\theta((k^1 + q^1)(p^1 - k^1 - q^1)) \\ & - \theta(k^1(p^1 - k^1 - q^1))\theta(q^1(p^1 - q^1))] a^\dagger(q^1)b^\dagger(p^1 - k^1 - q^1) + [\theta(p^1(k^1 - p^1 - q^1)) \\ & \times \theta(q^1(k^1 - q^1)) - \theta(p^1 q^1)\theta((p^1 + q^1)(k^1 - p^1 - q^1))] b(q^1)a(k^1 - p^1 - q^1) \Big\} = \delta(k^1 - p^1). \end{aligned} \quad (4.7)$$

This confirms Klaiber's result (4.1).

However, in order to identify the operators $c(k^1)$ and $c^\dagger(k^1)$ with operators of annihilation and creation of a boson with a momentum k^1 it is necessary to show that the commutators $[c(k^1), c(p^1)]$ and $[c^\dagger(k^1), c^\dagger(p^1)]$ vanish. In this case $c(k^1)$ and $c^\dagger(k^1)$ would be canonical quantum field operators. The calculation of the commutator $[c(k^1), c(p^1)]$ is analogous to $[c(k^1), c^\dagger(p^1)]$ and reads

$$\begin{aligned} [c(k^1), c(p^1)] &= -\frac{1}{\sqrt{k^0 p^0}} \int_{-\infty}^{\infty} dq^1 \int_{-\infty}^{\infty} d\ell^1 \Big\{ \theta(k^1 q^1)\theta(p^1 \ell^1) \\ & \times \left([a^\dagger(q^1)a(k^1 + q^1), a^\dagger(\ell^1)a(p^1 + \ell^1)] + [b^\dagger(q^1)b(k^1 + q^1), b^\dagger(\ell^1)b(p^1 + \ell^1)] \right) \\ & + \theta(k^1 q^1)\theta(\ell^1(p^1 - \ell^1)) [a^\dagger(q^1)a(k^1 + q^1) - b^\dagger(q^1)b(k^1 + q^1), b(p^1 - \ell^1)a(\ell^1)] \\ & + \theta(q^1(k^1 - q^1))\theta(p^1 \ell^1) [b(k^1 - q^1)a(q^1), a^\dagger(\ell^1)a(p^1 + \ell^1) - b^\dagger(\ell^1)b(p^1 + \ell^1)] \Big\} \\ &= -\frac{1}{\sqrt{k^0 p^0}} \int_{-\infty}^{\infty} dq^1 \Big\{ [\theta(k^1 q^1)\theta(p^1(k^1 + q^1)) - \theta(p^1 q^1)\theta(k^1(p^1 + q^1))] \\ & \times [a^\dagger(q^1)a(k^1 + p^1 + q^1) + b^\dagger(q^1)b(k^1 + p^1 + q^1)] + [\theta(k^1(q^1 - k^1)) \\ & \times \theta((q^1 - k^1)(k^1 + p^1 - q^1)) - \theta(k^1(p^1 - q^1))\theta(q^1(p^1 - q^1)) + \theta(q^1(k^1 - q^1)) \\ & \times \theta(p^1(k^1 - q^1)) - \theta(p^1(q^1 - p^1))\theta((q^1 - p^1)(k^1 + p^1 - q^1))] b(q^1)a(k^1 + p^1 - q^1) \Big\}. \end{aligned} \quad (4.8)$$

Accounting for the properties of the Heaviside functions one can show that all momentum integrals vanish. This yields

$$[c(k^1), c(p^1)] = 0. \quad (4.9)$$

This testifies that from the naive point of view $c(k^1)$ and $c^\dagger(k^1)$ are canonical quantum field operators.

In turn, the operators $d(k^1)$, $d^\dagger(k^1)$, $d_5(k^1)$ and $d_5^\dagger(k^1)$ do not satisfy canonical commutation relations (4.1) and (4.9). Moreover, the commutators of these operators are q -valued quantities. Hence, the only way to deal with canonical quantum field operators is to follow Klaiber and represent the vector fermion current $J^\mu(x)$ in the form of a gradient of the scalar fermionic field density.

For this aim we suggest to turn to (3.3) and rewrite the product $\varepsilon^{\mu\nu} k_\nu$ in the equivalent form

$$\varepsilon^{\mu\nu} k_\nu = -\varepsilon(k^1) k^\mu. \quad (4.10)$$

This allows to recast the vector current into the form

$$J^\mu(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[-ik^\mu C(k^1) e^{-ik \cdot x} + ik^\mu C^\dagger(k^1) e^{ik \cdot x} \right] = \frac{1}{\sqrt{\pi}} \partial^\mu S(x), \quad (4.11)$$

where $S(x)$ is a real scalar fermionic field density

$$S(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \left[C(k^1) e^{-ik \cdot x} + C^\dagger(k^1) e^{ik \cdot x} \right]. \quad (4.12)$$

The operators $C(k^1)$ and $C^\dagger(k^1)$ are defined by

$$\begin{aligned} C(k^1) &= d(k^1) - \varepsilon(k^1) d_5(k^1) = \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) \right. \\ &\quad \left. - b^\dagger(q^1) b(k^1 + q^1)] + \varepsilon(k^1) \theta(q^1(k^1 - q^1)) b(k^1 - q^1) a(q^1) \right\}, \\ C^\dagger(k^1) &= d^\dagger(k^1) - \varepsilon(k^1) d_5^\dagger(k^1) = -\frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) \right. \\ &\quad \left. - b^\dagger(k^1 + q^1) b(q^1)] + \varepsilon(k^1) \theta(q^1(k^1 - q^1)) a^\dagger(q^1) b^\dagger(k^1 - q^1) \right\}. \end{aligned} \quad (4.13)$$

The operators $C(k^1)$ and $C^\dagger(k^1)$ have standard properties under parity transformations

$$\mathcal{P}C(k^1)\mathcal{P}^\dagger = +C(-k^1) \quad , \quad \mathcal{P}C^\dagger(k^1)\mathcal{P}^\dagger = +C^\dagger(-k^1). \quad (4.14)$$

The axial-vector current $J_5^\mu(x)$ can be written in complete analogy with the vector current

$$J_5^\mu(x) = \frac{1}{\sqrt{\pi}} \partial^\mu P(x), \quad (4.15)$$

where $P(x)$ is a pseudoscalar fermionic field density defined by

$$P(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \varepsilon(k^1) \left[C(k^1) e^{-ik \cdot x} + C^\dagger(k^1) e^{ik \cdot x} \right]. \quad (4.16)$$

Following the calculation of the commutators $[c(k^1), c^\dagger(p^1)]$ and $[c(k^1), c(p^1)]$ represented in (4.7) and (4.8) one can show that $C(k^1)$ and $C^\dagger(p^1)$ are canonical quantum field operators obeying the commutation relations

$$\begin{aligned} [C(k^1), C(p^1)] &= [C^\dagger(k^1), C^\dagger(p^1)] = 0, \\ [C(k^1), C^\dagger(p^1)] &= \delta(k^1 - p^1). \end{aligned} \quad (4.17)$$

A one-parameter family of solutions of the equation of motion of the massless Thirring model reads now

$$\begin{aligned} \psi(x) &= e^{-i\alpha S(x) - i\beta \gamma^5 P(x)} \Psi(x), \\ \bar{\psi}(x) &= \bar{\Psi}(x) e^{i\alpha S(x) - i\beta \gamma^5 P(x)}. \end{aligned} \quad (4.18)$$

Using the relations

$$\begin{aligned} J^\mu(x) &= \frac{1}{\sqrt{\pi}} \partial^\mu S(x) = j^\mu(x) = \frac{1}{\sqrt{\pi}} \partial^\mu s(x), \\ J_5^\mu(x) &= \frac{1}{\sqrt{\pi}} \partial^\mu P(x) = j_5^\mu(x) = \frac{1}{\sqrt{\pi}} \partial^\mu p(x) \end{aligned} \quad (4.19)$$

and $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$ one can easily show that the solution (4.18) obeys the equation of motion (2.1) for $\alpha - \beta = g/\sqrt{\pi}$. Hence, the equation of motion of the massless Thirring model possesses a one-parameter family of solutions as claimed by Klaiber [4].

We would like to accentuate that this statement is only obtained for the classical solutions of the classical equation of motion of the massless Thirring model. In the next section we will show that the quantum version of this hypothesis is not valid.

Now we would like to discuss the definition of annihilation operator of antifermions with momentum p^1 given in Ref.[16]. According to Eq.(2.38) of Ref.[16] in order to use Thirring's solution for a free massless fermion field (2.13) with spinorial wave functions (2.30) one can define the annihilation operators of antifermions as follows

$$b(p^1) \rightarrow \tilde{b}(p^1) = \begin{cases} +b(p^1) & , \quad p^1 > 0, \\ -b(p^1) & , \quad p^1 < 0. \end{cases} \quad (4.20)$$

Formally, this relation can be defined in terms the sign function $\varepsilon(p^1)$, i.e. $\tilde{b}(p^1) = \varepsilon(p^1) b(p^1)$. The operator $b(p^1)$ should possess standard properties under the parity transformation $\mathcal{P}b(p^1)\mathcal{P}^\dagger = -b(-p^1)$ (2.19). In this case the expansion into plane waves of a free massless fermion field $\Psi(x)$ should read

$$\begin{aligned} \Psi(x^0, x^1) &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} u(p^0, p^1) \left[a(p^1) e^{-ip^0 x^0 + ip^1 x^1} + \tilde{b}^\dagger(p^1) e^{ip^0 x^0 - ip^1 x^1} \right] = \\ &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} u(p^0, p^1) \left[a(p^1) e^{-ip^0 x^0 + ip^1 x^1} + \varepsilon(p^1) b^\dagger(p^1) e^{ip^0 x^0 - ip^1 x^1} \right]. \end{aligned} \quad (4.21)$$

Now let us make a parity transformation of the Ψ -field (4.21) by using standard properties of the operators $a(p^1)$ and $b^\dagger(p^1)$ (2.19) and the spinorial function $u(p^0, p^1)$ (2.18). As a result we get

$$\begin{aligned} \mathcal{P}\Psi(x^0, x^1)\mathcal{P}^\dagger &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} u(p^0, p^1) \left[\mathcal{P}a(p^1)\mathcal{P}^\dagger e^{-ip^0 x^0 + ip^1 x^1} + \varepsilon(p^1) \right. \\ &\times \mathcal{P}b^\dagger(p^1)\mathcal{P}^\dagger e^{ip^0 x^0 - ip^1 x^1} \left. \right] = \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} \gamma^0 u(p^0, -p^1) \left[a(-p^1) e^{-ip^0 x^0 + ip^1 x^1} \right. \\ &\left. - \varepsilon(p^1) b^\dagger(-p^1) e^{ip^0 x^0 - ip^1 x^1} \right] = \gamma^0 \Psi(x^0, -x^1), \end{aligned} \quad (4.22)$$

where we have made a change of variables $p^1 \rightarrow -p^1$ and used the relation $\varepsilon(-p^1) = -\varepsilon(p^1)$. This agrees with standard properties of a free fermion field under parity transformation [2,3] (see also Appendix A).

According to the prescription (4.20) the operators $c(k^1)$ and $c^\dagger(k^1)$ (2.31) should be replaced by the operators $\tilde{c}(k^1)$ and $\tilde{c}^\dagger(k^1)$ defined by

$$\begin{aligned} \tilde{c}(k^1) &= \frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \theta(k^1 q^1) [a^\dagger(q^1) a(k^1 + q^1) - \varepsilon(q^1) \varepsilon(k^1 + q^1) b^\dagger(q^1) b(k^1 + q^1)] \right. \\ &\quad \left. + \theta(q^1(k^1 - q^1)) \varepsilon(k^1 - q^1) b(k^1 - q^1) a(q^1) \right\}, \\ \tilde{c}^\dagger(k^1) &= -\frac{i}{\sqrt{k^0}} \int_{-\infty}^{\infty} dq^1 \left\{ \theta(k^1 q^1) [a^\dagger(k^1 + q^1) a(q^1) - \varepsilon(q^1) \varepsilon(k^1 + q^1) b^\dagger(k^1 + q^1) b(q^1)] \right. \\ &\quad \left. + \theta(q^1(k^1 - q^1)) \varepsilon(k^1 - q^1) a^\dagger(q^1) b^\dagger(k^1 - q^1) \right\}. \end{aligned} \quad (4.23)$$

One can show that these operators $\tilde{c}(k^1)$ and $\tilde{c}^\dagger(k^1)$ are scalars under parity transformation

$$\begin{aligned}\mathcal{P}\tilde{c}(k^1)\mathcal{P}^\dagger &= \tilde{c}(-k^1), \\ \mathcal{P}\tilde{c}^\dagger(k^1)\mathcal{P}^\dagger &= \tilde{c}^\dagger(-k^1).\end{aligned}\tag{4.24}$$

One can also show that the operators $\tilde{c}(k^1)$ and $\tilde{c}^\dagger(k^1)$ obey canonical commutation relations

$$\begin{aligned}[\tilde{c}(k^1), \tilde{c}^\dagger(p^1)] &= \delta(k^1 - p^1), \\ [\tilde{c}(k^1), \tilde{c}(p^1)] &= [\tilde{c}^\dagger(k^1), \tilde{c}^\dagger(p^1)] = 0.\end{aligned}\tag{4.25}$$

Since $\varepsilon(p^1)u(p^0, p^1) = v(p^0, p^1)$, where $v(p^0, p^1)$ is defined by (2.15) the operators $\tilde{c}(k^1)$ and $\tilde{c}^\dagger(k^1)$ introduced in Ref.[16] give other representation of the operators $C(k^1)$ and $C^\dagger(k^1)$, which we have defined in this section.

5 Quantization of the massless Thirring model in Klaiber's operator formalism and quantum equation of motion

Following Klaiber for the quantization of massless Thirring fermion fields and the evaluation of correlation functions we have to represent the classical solution of the classical equation of motion (4.18) as follows

$$\begin{aligned}\psi(x) &= e^{-i\alpha S^{(-)}(x) - i\beta\gamma^5 P^{(-)}(x)} \Psi(x) e^{-i\alpha S^{(+)}(x) - i\beta\gamma^5 P^{(+)}(x)}, \\ \bar{\psi}(x) &= e^{+i\alpha S^{(-)}(x) - i\beta\gamma^5 P^{(-)}(x)} \bar{\Psi}(x) e^{+i\alpha S^{(+)}(x) - i\beta\gamma^5 P^{(+)}(x)},\end{aligned}\tag{5.1}$$

where $S^{(\pm)}(x)$ and $P^{(\pm)}(x)$ are positive and negative frequency parts of the operators $S(x)$ and $P(x)$, respectively:

$$\begin{aligned}S^{(+)}(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} C(k^1) e^{-ik \cdot x}, \\ S^{(-)}(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} C^\dagger(k^1) e^{+ik \cdot x}, \\ P^{(+)}(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \varepsilon(k^1) C(k^1) e^{-ik \cdot x}, \\ P^{(-)}(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \varepsilon(k^1) C^\dagger(k^1) e^{+ik \cdot x}.\end{aligned}\tag{5.2}$$

In analogy we define the positive and negative frequency parts $\Psi^{(+)}(x)$ and $\Psi^{(-)}(x)$ of the Ψ -field where $\Psi^{(+)}(y)$ is the positive frequency part of the Ψ -field

$$\begin{aligned}\Psi^{(+)}(y) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2p^0}} u(p^0, p^1) a(p^1) e^{-ip \cdot y}, \\ \Psi^{(-)}(y) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2p^0}} v(p^0, p^1) b^\dagger(p^1) e^{+ip \cdot y}.\end{aligned}\tag{5.3}$$

The operator relations (5.1) should be understood in matrix form as follows

$$\begin{aligned}\psi_a(x) &= \left(e^{-i\alpha S^{(-)}(x) - i\beta\gamma^5 P^{(-)}(x)} \right)_{cb} \Psi_c(x) \left(e^{-i\alpha S^{(+)}(x) - i\beta\gamma^5 P^{(+)}(x)} \right)_{ba}, \\ \bar{\psi}_a(x) &= \left(e^{+i\alpha S^{(-)}(x) - i\beta\gamma^5 P^{(-)}(x)} \right)_{cb} \bar{\Psi}_c(x) \left(e^{+i\alpha S^{(+)}(x) - i\beta\gamma^5 P^{(+)}(x)} \right)_{ba},\end{aligned}\quad (5.4)$$

where Latin indices run over $a = 1, 2$.

The operators $S^{(\pm)}(x)$ and $P^{(\pm)}(x)$ obey the c -valued commutation relations

$$\begin{aligned}[S^{(+)}(x), S^{(-)}(y)] &= [P^{(+)}(x), P^{(-)}(y)] = -i D^{(+)}(x - y), \\ [S^{(-)}(x), S^{(+)}(y)] &= [P^{(-)}(x), P^{(+)}(y)] = -i D^{(-)}(x - y), \\ [S^{(+)}(x), P^{(-)}(y)] &= -i D_5^{(+)}(x - y), \\ [S^{(-)}(x), P^{(+)}(y)] &= -i D_5^{(-)}(x - y),\end{aligned}\quad (5.5)$$

where the correlation functions in the r.h.s. of (5.3) are determined by [4] (see also [16])

$$\begin{aligned}D^{(\pm)}(x) &= \pm \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{2k^0} e^{\mp i k \cdot x} = \mp \frac{i}{4\pi} \ell n[-\mu^2 x^2 \pm i 0 \cdot \varepsilon(x^0)], \\ D_5^{(\pm)}(x) &= \pm \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{2k^0} \varepsilon(k^1) e^{\mp i k \cdot x} = \mp \frac{i}{4\pi} \ell n\left(\frac{x^0 + x^1 \mp i 0}{x^0 - x^1 \mp i 0}\right),\end{aligned}\quad (5.6)$$

where μ is an infrared cut-off and $x^2 = (x^0)^2 - (x^1)^2$. The detailed calculation of the correlation functions $D^{(\pm)}(x)$ and $D_5^{(\pm)}(x)$ we adduce in Appendix C. Under Lorentz transformations

$$\begin{aligned}x^0 \rightarrow \tilde{x}^0 &= \frac{x^0 - vx^1}{\sqrt{1-v^2}}, \\ x^1 \rightarrow \tilde{x}^1 &= \frac{x^1 - vx^0}{\sqrt{1-v^2}},\end{aligned}\quad (5.7)$$

where v is the velocity of a Lorentz frame, the correlation functions $D^{(\pm)}(x)$ and $D_5^{(\pm)}(x)$ behave as follows

$$\begin{aligned}D^{(\pm)}(x) \rightarrow D^{(\pm)}(\tilde{x}) &= D^{(\pm)}(x), \\ D_5^{(\pm)}(x) \rightarrow D_5^{(\pm)}(\tilde{x}) &= D_5^{(\pm)}(x) \pm \frac{i}{4\pi} \ell n\left(\frac{1+v}{1-v}\right).\end{aligned}\quad (5.8)$$

Hence, the correlation function $D^{(\pm)}(x)$ is covariant and $D_5^{(\pm)}(x)$ is non-covariant under Lorentz transformations, respectively, since $x^2 \rightarrow \tilde{x}^2 = x^2$.

Now let us evaluate of the commutators $[S^{(\pm)}(x), \Psi(y)]$ and $[P^{(\pm)}(x), \Psi(y)]$. The knowledge of these commutators is important for the correct (i) derivation of the quantum equation of motion for the massless Thirring model from the operator field representation (5.1) and (ii) evaluation of correlation functions of massless Thirring fermion fields.

Using (5.2) and (2.13) the evaluation of the commutator $[S^{(+)}(x), \Psi(y)]$ runs in the way

$$\begin{aligned}[S^{(+)}(x), \Psi(y)] &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2k^0}} \int_{-\infty}^{\infty} dp^1 \left[\begin{aligned} &\begin{pmatrix} \theta(+p^1) \\ \theta(-p^1) \end{pmatrix} [C(k^1), a(p^1)] e^{-ik \cdot x - ip \cdot y} \\ &+ \begin{pmatrix} +\theta(+p^1) \\ -\theta(-p^1) \end{pmatrix} [C(k^1), b^\dagger(p^1)] e^{-ik \cdot x + ip \cdot y} \end{aligned} \right].\end{aligned}\quad (5.9)$$

According to the definition of the C -operator (4.13) and the canonical anti-commutation relations (2.14) we get

$$\begin{aligned} [C(k^1), a(p^1)] &= -\frac{i}{\sqrt{k^0}} \theta(k^1 p^1) a(k^1 + p^1), \\ [C(k^1), b^\dagger(p^1)] &= -\frac{i}{\sqrt{k^0}} \{ \theta(k^1(p^1 - k^1)) b^\dagger(p^1 - k^1) + \varepsilon(k^1) \theta(p^1(k^1 - p^1)) a(k^1 - p^1) \} \end{aligned} \quad (5.10)$$

Substituting (5.10) in (5.9), making the corresponding changes of variables and using the properties of the Heaviside functions we end up with the expression

$$\begin{aligned} [S^{(+)}(x), \Psi(y)] &= \\ &= -\sqrt{\pi} \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{k^0} e^{-ik \cdot (x-y)} \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \left[\left(\frac{\theta(+k^1)\theta(+p^1)\theta(p^1 - k^1)}{\theta(-k^1)\theta(-p^1)\theta(k^1 - p^1)} \right) a(p^1) e^{-ip \cdot y} \right. \\ &\quad \left. + \left(\frac{\theta(+k^1)\theta(+p^1)\theta(k^1 - p^1)}{\theta(-k^1)\theta(-p^1)\theta(p^1 - k^1)} \right) a(p^1) e^{-ip \cdot y} + \left(\frac{+\theta(+k^1)\theta(+p^1)}{-\theta(-k^1)\theta(-p^1)} \right) b^\dagger(p^1) e^{+ip \cdot y} \right] = \\ &= -\sqrt{\pi} \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{k^0} e^{-ik \cdot (x-y)} \\ &\quad \times \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \left[\left(\frac{\theta(+k^1)\theta(+p^1)}{\theta(-k^1)\theta(-p^1)} \right) a(p^1) e^{-ip \cdot y} + \left(\frac{+\theta(+k^1)\theta(+p^1)}{-\theta(-k^1)\theta(-p^1)} \right) b^\dagger(p^1) e^{+ip \cdot y} \right]. \end{aligned} \quad (5.11)$$

Due to the relations

$$\theta(k^1) = \frac{1 + \varepsilon(k^1)}{2}, \quad \theta(-k^1) = \frac{1 - \varepsilon(k^1)}{2} \quad (5.12)$$

the r.h.s. of (5.11) can be recast into the form

$$\begin{aligned} [S^{(+)}(x), \Psi(y)] &= -\sqrt{\pi} \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{2k^0} e^{-ik \cdot (x-y)} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2p^0}} \left[u(p^0, p^1) a(p^1) e^{-ip \cdot y} \right. \\ &\quad \left. + \varepsilon(k^1) \gamma^5 u(p^0, p^1) a(p^1) e^{-ip \cdot y} + v(p^0, p^1) b^\dagger(p^1) e^{+ip \cdot y} + \varepsilon(k^1) \gamma^5 v(p^0, p^1) b^\dagger(p^1) e^{+ip \cdot y} \right] \\ &= -\sqrt{\pi} [D^{(+)}(x-y) + \gamma^5 D_5^{(+)}(x-y)] \Psi(y). \end{aligned} \quad (5.13)$$

Thus, we have obtained

$$[S^{(+)}(x), \Psi(y)] = -\sqrt{\pi} [D^{(+)}(x-y) + \gamma^5 D_5^{(+)}(x-y)] \Psi(y). \quad (5.14)$$

This is in agreement with Klaiber's commutator (see (IV.13) of Ref.[4]).

In a similar way we calculate the commutators

$$\begin{aligned} [S^{(-)}(x), \Psi(y)] &= -\sqrt{\pi} [D^{(-)}(x-y) + \gamma^5 D_5^{(-)}(x-y)] \Psi(y), \\ [P^{(+)}(x), \Psi(y)] &= -\sqrt{\pi} [D_5^{(+)}(x-y) + \gamma^5 D^{(+)}(x-y)] \Psi(y), \\ [P^{(-)}(x), \Psi(y)] &= -\sqrt{\pi} [D_5^{(-)}(x-y) + \gamma^5 D^{(-)}(x-y)] \Psi(y). \end{aligned} \quad (5.15)$$

For the further analysis we need the commutation relations between the scalar and pseudoscalar densities $S^{(\pm)}(x)$ and $P^{(\pm)}(x)$ and $\bar{\Psi}(y)$. These commutators read

$$\begin{aligned} [S^{(+)}(x), \bar{\Psi}(y)] &= +\sqrt{\pi} [D^{(+)}(x-y) - \gamma^5 D_5^{(+)}(x-y)] \bar{\Psi}(y), \\ [S^{(-)}(x), \bar{\Psi}(y)] &= +\sqrt{\pi} [D^{(-)}(x-y) - \gamma^5 D_5^{(-)}(x-y)] \bar{\Psi}(y), \\ [P^{(+)}(x), \bar{\Psi}(y)] &= +\sqrt{\pi} [D_5^{(+)}(x-y) - \gamma^5 D^{(+)}(x-y)] \bar{\Psi}(y), \\ [P^{(-)}(x), \bar{\Psi}(y)] &= +\sqrt{\pi} [D_5^{(-)}(x-y) - \gamma^5 D^{(-)}(x-y)] \bar{\Psi}(y). \end{aligned} \quad (5.16)$$

For the evaluation of correlation functions one needs the relations

$$\begin{aligned}
& e^{+i\alpha S^{(+)}(x) \pm i\beta\gamma^5 P^{(+)}(x)} \Psi(y) e^{-i\alpha S^{(+)}(x) \mp i\beta\gamma^5 P^{(+)}(x)} = \\
& = e^{-i\sqrt{\pi}(\alpha \pm \beta) [D^{(+)}(x-y) + \gamma^5 D_5^{(+)}(x-y)]} \Psi(y), \\
& e^{+i\alpha S^{(-)}(x) \pm i\beta\gamma^5 P^{(-)}(x)} \Psi(y) e^{-i\alpha S^{(-)}(x) \mp i\beta\gamma^5 P^{(-)}(x)} = \\
& = e^{-i\sqrt{\pi}(\alpha \pm \beta) [D^{(-)}(x-y) + \gamma^5 D_5^{(-)}(x-y)]} \Psi(y), \\
& e^{+i\alpha S^{(+)}(x) \pm i\beta\gamma^5 P^{(+)}(x)} \bar{\Psi}(y) e^{-i\alpha S^{(+)}(x) \mp i\beta\gamma^5 P^{(+)}(x)} = \\
& = \bar{\Psi}(y) e^{+i\sqrt{\pi}(\alpha \mp \beta) [D^{(+)}(x-y) - \gamma^5 D_5^{(+)}(x-y)]}, \\
& e^{+i\alpha S^{(-)}(x) \pm i\beta\gamma^5 P^{(-)}(x)} \bar{\Psi}(y) e^{-i\alpha S^{(-)}(x) \mp i\beta\gamma^5 P^{(-)}(x)} = \\
& = \bar{\Psi}(y) e^{+i\sqrt{\pi}(\alpha \mp \beta) [D^{(-)}(x-y) - \gamma^5 D_5^{(-)}(x-y)]}. \tag{5.17}
\end{aligned}$$

Now we are able to derive the quantum equation of motion for the massless Thirring model. For this aim we act with the operator $i\gamma^\mu \partial_\mu$ on the operator of the $\psi(x)$ -field defined by (5.1).

We start the derivation of the quantum equation of motion from the transformation of the ψ -field defined by (5.1). We suggest to use the space-time splitting method suggested by Schwinger [22]. Within Schwinger's splitting method we understand the $\psi(x)$ -field given by (5.1) as a limit

$$\psi(x) = \lim_{\epsilon \rightarrow 0} e^{-iQ^{(-)}(x+\epsilon)} \Psi(x) e^{-iQ^{(+)}(x-\epsilon)}, \tag{5.18}$$

where ϵ is an infinitesimal space-like 2-vector, $\epsilon^2 < 0$ [4,21]. Then, for convenience we have introduced the notations: $Q^{(\pm)}(x \mp \epsilon) = \alpha S^{(\pm)}(x \mp \epsilon) + \beta\gamma^5 P^{(\pm)}(x \mp \epsilon)$.

Now we have to calculate the quantity $i\gamma^\mu \partial_\mu \psi(x)$. The differentiation of exponentials depending of operators $Q^{(\mp)}(x \pm \epsilon)$ we would perform by using the formula

$$i\partial_\mu \left(e^{-iQ^{(\mp)}(x \pm \epsilon)} \right) = \int_0^1 dt e^{-i(1-t)Q^{(\mp)}(x \pm \epsilon)} \partial_\mu Q^{(\mp)}(x \pm \epsilon) e^{-itQ^{(\mp)}(x \pm \epsilon)}. \tag{5.19}$$

Applying (5.19) to the calculation of $i\gamma^\mu \partial_\mu \psi(x)$ we obtain

$$\begin{aligned}
i\gamma^\mu \partial_\mu \psi(x) &= \lim_{\epsilon \rightarrow 0} \left[\gamma^\mu \int_0^1 dt e^{-i(1-t)Q^{(-)}(x+\epsilon)} \partial_\mu Q^{(-)}(x+\epsilon) \Psi(x) e^{-iQ^{(+)}(x-\epsilon)} \right. \\
&+ \gamma^\mu e^{-iQ^{(-)}(x+\epsilon)} \Psi(x) \int_0^1 dt e^{-i(1-t)Q^{(+)}(x-\epsilon)} \partial_\mu Q^{(+)}(x-\epsilon) \Psi(x) \\
&\times e^{-itQ^{(+)}(x-\epsilon)} \Big] = \lim_{\epsilon \rightarrow 0} \left[\gamma^\mu \partial_\mu Q^{(-)}(x+\epsilon) e^{-iQ^{(-)}(x+\epsilon)} \Psi(x) e^{-iQ^{(+)}(x-\epsilon)} \right. \\
&+ \gamma^\mu e^{-iQ^{(-)}(x+\epsilon)} \Psi(x) e^{-iQ^{(+)}(x-\epsilon)} \partial_\mu Q^{(+)}(x-\epsilon) \Big] = \partial_\mu Q^{(-)}(x) \gamma^\mu \psi(x) \\
&+ \gamma^\mu \psi(x) \partial_\mu Q^{(+)}(x) = \sqrt{\pi}(\alpha - \beta) J_\mu^{(-)}(x) \gamma^\mu \psi(x) + \sqrt{\pi}(\alpha - \beta) \gamma^\mu \psi(x) J_\mu^{(+)}(x) = \\
&= \sqrt{\pi}(\alpha - \beta) : J_\mu(x) \gamma^\mu \psi(x) : . \tag{5.20}
\end{aligned}$$

Here we have used the Wick definition of a normal-ordered product [24,25]

$$:J_\mu(x)\psi(x):=J_\mu^{(-)}(x)\psi(x)+\psi(x)J_\mu^{(+)}(x). \quad (5.21)$$

Thus, we have got

$$i\gamma^\mu\partial_\mu\psi(x)=\sqrt{\pi}(\alpha-\beta):J_\mu(x)\gamma^\mu\psi(x): \quad (5.22)$$

In order to transform this equation to the equation of motion of the massless Thirring model we have to find the relation between the quantum currents $J_\mu(x)=:\bar{\Psi}(x)\gamma_\mu\Psi(x):$ and $j_\mu(x)=:\bar{\psi}(x)\gamma_\mu\psi(x):$.

The normal-ordered current $j_\mu(x)=:\bar{\psi}(x)\gamma_\mu\psi(x):$ we define according to Klaiber as follows

$$j_\mu(x)=:\bar{\psi}(x)\gamma_\mu\psi(x):=\frac{1}{2}\lim_{\epsilon\rightarrow 0}(\gamma^\mu)_{ab}[\bar{\psi}_a(x+\epsilon)\psi_b(x)-\psi_b(x+\epsilon)\bar{\psi}_b(x)], \quad (5.23)$$

where ϵ is an infinitesimal 2-vector.

First, let us treat the quantity $\bar{\psi}(x+\epsilon)\gamma^\mu\psi(x)=\bar{\psi}_a(x+\epsilon)(\gamma^\mu)_{ab}\psi_b(x):$

$$\begin{aligned} \bar{\psi}_a(x+\epsilon)(\gamma^\mu)_{ab}\psi_b(x) &= \left(e^{+i\bar{Q}^{(-)}(x+\epsilon)}\right)_{cd}\bar{\Psi}_c(x+\epsilon)\left(e^{+i\bar{Q}^{(+)}(x+\epsilon)}\right)_{da} \\ &\quad \times (\gamma^\mu)_{ab}\left(e^{-iQ^{(-)}(x)}\right)_{fe}\Psi_e(x)\left(e^{-iQ^{(+)}(x)}\right)_{fb}, \end{aligned} \quad (5.24)$$

where we have denoted $\bar{Q}^{(+)}(x+\epsilon)=\alpha S^{(+)}(x+\epsilon)-\beta\gamma^5 P^{(+)}(x+\epsilon)$.

Using the commutation relations (5.5) and the relations (5.17) we recast the r.h.s. of (5.24) into the form

$$\begin{aligned} \bar{\psi}_a(x+\epsilon)(\gamma^\mu)_{ab}\psi_b(x) &= \left(e^{+i\bar{Q}^{(-)}(x+\epsilon)-i\bar{Q}^{(-)}(x)}\right)_{cd}\bar{\Psi}_c(x+\epsilon)(\gamma^\mu)_{da} \\ &\quad \times \left(e^{-i[(\alpha^2+\beta^2)+2\sqrt{\pi}(\alpha+\beta)]D^{(+)}(\epsilon)-2i[\alpha\beta+\sqrt{\pi}(\alpha+\beta)]\gamma^5 D_5^{(+)}(\epsilon)}\right)_{ab} \\ &\quad \times \Psi_e(x)\left(e^{+iQ^{(+)}(x+\epsilon)-iQ^{(+)}(x)}\right)_{eb}. \end{aligned} \quad (5.25)$$

This expression contains the scalar, $D^{(+)}(\epsilon)$, and pseudoscalar, $\gamma^5 D_5^{(+)}(\epsilon)$, divergences at the limit $\epsilon \rightarrow 0$. Since pseudoscalar divergences can be removed by neither renormalization of wave functions of fermion fields nor chiral rotations, we should impose the constraint

$$\alpha\beta+\sqrt{\pi}(\alpha+\beta)=0. \quad (5.26)$$

As we have shown in (5.8) the functions $D_5^{(+)}(\epsilon)$ are non-covariant under Lorentz transformations. Therefore, removal of contributions of these functions is required too by Lorentz covariance. This agrees with Klaiber's constraint making correlation functions covariant under Lorentz transformations (see section V of Ref.[4]).

Due to the constraint (5.26) the relation (5.25) takes the form

$$\begin{aligned} \bar{\psi}_a(x+\epsilon)(\gamma^\mu)_{ab}\psi_b(x) &= e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)}\left(e^{+i\bar{Q}^{(-)}(x+\epsilon)-i\bar{Q}^{(-)}(x)}\right)_{cd} \\ &\quad \times \bar{\Psi}_c(x+\epsilon)(\gamma^\mu)_{da}\Psi_e(x)\left(e^{+iQ^{(+)}(x+\epsilon)-iQ^{(+)}(x)}\right)_{ea}. \end{aligned} \quad (5.27)$$

Now we have to rewrite the product $\bar{\Psi}_c(x+\epsilon)\Psi_e(x)$ in the normal-ordered form. According to Wick's theorem [24] we get

$$\begin{aligned}\bar{\Psi}_c(x+\epsilon)\Psi_e(x) &= :\bar{\Psi}_c(x+\epsilon)\Psi_e(x): + \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{dk^1}{2k^0} (\hat{k})_{ec} e^{-ik \cdot \epsilon} = \\ &= :\bar{\Psi}_c(x+\epsilon)\Psi_e(x): + (\hat{\epsilon})_{ec} \delta(\epsilon^2) + \frac{1}{2\pi i} \frac{(\hat{\epsilon})_{ec}}{\epsilon^2}.\end{aligned}\quad (5.28)$$

Recall, that the momentum integral in (5.28) has been calculated in [6] (see Eq.(6.6) of Ref.[6]).

Substituting (5.5) in (5.27), expanding the exponentials in powers of ϵ and keeping only finite and leading divergent contributions we arrive at the expression

$$\begin{aligned}\bar{\psi}(x+\epsilon)\gamma^\mu\psi(x) &= e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \left[\frac{1}{\pi i} \frac{\epsilon^\mu}{\epsilon^2} + \epsilon^\mu \delta(\epsilon^2) + :\bar{\Psi}(x)\gamma^\mu\Psi(x): \right. \\ &+ \frac{1}{2\pi} \frac{\epsilon^\nu}{\epsilon^2} \text{tr}\{\hat{\epsilon}\gamma^\mu[\alpha\partial_\nu S(x) + \beta\gamma^5\partial_\nu P(x)]\} \left. \right] = e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \left[\frac{1}{\pi i} \frac{\epsilon^\mu}{\epsilon^2} + \epsilon^\mu \delta(\epsilon^2) \right. \\ &+ J^\mu(x) + \frac{\alpha}{\sqrt{\pi}} \frac{\epsilon^\mu\epsilon^\nu}{\epsilon^2} J_\nu(x) - \frac{\beta}{\sqrt{\pi}} \frac{\epsilon_\lambda\epsilon_\nu}{\epsilon^2} \varepsilon^{\lambda\mu}\varepsilon^{\nu\rho} J_\rho(x) \left. \right].\end{aligned}\quad (5.29)$$

Using then the identity

$$\varepsilon^{\lambda\mu}\varepsilon^{\nu\rho} = g^{\mu\nu}g^{\lambda\rho} - g^{\mu\rho}g^{\lambda\nu}\quad (5.30)$$

we get

$$\begin{aligned}\bar{\psi}(x+\epsilon)\gamma^\mu\psi(x) &= \frac{4i\epsilon^\mu}{(\alpha-\beta)^2} \frac{d}{d\epsilon^2} \left[e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \right] \\ &+ e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \left[\left(1 + \frac{\beta}{\sqrt{\pi}}\right) J^\mu(x) + \frac{\alpha-\beta}{\sqrt{\pi}} \frac{\epsilon^\mu\epsilon^\nu}{\epsilon^2} J_\nu(x) \right],\end{aligned}\quad (5.31)$$

where we have used the relation

$$e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \left[\frac{1}{\pi i} \frac{\epsilon^\mu}{\epsilon^2} + \epsilon^\mu \delta(\epsilon^2) \right] = \frac{4i\epsilon^\mu}{(\alpha-\beta)^2} \frac{d}{d\epsilon^2} \left[e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \right],\quad (5.32)$$

which follows from the definition of $D^{(+)}(\epsilon)$ given by (5.5).

In a similar way we can treat the second term in the r.h.s. of (5.23). As result we get

$$\begin{aligned}(\gamma^\mu)_{ab}\psi_b(x+\epsilon)\bar{\psi}_a(x) &= e^{-i(\alpha-\beta)^2 D^{(+)}(\epsilon)} \left(e^{-i\bar{Q}^{(-)}(x+\epsilon)} + i\bar{Q}^{(-)}(x) \right)_{cd} \\ &\times \Psi_c(x+\epsilon)(\gamma^\mu)_{db}\bar{\Psi}_a(x) \left(e^{-i\bar{Q}^{(+)}(x+\epsilon)} + i\bar{Q}^{(+)}(x) \right)_{ab}.\end{aligned}\quad (5.33)$$

In the normal-ordered form the product $\Psi_c(x+\epsilon)(\gamma^\mu)_{db}\bar{\Psi}_a(x)$ is defined by

$$\Psi_c(x+\epsilon)\bar{\Psi}_a(x) = -:\bar{\Psi}_a(x)\Psi_c(x+\epsilon): + (\hat{\epsilon})_{ca} \delta(\epsilon^2) + \frac{1}{2\pi i} \frac{(\hat{\epsilon})_{ca}}{\epsilon^2}.\quad (5.34)$$

Substituting (5.34) in (5.33) and expanding in powers of ϵ , calculating traces over Dirac matrices and using the identity (5.30) we obtain

$$\begin{aligned} (\gamma^\mu)_{ab} \psi_b(x + \epsilon) \bar{\psi}_a(x) &= \frac{4i\epsilon^\mu}{(\alpha - \beta)^2} \frac{d}{d\epsilon^2} \left[e^{-i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \right] \\ &- e^{-i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \left[\left(1 + \frac{\beta}{\sqrt{\pi}} \right) J^\mu(x) + \frac{\alpha - \beta}{\sqrt{\pi}} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^2} J_\nu(x) \right]. \end{aligned} \quad (5.35)$$

Thus, the normal-ordered vector current $j^\mu(x)$ is defined by

$$\begin{aligned} j^\mu(x) &= : \bar{\psi}(x) \gamma^\mu \psi(x) : = \\ &= e^{-i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \left[\left(1 + \frac{\beta}{\sqrt{\pi}} \right) J^\mu(x) + \frac{\alpha - \beta}{\sqrt{\pi}} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^2} J_\nu(x) \right]. \end{aligned} \quad (5.36)$$

In order to remove the non-covariant term proportional to $\epsilon^\mu \epsilon^\nu / \epsilon^2$ we can follow the standard point-splitting procedure [22,25] and average over all directions of the ϵ^μ . However, Klaiber has invented another method which leads, unfortunately, to violation of chiral symmetry at intermediate steps.

In fact, Klaiber suggested to define a renormalized vector current $K^\mu(x)$ as follows

$$\begin{aligned} K^\mu(x) &= \lim_{\delta \rightarrow 0} \lim_{\epsilon \rightarrow 0} e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{1}{2} (\gamma_\nu)_{ab} \{ \bar{\psi}_a(x + \epsilon) \psi_b(x) [g^{\mu\nu} (1 - i g \epsilon^\lambda K_\lambda(x + \delta) \\ &- i \sigma \epsilon^\nu K^\mu(x + \delta)) - \psi_b(x + \epsilon) \bar{\psi}_a(x) [g^{\mu\nu} (1 + i g \epsilon^\lambda K_\lambda(x + \delta) + i \sigma \epsilon^\nu K^\mu(x + \delta))] \}, \end{aligned} \quad (5.37)$$

where σ is a real parameter $\sigma \in \mathbb{R}^1$.

Using equations (5.28) and (5.34) we recast the r.h.s. of (5.37) into the form

$$K^\mu(x) = -\frac{\sigma}{\pi} K^\mu(x) + \left(1 + \frac{\beta}{\sqrt{\pi}} \right) J^\mu(x) + \left[\frac{\alpha - \beta}{\sqrt{\pi}} J_\nu(x) - \frac{g}{\pi} K_\nu(x) \right] \lim_{\epsilon \rightarrow 0} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^2}. \quad (5.38)$$

We would like to emphasize that the limit $\delta \rightarrow 0$ is not singular and can be taken from the very beginning. So its presence is rather formal. The requirement of the vanishing of the non-covariant terms gives

$$K_\nu(x) = \sqrt{\pi} \frac{\alpha - \beta}{g} J_\nu(x). \quad (5.39)$$

Substituting (5.39) into (5.22) we get

$$i\gamma^\mu \partial_\mu \psi(x) = g : K_\mu(x) \gamma^\mu \psi(x) :. \quad (5.40)$$

According to Klaiber this is the quantum equation of motion for the massless Thirring model quantized in the chiral symmetric phase. However, equation (5.38) together with (5.39) leads to a relation between the parameters α , β and σ

$$1 + \frac{\beta}{\sqrt{\pi}} = \frac{\pi + \sigma}{g} \frac{\alpha - \beta}{\sqrt{\pi}}. \quad (5.41)$$

Due to the relation (5.25) this leaves only one free parameter in the approach confirming the one-parameter family of solutions of the massless Thirring model [4].

Equation (5.25) can be fulfilled identically by [4]

$$\alpha = \sqrt{\pi}(\rho - 1) \quad , \quad \beta = \sqrt{\pi}\left(\frac{1}{\rho} - 1\right), \quad (5.42)$$

where $\rho \in \mathbb{R}^1$ and $\rho \neq 0$ [4]. The solution of equation (5.41) with respect to ρ is equal to

$$\rho = \pm \sqrt{1 + \frac{g}{\pi + \sigma}}. \quad (5.43)$$

This agrees with Klaiber's result (see Eq.(X.13) of Ref.[4]).

However, we would like to emphasize that this method is not innocent and leads to a breaking of chiral symmetry. In order to show this we have to define the axial vector current $K_5^\mu(x)$ in a way analogous to the vector current. First of all notice that the δ -limit is non-singular and, without loss of generality, we can take the limit $\delta \rightarrow 0$ from the very beginning. This reduces the r.h.s. of (5.37) to the form

$$K^\mu(x) = \lim_{\epsilon \rightarrow 0} e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{1}{2} (\gamma_\nu)_{ab} \{ \bar{\psi}_a(x + \epsilon) \psi_b(x) [g^{\mu\nu}(1 - i g \epsilon^\lambda K_\lambda(x)) - i \sigma \epsilon^\nu K^\mu(x)] - \psi_b(x + \epsilon) \bar{\psi}_a(x) [g^{\mu\nu}(1 + i g \epsilon^\lambda K_\lambda(x)) + i \sigma \epsilon^\nu K^\mu(x)] \}, \quad (5.44)$$

which can be rewritten as follows (see Appendix D)

$$K^\mu(x) = \lim_{\epsilon \rightarrow 0} e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \times \frac{1}{2} \{ \bar{\psi}_a(x + \epsilon) (\gamma_\nu)_{ab} e^{-i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \psi_b(x) - i \sigma (\gamma_\nu)_{ab} \bar{\psi}_a(x + \epsilon) \psi_b(x) \epsilon^\nu K^\mu(x) \} - (\gamma_\nu)_{ab} \psi_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\psi}_a(x) + i \sigma (\gamma_\nu)_{ab} \psi_b(x + \epsilon) \bar{\psi}_a(x) \epsilon^\nu K^\mu(x) \}. \quad (5.45)$$

The appearance of the vector field in the r.h.s. of (5.45) can be explained by the requirement of gauge invariance. The vector current $j^\mu(x) =: \bar{\psi}(x) \gamma^\mu \psi(x) :$ and the axial-vector current $j_5^\mu(x) =: \bar{\psi}(x) \gamma^\mu \gamma^5 \psi(x) :$ are invariant under gauge transformations

$$\begin{aligned} \psi(x) &\rightarrow \psi'(x) = e^{-i\omega(x)} \psi(x), \\ \bar{\psi}(x) &\rightarrow \bar{\psi}'(x) = \bar{\psi} e^{+i\omega(x)}, \end{aligned} \quad (5.46)$$

where $\omega(x)$ is a gauge function. Gauge invariance breaks down for the products $\bar{\psi}_a(x + \epsilon) \psi_b(x)$ and $\psi_b(x + \epsilon) \bar{\psi}_a(x)$

$$\begin{aligned} \bar{\psi}_a(x + \epsilon) \psi_b(x) &\rightarrow \bar{\psi}'_a(x + \epsilon) \psi'_b(x) = \bar{\psi}_a(x + \epsilon) \psi_b(x) e^{+i\omega(x + \epsilon) - i\omega(x)}, \\ \psi_b(x + \epsilon) \bar{\psi}_a(x) &\rightarrow \psi'_b(x + \epsilon) \bar{\psi}'_a(x) = \psi_b(x + \epsilon) \bar{\psi}_a(x) e^{-i\omega(x + \epsilon) + i\omega(x)}. \end{aligned} \quad (5.47)$$

Since a difference of gauge functions $\omega(x + \epsilon) - \omega(x)$ can be arbitrary large despite the infinitesimal magnitude of ϵ [26], the violation of gauge invariance can be arbitrarily strong. The presence of the vector field $K_\lambda(x)$ restores gauge invariance of the products (5.47):

$$\begin{aligned} \bar{\psi}_a(x + \epsilon) e^{-i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \psi_b(x) &= \bar{\psi}'_a(x + \epsilon) e^{-i g \int_x^{x+\epsilon} dz^\lambda K'_\lambda(z)} \psi'_b(x), \\ \psi_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\psi}_a(x) &\rightarrow \psi'_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K'_\lambda(z)} \bar{\psi}'_a(x), \end{aligned} \quad (5.48)$$

if the fields $K_\lambda(z)$ and $K'_\lambda(z)$ are related by

$$K'_\lambda(z) = K_\lambda(z) + \frac{1}{g} \partial_\lambda \omega(z). \quad (5.49)$$

As a result gauge invariance of the vector current defined by the expression (5.44) is broken only by non-covariant terms proportional to σ . This violation is of order $O(\epsilon)$.

Now we are able to define the axial-vector current $K_5^\mu(x)$

$$K_5^\mu(x) = \lim_{\epsilon \rightarrow 0} e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{1}{2} (\gamma_\nu \gamma^5)_{ab} \{ \bar{\psi}_a(x + \epsilon) \psi_b(x) [g^{\mu\nu} (1 - i g \epsilon^\lambda K_\lambda(x)) - i \sigma \epsilon^\nu K^\mu(x)] - \psi_b(x + \epsilon) \bar{\psi}_a(x) [g^{\mu\nu} (1 + i g \epsilon^\lambda K_\lambda(x)) + i \sigma \epsilon^\nu K^\mu(x)] \} \quad (5.50)$$

and consider the divergence of the axial-vector current $\partial_\mu K_5^\mu(x)$. The terms proportional to $\epsilon^\lambda K_\lambda(x)$ are invented to retain gauge invariance of the axial-vector current, and the non-covariant terms proportional to $\epsilon^\nu K^\mu(x)$ provide a smooth violation of gauge invariance of order $O(\epsilon)$.

We should emphasize that the correct definition of the quantum vector current $K^\mu(x)$ should give us the quantum axial-vector current $K_5^\mu(x)$ related to $K^\mu(x)$ by the standard relation: $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$.

According to the definition of the axial-vector current (5.50) the divergence $\partial_\mu K_5^\mu(x)$ is determined by

$$\begin{aligned} \partial_\mu K_5^\mu(x) &= \lim_{\epsilon \rightarrow 0} e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} (-i) g \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} [\bar{\psi}_a(x + \epsilon) \psi_b(x) + \psi_b(x + \epsilon) \bar{\psi}_a(x)] \\ &\quad \times \epsilon^\lambda \partial_\mu K_\lambda(x), \end{aligned} \quad (5.51)$$

where we have taken into account the conservation of the vector current $\partial_\mu K^\mu(x) = 0$.

Using the definition of the self-coupled fermionic fields ψ in terms of the free fermionic fields Ψ given by (5.18), the commutation relations (5.5) and Eqs.(5.13)–(5.17) we bring the r.h.s. of (5.51) to the form

$$\partial_\mu K_5^\mu(x) = \lim_{\epsilon \rightarrow 0} \frac{g}{2i} (\gamma^\mu \gamma^5)_{ab} [\bar{\Psi}_a(x + \epsilon) \Psi_b(x) + \Psi_b(x + \epsilon) \bar{\Psi}_a(x)] \epsilon^\lambda \partial_\mu K_\lambda(x). \quad (5.52)$$

The product of the Ψ -fields is equal to

$$\begin{aligned} &\bar{\Psi}_a(x + \epsilon) \Psi_b(x) + \Psi_b(x + \epsilon) \bar{\Psi}_a(x) = \\ &=: \bar{\Psi}_a(x + \epsilon) \Psi_b(x) : - : \bar{\Psi}_a(x) \Psi_b(x + \epsilon) : + \frac{1}{\pi i} \frac{(\hat{\epsilon})_{ba}}{\epsilon^2}, \end{aligned} \quad (5.53)$$

where we have dropped an insignificant contribution of the δ -function $\delta(\epsilon^2)$.

Substituting (5.53) in (5.52) we obtain

$$\partial_\mu K_5^\mu(x) = \frac{g}{\pi} \partial_\mu K_\lambda(x) \lim_{\epsilon \rightarrow 0} \frac{1}{2} \text{tr}(\gamma^5 \gamma^\mu \hat{\epsilon}) \frac{\epsilon^\lambda}{\epsilon^2} = \frac{g}{\pi} \partial_\mu K_\lambda(x) \varepsilon^{\mu\nu} \lim_{\epsilon \rightarrow 0} \frac{\epsilon_\nu \epsilon^\lambda}{\epsilon^2}. \quad (5.54)$$

Hence, non-covariant terms are not canceled in the divergence of the quantum axial-vector current that gives $\partial_\mu K_5^\mu(x) \neq 0$. The former breaks explicitly chiral symmetry.

Chiral symmetry can be restored only after averaging over the directions of the 2-vector ϵ [22,25]. This yields

$$\partial_\mu K_5^\mu(x) = \frac{g}{2\pi} \varepsilon^{\mu\nu} \partial_\mu K_\nu(x) = \frac{\alpha - \beta}{2\sqrt{\pi}} \varepsilon^{\mu\nu} \partial_\mu J_\nu(x). \quad (5.55)$$

The r.h.s. of (5.55) vanishes due to the conservation of the axial-vector current of the free massless fermionic field Ψ , $\varepsilon^{\mu\nu} \partial_\mu J_\nu(x) = 0$.

Thereby, Klaiber's method does not provide a cancellation of non-covariant terms in the divergence of the quantum axial-vector current that breaks chiral symmetry explicitly.

Now let us show that the quantum axial-vector current defined by (5.50) does not satisfy the standard relation $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$. Making use of the sequence of actions that we have made for the derivation of the quantum vector current $K^\mu(x)$ we reduce the r.h.s. of (5.50) to the form

$$K_5^\mu(x) = J_5^\mu(x) + \frac{\beta}{\sqrt{\pi}} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^2} J_{5\nu}(x) + \varepsilon^{\mu\nu} \frac{\epsilon_\nu \epsilon_\lambda}{\epsilon^2} \left[-\frac{\alpha}{\sqrt{\pi}} J^\lambda(x) + \frac{g}{\pi} K^\lambda(x) \right]. \quad (5.56)$$

With the relation (5.39) the quantum axial-vector current (5.56) takes the form

$$K_5^\mu(x) = J_5^\mu(x) + \frac{\beta}{\sqrt{\pi}} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^2} J_{5\nu}(x) - \frac{\beta}{\sqrt{\pi}} \varepsilon^{\mu\nu} \frac{\epsilon_\nu \epsilon_\lambda}{\epsilon^2} J^\lambda(x). \quad (5.57)$$

Thus, Klaiber's method for the removal of non-covariant contributions fails for the covariant definition of the quantum axial-vector current. Averaging over all directions of a 2-vector ϵ we arrive at the expression

$$K_5^\mu(x) = \left(1 + \frac{\beta}{\sqrt{\pi}}\right) J_5^\mu(x), \quad (5.58)$$

which does not obey the standard relation $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$ if the quantum vector current $K_\nu(x)$ is given by

$$K^\mu(x) = \left[\left(1 + \frac{\beta}{\sqrt{\pi}}\right) - \frac{\sigma}{g} \frac{\alpha - \beta}{\sqrt{\pi}} \right] J^\mu(x). \quad (5.59)$$

In order to fulfill the relation $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$ one should set $\sigma = 0$. Compared to Klaiber's discussion this condition leads to a substantial narrowing of the class of solutions of the massless Thirring model.

Hence, due to the impossibility to remove non-covariant terms simultaneously in the vector and axial-vector currents and the explicit breakdown of chiral symmetry Klaiber's parameterization based on the definition of the quantum vector current (5.37) seems to be incorrect. Therefore, it cannot be accepted even if for $\sigma = 0$, the case that Klaiber adduced finally in [4], and that has been used then by Coleman for his proof of equivalence between the massive Thirring model and the sine-Gordon model [18].

In turn, if we follow the standard point-splitting prescription [22,25] and average over all directions of a 2-vector ϵ in (5.36) and (5.56), the former, of course, without the contribution of $K_\lambda(x)$, one arrives at the definition of the quantum currents

$$\begin{aligned} j^\mu(x) &= e^{-i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \left(1 + \frac{\alpha + \beta}{\sqrt{\pi}}\right) J_\nu(x), \\ j_5^\mu(x) &= e^{-i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \left(1 + \frac{\alpha + \beta}{\sqrt{\pi}}\right) J_5^\mu(x). \end{aligned} \quad (5.60)$$

which agree completely with the standard relation $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$.

Inserting (5.60) in (5.22) we obtain

$$i\gamma^\mu \partial_\mu \psi(x) = e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{\sqrt{\pi}(\alpha - \beta)}{1 + \frac{\alpha + \beta}{2\sqrt{\pi}}} :j^\mu(x) \gamma_\mu \psi(x): . \quad (5.61)$$

If we define the renormalized quantum vector and axial-vector currents as follows [4]

$$\begin{aligned} \mathcal{J}^\mu(x) &= e^{+i(\alpha - \beta)^2 D^{(+)}(\epsilon)} j^\mu(x) = \left(1 + \frac{\alpha + \beta}{\sqrt{\pi}}\right) J_\nu(x), \\ \mathcal{J}_5^\mu(x) &= e^{+i(\alpha - \beta)^2 D^{(+)}(\epsilon)} j_5^\mu(x) = \left(1 + \frac{\alpha + \beta}{\sqrt{\pi}}\right) J_5^\mu(x), \end{aligned} \quad (5.62)$$

we arrive at the quantum equation of motion

$$i\gamma^\mu \partial_\mu \psi(x) = \frac{\sqrt{\pi}(\alpha - \beta)}{1 + \frac{\alpha + \beta}{2\sqrt{\pi}}} : \mathcal{J}^\mu(x) \gamma_\mu \psi(x) : . \quad (5.63)$$

In terms of the parameter ρ (5.42) the equation of motion (5.63) reads

$$i\gamma^\mu \partial_\mu \psi(x) = 2\pi \frac{\rho^2 - 1}{\rho^2 + 1} : \mathcal{J}^\mu(x) \gamma_\mu \psi(x) : . \quad (5.64)$$

Since the constant factor in front of the product $: \mathcal{J}^\mu(x) \gamma_\mu \psi(x) :$ should be equal to g , we obtain the equation

$$\frac{2\pi}{g} \frac{\rho^2 - 1}{\rho^2 + 1} = 1. \quad (5.65)$$

The solution of this equation defines the parameter ρ :

$$\rho = \sqrt{\frac{2\pi + g}{2\pi - g}}. \quad (5.66)$$

This confirms Klaiber's statement [27] concerning the presence of singularities of the massless Thirring model at $g = \pm 2\pi$ in the g -plane.

We would like to emphasize that in our definition of quantum vector and axial-vector currents the parameter ρ should be only positive. Indeed, for negative ρ in the limit of a free fermionic field theory the renormalized vector and axial-vector currents given by (5.62) vanish. Therefore, for real ρ the coupling constant g should range between $-2\pi < g < 2\pi$.

The main conclusion of this result is the absence of a one-parameter family of solutions for the massless Thirring model with fermionic fields quantized in the chiral symmetric phase. This statement is valid for both Klaiber's parameterization in terms of the parameter σ fixed to the value $\sigma = 0$ and ours given by (5.66).

In order to retain the existence of a one-parameter family of solutions of the quantized version of the massless Thirring model we suggest (i) to renormalize the wave function of the fermionic field $\psi(x) \rightarrow Z_2^{1/2} \psi(x)$

$$i\gamma^\mu \partial_\mu \psi(x) = e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{\sqrt{\pi}(\alpha - \beta)}{1 + \frac{\alpha + \beta}{2\sqrt{\pi}}} :j^\mu(x) \gamma_\mu \psi(x): . \quad (5.67)$$

and denote

$$g = Z_2 e^{i(\alpha - \beta)^2 D^{(+)}(\epsilon)} \frac{\sqrt{\pi}(\alpha - \beta)}{1 + \frac{\alpha + \beta}{2\sqrt{\pi}}} . \quad (5.68)$$

This yields the quantum equation of motion

$$i\gamma^\mu \partial_\mu \psi(x) = g :j^\mu(x) \gamma_\mu \psi(x): \quad (5.69)$$

without any constraints on the parameters. Moreover, by using the definition of the parameters α and β in terms of the parameter ρ one can easily show that the ρ -depending factor can be absorbed by a redefinition of the infrared cut-off regularizing the function $D^{(+)}(\epsilon)$. Thus, this derivation of the quantum equation of motion of the massless Thirring model introduces neither new arbitrary parameters nor additional constraints. The parameters α and β defining a one-parameter family of classical solutions (4.18) are related by $\alpha - \beta = g/\sqrt{\pi}$. In turn, for the quantum solutions (5.1), the parameters α and β obey only the relation (5.26).

In section 11 we show that Klaiber's operator approach applied to the evaluation of correlation functions of massless Thirring fermion fields is inconsistent with non-perturbative ultra-violet renormalizability of the massless Thirring model for any parameters α and β related to each other in the operator formalism. Non-perturbative renormalizability of the massless Thirring model demands contributions induced by local chiral rotations in terms of chiral Jacobians entering with an arbitrary parameter related to a regularization procedure [11].

6 Correlation functions of left(right)handed fermion densities. Path-integral approach

For the derivation of an equivalence between the massive Thirring model and the sine-Gordon model Coleman analysed the following set of n -point correlation functions of left(right)handed fermion densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ [17]

$$\left\langle 0 \left| T \left(\prod_{i=1}^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) \right| 0 \right\rangle . \quad (6.1)$$

This set of correlation functions can be obtained by using the generating functional (1.23) and the generating functional (1.32). This yields

$$\left\langle 0 \left| T \left(\prod_i^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) \right| 0 \right\rangle =$$

$$\begin{aligned}
&= \prod_{i=1}^n \frac{1}{i} \frac{\delta}{\delta J(x_i)} \frac{1}{i} \frac{\delta}{\delta \bar{J}(x_i)} \frac{1}{i} \frac{\delta}{\delta J(y_i)} \frac{1}{i} \frac{\delta}{\delta \bar{J}(y_i)} Z_{\text{Th}}[J, \bar{J}] \Big|_{J = \bar{J} = 0} = \\
&= \prod_{i=1}^n \frac{1}{i} \frac{\delta}{\delta \sigma_+(x_i)} \frac{1}{i} \frac{\delta}{\delta \sigma_-(y_i)} Z_{\text{Th}}[\sigma_-, \sigma_+] \Big|_{\sigma_- = \sigma_+ = 0}.
\end{aligned} \tag{6.2}$$

As a result the n -point correlation functions (6.1) can be represented by the path integral

$$\begin{aligned}
&\langle 0 | T \left(\prod_i^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) | 0 \rangle = \\
&= \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} e^{i \int d^2x \bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x)} \prod_{i=1}^n \bar{\Psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_i) \bar{\Psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_i) \\
&\times \int \mathcal{D}\xi e^{-i \frac{1}{2} \int d^2x \partial_\mu \xi(x) \partial^\mu \xi(x)} \prod_{k=1}^n e^{2i \lambda(\alpha_J) \xi(x_k) - 2i \lambda(\alpha_J) \xi(y_k)}.
\end{aligned} \tag{6.3}$$

For the integration over fermionic degrees of freedom we suggest to consider the following generating functional

$$Z_F[J, \bar{J}] = \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} e^{i \int d^2x [\bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x) + \bar{J}(x) \Psi(x) + \bar{\Psi}(x) J(x)]}, \tag{6.4}$$

where $J(x)$ and $\bar{J}(x) = J^\dagger(x) \gamma^0$ are column and row matrices of external sources of fermion fields $\bar{\Psi}(x)$ and $\Psi(x)$, respectively, with components $J(x) = (J_1(x), J_2(x))$ and $\bar{J}(x) = (J_2^\dagger(x), J_1^\dagger(x))$. The fermionic field $\Psi(x)$ is a column matrix with components $\Psi(x) = (\Psi_1(x), \Psi_2(x))$. It is convenient to rewrite the r.h.s. of (6.4) in terms of the fermionic field components

$$\begin{aligned}
&Z_F[J, \bar{J}] = \\
&= \int \mathcal{D}\Psi_1 \mathcal{D}\Psi_1^\dagger \exp i \int d^2x \left[\Psi_1^\dagger(x) i \left(\frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1} \right) \Psi_1(x) + \Psi_1^\dagger(x) J_2(x) + J_2^\dagger(x) \Psi_1(x) \right] \\
&\times \int \mathcal{D}\Psi_2 \mathcal{D}\Psi_2^\dagger \exp i \int d^2x \left[\Psi_2^\dagger(x) i \left(\frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1} \right) \Psi_2(x) + \Psi_2^\dagger(x) J_1(x) + J_1^\dagger(x) \Psi_2(x) \right].
\end{aligned} \tag{6.5}$$

Integrating out the fermion fields we obtain

$$Z_F[J, \bar{J}] = e^{i \iint d^2x d^2y J_2^\dagger(x) S_{11}(x - y) J_2(y)} e^{i \iint d^2x d^2y J_1^\dagger(x) S_{22}(x - y) J_1(y)}, \tag{6.6}$$

where $S_{11}(x - y)$ and $S_{22}(x - y)$ are the Green functions

$$\begin{aligned}
S_{11}(x - y) &= i \langle 0 | T(\Psi_1(x) \Psi_1^\dagger(y)) | 0 \rangle = \\
&= \frac{1}{2\pi} \frac{1}{v - V - i 0 \cdot \varepsilon(x^0 - y^0)} = \frac{1}{2\pi} \frac{u - U}{(x - y)^2 - i 0}, \\
S_{22}(x - y) &= i \langle 0 | T(\Psi_2(x) \Psi_2^\dagger(y)) | 0 \rangle = \\
&= \frac{1}{2\pi} \frac{1}{u - U - i 0 \cdot \varepsilon(x^0 - y^0)} = \frac{1}{2\pi} \frac{v - V}{(x - y)^2 - i 0},
\end{aligned} \tag{6.7}$$

where $v = x^0 - x^1$, $V = y^0 - y^1$, $u = x^0 + x^1$ and $U = y^0 + y^1$, and $\varepsilon(x^0 - y^0)$ is a sign function.

In terms of $S_{11}(x-y)$ and $S_{22}(x-y)$ the covariant Green function $S_F(x-y)$ is defined by

$$\begin{aligned} S_F(x-y) &= i\langle 0 | T(\psi(x)\bar{\psi}(y)) | 0 \rangle = \\ &= \begin{pmatrix} 0 & S_{11}(x-y) \\ S_{22}(x-y) & 0 \end{pmatrix} = \frac{1}{2\pi} \frac{1}{(x-y)^2 - i0} \begin{pmatrix} 0 & u-U \\ v-V & 0 \end{pmatrix} = \\ &= \frac{1}{2\pi} \frac{1}{(x-y)^2 - i0} [\gamma^0(x^0 - y^0) - \gamma^1(x^1 - y^1)] = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0}. \end{aligned} \quad (6.8)$$

For the evaluation of the Green functions (6.7) we have used expansions of the fermion fields $\Psi_1(x)$ and $\Psi_2(x)$ into plane waves [6]:

$$\begin{aligned} \Psi_1(x) &= \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2\pi}} \theta(+k^1) \left[a(k^1) e^{-ik^0 x^0 + ik^1 x^1} + b^\dagger(k^1) e^{ik^0 x^0 - ik^1 x^1} \right], \\ \Psi_2(x) &= \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{2\pi}} \theta(-k^1) \left[a(k^1) e^{-ik^0 x^0 + ik^1 x^1} - b^\dagger(k^1) e^{ik^0 x^0 - ik^1 x^1} \right], \end{aligned} \quad (6.9)$$

where $a(k^1)(a^\dagger(k^1))$ and $b(k^1)(b^\dagger(k^1))$ are annihilation (creation) operators of fermions and anti-fermions with momentum k^1 .

The evaluation of the Green functions (6.7) runs in the way

$$\begin{aligned} S_{11}(x-y) &= i\langle 0 | T(\Psi_1(x)\Psi_1^\dagger(y)) | 0 \rangle = \\ &= i\theta(x^0 - y^0) \langle 0 | \Psi_1(x)\Psi_1^\dagger(y) | 0 \rangle - i\theta(y^0 - x^0) \langle 0 | \Psi_1^\dagger(y)\Psi_1(x) | 0 \rangle = \\ &= i\theta(x^0 - y^0) \int_{-\infty}^{\infty} \frac{dk^1}{2\pi} \theta(+k^1) e^{-ik^0(x^0 - y^0) + ik^1(x^1 - y^1)} \\ &\quad - i\theta(y^0 - x^0) \int_0^{\infty} \frac{dk^1}{2\pi} \theta(+k^1) e^{ik^0(x^0 - y^0) - ik^1(x^1 - y^1)} = \\ &= i\theta(x^0 - y^0) \int_0^{\infty} \frac{dk^1}{2\pi} e^{-ik^0(x^0 - y^0) + ik^1(x^1 - y^1)} \\ &\quad - i\theta(y^0 - x^0) \int_0^{\infty} \frac{dk^1}{2\pi} e^{ik^0(x^0 - y^0) - ik^1(x^1 - y^1)} = \\ &= \theta(x^0 - y^0) \frac{1}{2\pi} \frac{1}{v - V - i0} + \theta(y^0 - x^0) \frac{1}{2\pi} \frac{1}{v - V + i0} = \\ &= \theta(x^0 - y^0) \frac{1}{2\pi} \left(\text{P} \frac{1}{v - V} + \pi i \delta(v - V) \right) \\ &\quad + \theta(y^0 - x^0) \frac{1}{2\pi} \left(\text{P} \frac{1}{v - V} - \pi i \delta(v - V) \right) = \\ &= \frac{1}{2\pi} \left(\text{P} \frac{1}{v - V} + \pi i \varepsilon(x^0 - y^0) \delta(v - V) \right) = \frac{1}{2\pi} \frac{1}{v - V - i0 \cdot \varepsilon(x^0 - y^0)}, \\ S_{22}(x-y) &= i\langle 0 | T(\Psi_2(x)\Psi_2^\dagger(y)) | 0 \rangle = \\ &= i\theta(x^0 - y^0) \langle 0 | \Psi_2(x)\Psi_2^\dagger(y) | 0 \rangle - i\theta(y^0 - x^0) \langle 0 | \Psi_2^\dagger(y)\Psi_2(x) | 0 \rangle = \\ &= i\theta(x^0 - y^0) \int_{-\infty}^{\infty} \frac{dk^1}{2\pi} \theta(-k^1) e^{-ik^0(x^0 - y^0) + ik^1(x^1 - y^1)} \end{aligned}$$

$$\begin{aligned}
& -i\theta(y^0 - x^0) \int_0^\infty \frac{dk^1}{2\pi} \theta(-k^1) e^{ik^0(x^0 - y^0) - ik^1(x^1 - y^1)} = \\
& = i\theta(x^0 - y^0) \int_0^\infty \frac{dk^1}{2\pi} e^{-ik^0(x^0 - y^0) - ik^1(x^1 - y^1)} \\
& -i\theta(y^0 - x^0) \int_0^\infty \frac{dk^1}{2\pi} e^{ik^0(x^0 - y^0) + ik^1(x^1 - y^1)} = \\
& = \theta(x^0 - y^0) \frac{1}{2\pi} \frac{1}{u - U - i0} + \theta(y^0 - x^0) \frac{1}{2\pi} \frac{1}{u - U + i0} = \\
& = \theta(x^0 - y^0) \frac{1}{2\pi} \left(\text{P} \frac{1}{u - U} + \pi i \delta(u - U) \right) \\
& + \theta(y^0 - x^0) \frac{1}{2\pi} \left(\text{P} \frac{1}{u - U} - \pi i \delta(u - U) \right) = \\
& = \frac{1}{2\pi} \left(\text{P} \frac{1}{u - U} + \pi i \varepsilon(x^0 - y^0) \delta(u - U) \right) = \frac{1}{2\pi} \frac{1}{u - U - i0 \cdot \varepsilon(x^0 - y^0)}, \quad (6.10)
\end{aligned}$$

where the symbol P denotes the principle value and $\varepsilon(x^0 - y^0) = \theta(x^0 - y^0) - \theta(y^0 - x^0)$.

The Green functions $S_{11}(x - y)$ and $S_{22}(x - y)$ obey the equations

$$\begin{aligned}
i \left(\frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1} \right) S_{11}(x - y) &= -\delta^{(2)}(x - y), \\
i \left(\frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1} \right) S_{22}(x - y) &= -\delta^{(2)}(x - y). \quad (6.11)
\end{aligned}$$

One can show that direct solutions of these equations leads to the expressions given by (6.10).

The correlation function (6.3) can be rewritten as follows

$$\begin{aligned}
& \langle 0 | \text{T} \left(\prod_i^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) | 0 \rangle = \int \mathcal{D}\Psi_1 \mathcal{D}\Psi_1^\dagger \mathcal{D}\Psi_2 \mathcal{D}\Psi_2^\dagger \\
& \times \exp i \int d^2x \left[\Psi_1^\dagger(x) i \left(\frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1} \right) \Psi_1(x) + \Psi_2^\dagger(x) i \left(\frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1} \right) \Psi_2(x) \right] \\
& \times \prod_{i=1}^n \Psi_1^\dagger(x_i) \Psi_2(x_i) \Psi_2^\dagger(y_i) \Psi_1(y_i) \int \mathcal{D}\xi e^{-i \frac{1}{2} \int d^2x \partial_\mu \xi(x) \partial^\mu \xi(x)} \\
& \times \prod_{k=1}^n e^{2i \lambda(\alpha_J) \xi(x_k) - 2i \lambda(\alpha_J) \xi(y_k)}. \quad (6.12)
\end{aligned}$$

Using the generating functional (6.6) the integration over fermionic degrees of freedom gives (see (8.35) of Ref.[6])

$$\begin{aligned}
& \int \mathcal{D}\Psi_1 \mathcal{D}\Psi_1^\dagger \mathcal{D}\Psi_2 \mathcal{D}\Psi_2^\dagger \exp i \int d^2x \left[\Psi_1^\dagger(x) i \left(\frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1} \right) \Psi_1(x) \right. \\
& \left. + \Psi_2^\dagger(x) i \left(\frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1} \right) \Psi_2(x) \right] \prod_{i=1}^n \Psi_1^\dagger(x_i) \Psi_2(x_i) \Psi_2^\dagger(y_i) \Psi_1(y_i) =
\end{aligned}$$

$$= \frac{1}{(2\pi)^{2n}} \frac{\prod_{j < k}^n [-(x_j - x_k)^2 + i0] \prod_{j < k}^n [-(y_j - y_k)^2 + i0]}{\prod_{j=1}^n \prod_{k=1}^n [-(x_j - y_k)^2 + i0]}. \quad (6.13)$$

Now the correlation function (6.12) is equal to

$$\begin{aligned} & \langle 0 | T \left(\prod_i^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) | 0 \rangle = \\ &= \frac{1}{(2\pi)^{2n}} \frac{\prod_{j < k}^n [-(x_j - x_k)^2 + i0] \prod_{j < k}^n [-(y_j - y_k)^2 + i0]}{\prod_{j=1}^n \prod_{k=1}^n [-(x_j - y_k)^2 + i0]} \\ & \times \int \mathcal{D}\xi e^{-i \frac{1}{2} \int d^2x \partial_\mu \xi(x) \partial^\mu \xi(x)} \prod_{k=1}^n e^{2i \lambda(\alpha_J) \xi(x_k) - 2i \lambda(\alpha_J) \xi(y_k)}. \end{aligned} \quad (6.14)$$

For the evaluation of the path integral over the ξ -field we apply the infrared regularization suggested by Coleman [17] and discussed then in [8] (see also [6]):

$$\begin{aligned} & \int \mathcal{D}\xi e^{-i \frac{1}{2} \int d^2x \partial_\mu \xi(x) \partial^\mu \xi(x)} \prod_{k=1}^n e^{i \lambda(\alpha_J) \xi(x_k) - i \lambda(\alpha_J) \xi(y_k)} \\ & \rightarrow \lim_{\mu \rightarrow 0} \int \mathcal{D}\xi e^{-i \frac{1}{2} \int d^2x [\partial_\mu \xi(x) \partial^\mu \xi(x) - \mu^2 \xi^2(x)]} \prod_{k=1}^n e^{i \lambda(\alpha_J) \xi(x_k) - i \lambda(\alpha_J) \xi(y_k)} = \\ & \rightarrow \lim_{\mu \rightarrow 0} \int \mathcal{D}\xi e^{i \frac{1}{2} \int d^2x \xi(x) (\square + \mu^2) \xi(x)} \prod_{k=1}^n e^{2i \lambda(\alpha_J) \xi(x_k) - 2i \lambda(\alpha_J) \xi(y_k)}. \end{aligned} \quad (6.15)$$

The result of the integration over the ξ -field reads (see (2.6) of Ref.[6])

$$\begin{aligned} & \int \mathcal{D}\xi e^{i \frac{1}{2} \int d^2x \xi(x) (\square + \mu^2) \xi(x)} \prod_{k=1}^n e^{2i \lambda(\alpha_J) \xi(x_k) - 2i \lambda(\alpha_J) \xi(y_k)} = \\ &= \exp \left\{ -n 4 \lambda^2(\alpha_J) i \Delta(0) \right\} \exp \left\{ -4 \lambda^2(\alpha_J) \sum_{j < k}^n i \Delta(x_j - x_k) \right. \\ & \left. - 4 \lambda^2(\alpha_J) \sum_{j < k}^n i \Delta(y_j - y_k) + 4 \lambda^2(\alpha_J) \sum_{j=1}^n \sum_{k=1}^n i \Delta(x_j - y_k) \right\}, \end{aligned} \quad (6.16)$$

where the Green function $\Delta(x - y)$ is determined by

$$\Delta(x - y) = i \langle 0 | T(\xi(x) \xi(y)) | 0 \rangle \quad (6.17)$$

and obeys the equation

$$(\square + \mu^2) \Delta(x - y) = \delta^{(2)}(x - y). \quad (6.18)$$

In the limit $\mu \rightarrow 0$ the Green function $\Delta(x - y)$ is given by the expression (for a detailed calculation see Appendix C)

$$i\Delta(x - y) = \frac{1}{4\pi} \ell n[-\mu^2(x - y)^2 + i0]. \quad (6.19)$$

For $(x - y) = 0$ the Green function $i\Delta(0)$ amounts to

$$i\Delta(0) = -\frac{1}{4\pi} \ell n\left(\frac{\Lambda^2}{\mu^2}\right), \quad (6.20)$$

where Λ is an ultra-violet cut-off.

It is important to emphasize that the path integrals over the ξ -field are ill-defined in the infrared region

$$\begin{aligned} & \int \mathcal{D}\xi e^{i\frac{1}{2} \int d^2x \xi(x)(\square + \mu^2) \xi(x)} \prod_{k=1}^p \prod_{j=1}^n e^{2i\lambda(\alpha_J) \xi(x_k) - 2i\lambda(\alpha_J) \xi(y_j)} = \\ & = \exp \left\{ -\frac{1}{2} (p + n) 4\lambda^2(\alpha_J) i\Delta(0) \right\} \exp \left\{ -4\lambda^2(\alpha_J) \sum_{j < k}^p i\Delta(x_j - x_k) \right. \\ & \quad \left. - 4\lambda^2(\alpha_J) \sum_{j < k}^n i\Delta(y_j - y_k) + 4\lambda^2(\alpha_J) \sum_{j=1}^p \sum_{k=1}^n i\Delta(x_j - y_k) \right\} \\ & \sim (\mu^2)^{-(p-n)^2 \lambda^2(\alpha_J)/2\pi} \end{aligned} \quad (6.21)$$

and diverge in the limit $\mu^2 \rightarrow 0$ for $p \neq n$. Such a behaviour of correlation functions with $p \neq n$ is related to the ghost-nature of the ξ -field having an incorrect sign of the kinetic term.

After the integration over the ξ -field the correlation function (6.1) reads

$$\begin{aligned} & \langle 0 | T \left(\prod_i^n \bar{\psi}(x_i) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_i) \bar{\psi}(y_i) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_i) \right) | 0 \rangle = \left(\frac{\Lambda}{2\pi} \right)^{2n} \\ & \times \frac{\prod_{j < k}^n [-\Lambda^2(x_j - x_k)^2 + i0]^{1 + d_{\bar{\psi}\psi}(g, \alpha_J)} \prod_{j < k}^n [-\Lambda^2(y_j - y_k)^2 + i0]^{1 + d_{\bar{\psi}\psi}(g, \alpha_J)}}{\prod_{j=1}^n \prod_{k=1}^n [-\Lambda^2(x_j - y_k)^2 + i0]^{1 + d_{\bar{\psi}\psi}(g, \alpha_J)}}, \end{aligned} \quad (6.22)$$

where $d_{\bar{\psi}\psi}(g, \alpha_J)$ are dynamical dimensions of the left(right)handed fermion densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$. According to Wilson's analysis [28] $d_{\bar{\psi}\psi}(g, \alpha_J)$ is equal to

$$d_{\bar{\psi}\psi}(g, \alpha_J) = -\frac{g/\pi}{1 + \alpha_J g/\pi}. \quad (6.23)$$

The dependence of $d_{\bar{\psi}\psi}(g, \alpha_J)$ on the arbitrary parameter α_J agrees with a one-parameter family of solutions of the massless Thirring model.

For the two-point correlation function we obtain

$$\begin{aligned} & \left\langle 0 \left| T \left(\bar{\psi}(x) \left(\frac{1-\gamma^5}{2} \right) \psi(x) \bar{\psi}(y) \left(\frac{1+\gamma^5}{2} \right) \psi(y) \right) \right| 0 \right\rangle = \\ & = \frac{\Lambda^2}{4\pi^2} [-\Lambda^2(x-y)^2 + i0]^{-1-d_{\bar{\psi}\psi}(g, \alpha_J)}. \end{aligned} \quad (6.24)$$

We would like to emphasize that the correlation functions (6.22) evaluated within the path-integral approach do not depend on an infrared cut-off. This is unlike the correlation functions evaluated within Klaiber's operator formalism. In sections 8 and 9 we consider the evaluation of $2n$ -point Green functions and n -point correlation functions of the left(right)handed fermion densities within Klaiber's operator formalism and discuss the dependence of these expressions on the infrared cut-off.

7 Bosonization of generating functionals $Z_{\text{Th}}[\sigma_-, \sigma_+]$ and $Z_{\text{Th}}[\sigma]$

The expression (6.22) for the n -point correlation function (6.1) can be obtained using the generating functional $Z_{\text{Th}}[\sigma_-, \sigma_+]$ in (1.32). In this section we show that the fermionic degrees of freedom in the generating functionals $Z_{\text{Th}}[\sigma_-, \sigma_+]$ and $Z_{\text{Th}}[\sigma]$ can be fully replaced by bosonic degrees. This means that these generating functionals admit full bosonization.

For this aim we suggest to expand $Z_{\text{Th}}[\sigma_-, \sigma_+]$ in powers of $\sigma_{\mp}$. This yields

$$\begin{aligned} Z_{\text{Th}}[\sigma_-, \sigma_+] &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{i^n}{n!} \frac{i^m}{m!} \prod_{i=1}^n \prod_{j=1}^m \iint d^2x_i d^2y_j \sigma_+(x_i) \sigma_-(y_j) \\ &\times \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\xi \exp i \int d^2z \left\{ \bar{\Psi}(z) i \gamma^\mu \partial_\mu \Psi(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right\} \\ &\times \left[\bar{\Psi}(x_i) \left(\frac{1-\gamma^5}{2} \right) \Psi(x_i) \right] \left[\bar{\Psi}(y_j) \left(\frac{1+\gamma^5}{2} \right) \Psi(y_j) \right] e^{-2i\lambda(\alpha_J) \xi(x_i)} e^{2i\lambda(\alpha_J) \xi(y_j)} = \\ &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{i^n}{n!} \frac{i^m}{m!} \prod_{i=1}^n \prod_{j=1}^m \iint d^2x_i d^2y_j \sigma_+(x_i) \sigma_-(y_j) \\ &\times \left\langle 0 \left| T \left(\left[\bar{\Psi}(x_i) \left(\frac{1-\gamma^5}{2} \right) \Psi(x_i) \right] \left[\bar{\Psi}(y_j) \left(\frac{1+\gamma^5}{2} \right) \Psi(y_j) \right] \right) \right| 0 \right\rangle \\ &\times \int \mathcal{D}\xi \exp \left\{ -i \int d^2z \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) - 2i\lambda(\alpha_J) \xi(x_i) + 2i\lambda(\alpha_J) \xi(y_j) \right\}. \end{aligned} \quad (7.1)$$

The time-ordered vacuum expectation value of products of free fermion fields Ψ and $\bar{\Psi}$ can be represented in terms of a time-ordered vacuum expectation value of products of a massless pseudoscalar (or scalar) field ϑ . Using Eq.(2.10) of Ref.[6] we get

$$\left\langle 0 \left| T \left(\prod_{i=1}^n \prod_{j=1}^m \left[\bar{\Psi}(x_i) \left(\frac{1-\gamma^5}{2} \right) \Psi(x_i) \right] \left[\bar{\Psi}(y_j) \left(\frac{1+\gamma^5}{2} \right) \Psi(y_j) \right] \right) \right| 0 \right\rangle =$$

$$\begin{aligned}
&= \delta_{nm} \frac{1}{(2\pi)^{n+m}} \frac{\prod_{i < k}^n [-(x_i - x_k)^2] \prod_{j < k}^m [-(y_j - y_k)^2]}{\prod_{i=1}^n \prod_{j=1}^m [-(x_i - y_j)^2]} = \\
&= \prod_{i=1}^n \prod_{j=1}^m \left\langle 0 \left| T \left(\frac{\Lambda}{2\pi} e^{i\sqrt{4\pi} \vartheta(x_i)} \frac{\Lambda}{2\pi} e^{-i\sqrt{4\pi} \vartheta(y_j)} \right) \right| 0 \right\rangle, \tag{7.2}
\end{aligned}$$

where δ_{nm} is the Kronecker symbol and Λ is an ultra-violet cut-off.

Substituting (7.2) and (7.1) we get

$$\begin{aligned}
Z_{\text{Th}}[\sigma_-, \sigma_+] &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{i^n}{n!} \frac{i^m}{m!} \prod_{i=1}^n \prod_{j=1}^m \iint d^2 x_i d^2 y_j \sigma_+(x_i) \sigma_-(y_j) \\
&\times \left\langle 0 \left| T \left(\frac{\Lambda}{2\pi} e^{i\sqrt{4\pi} \vartheta(x_i)} \frac{\Lambda}{2\pi} e^{-i\sqrt{4\pi} \vartheta(y_j)} \right) \right| 0 \right\rangle \\
&\times \int \mathcal{D}\xi \exp \left\{ -i \int d^2 z \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) - 2i\lambda(\alpha_J) \xi(x_i) + 2i\lambda(\alpha_J) \xi(y_j) \right\}. \tag{7.3}
\end{aligned}$$

Since the vacuum expectation value can be represented in terms of a path integral over the ϑ -field (see Eq.(2.5) of Ref.[6]), the generating functional (7.3) takes the form

$$\begin{aligned}
Z_{\text{Th}}[\sigma_-, \sigma_+] &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{i^n}{n!} \frac{i^m}{m!} \prod_{i=1}^n \prod_{j=1}^m \iint d^2 x_i d^2 y_j \sigma_+(x_i) \sigma_-(y_j) \\
&\times \int \mathcal{D}\vartheta \mathcal{D}\xi \exp i \int d^2 z \left\{ \frac{1}{2} \partial_\mu \vartheta(z) \partial^\mu \vartheta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right\} \\
&\times \frac{\Lambda}{2\pi} e^{i\sqrt{4\pi} \vartheta(x_i)} \frac{\Lambda}{2\pi} e^{-i\sqrt{4\pi} \vartheta(y_j)} e^{-2i\lambda(\alpha_J) \xi(x_i) + 2i\lambda(\alpha_J) \xi(y_j)}. \tag{7.4}
\end{aligned}$$

Summing up the series we get

$$\begin{aligned}
Z_{\text{Th}}[\sigma_-, \sigma_+] &= \int \mathcal{D}\vartheta \mathcal{D}\xi \exp i \int d^2 z \left\{ \frac{1}{2} \partial_\mu \vartheta(z) \partial^\mu \vartheta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right. \\
&+ \sigma_-(z) \frac{\Lambda}{2\pi} e^{-i\sqrt{4\pi} \vartheta(z)} + 2i\lambda(\alpha_J) \xi(z) + \sigma_+(z) \frac{\Lambda}{2\pi} e^{+i\sqrt{4\pi} \vartheta(z)} - 2i\lambda(\alpha_J) \xi(z) \left. \right\}. \tag{7.5}
\end{aligned}$$

In order to remove unphysical degrees of freedom from the path integral (7.5) we suggest to make a change of variables

$$\begin{aligned}
\Theta(x) &= \frac{\sqrt{4\pi} \vartheta(x) - 2\lambda(\alpha_J) \xi(x)}{\sqrt{4\pi - 4\lambda^2(\alpha_J)}}, \\
\Phi(x) &= \frac{2\lambda(\alpha_J) \vartheta(x) - \sqrt{4\pi} \xi(x)}{\sqrt{4\pi - 4\lambda^2(\alpha_J)}} \tag{7.6}
\end{aligned}$$

Now the generating functional (7.5) reads

$$Z_{\text{Th}}[\sigma_-, \sigma_+] = \int \mathcal{D}\Theta \mathcal{D}\Phi \exp i \int d^2 z \left\{ \frac{1}{2} \partial_\mu \Theta(z) \partial^\mu \Theta(z) - \frac{1}{2} \partial_\mu \Phi(z) \partial^\mu \Phi(z) \right\}$$

$$\begin{aligned}
& +\sigma_-(z) \frac{\Lambda}{2\pi} \exp \left(-i\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(z) \right) \\
& +\sigma_+(z) \frac{\Lambda}{2\pi} \exp \left(+i\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(z) \right) \Big\}, \tag{7.7}
\end{aligned}$$

where we have used the relation

$$\sqrt{4\pi - 4\lambda^2(\alpha_J)} = \sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \tag{7.8}$$

with $d_{\bar{\psi}\psi}(g,\alpha_J)$ given by (6.23).

Integrating out the Φ -field we define the generating functional $Z_{\text{Th}}[\sigma_-, \sigma_+]$ by the path integral over the Θ -field only

$$\begin{aligned}
Z_{\text{Th}}[\sigma_-, \sigma_+] &= \int \mathcal{D}\Theta \exp i \int d^2z \left\{ \frac{1}{2} \partial_\mu \Theta(z) \partial^\mu \Theta(z) \right. \\
& +\sigma_-(z) \frac{\Lambda}{2\pi} \exp \left(-i\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(z) \right) \\
& \left. +\sigma_+(z) \frac{\Lambda}{2\pi} \exp \left(+i\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(z) \right) \right\} \tag{7.9}
\end{aligned}$$

It is important to accentuate that the generating functional (7.9) is in terms of physical degrees of freedom only. It is easy to verify that substituting the generating functional (7.9) and integrating over the Θ -field one obtains the result (6.22).

Setting $\sigma_-(z) = \sigma_+(z) = \sigma(z)$ in (7.9) we obtain the bosonized version of the generating functional $Z_{\text{Th}}[\sigma]$. It reads

$$\begin{aligned}
Z_{\text{Th}}[\sigma] &= \int \mathcal{D}\Theta \exp i \int d^2z \left\{ \frac{1}{2} \partial_\mu \Theta(z) \partial^\mu \Theta(z) \right. \\
& \left. +\sigma(z) \frac{\Lambda}{\pi} \cos \left(\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(z) \right) \right\}. \tag{7.10}
\end{aligned}$$

Using the generating functional (7.10) we are able to evaluate the two-point correlation function of scalar fermion densities

$$\begin{aligned}
\langle 0 | T(\bar{\psi}(x)\psi(x)\bar{\psi}(y)\psi(y)) | 0 \rangle &= \frac{1}{i} \frac{\delta}{\delta\sigma(x)} \frac{1}{i} \frac{\delta}{\delta\sigma(y)} Z_{\text{Th}}[\sigma] \Big|_{\sigma=0} = \\
&= \frac{\Lambda^2}{\pi^2} \int \mathcal{D}\Theta \exp i \int d^2z \left\{ \frac{1}{2} \partial_\mu \Theta(z) \partial^\mu \Theta(z) \right\} \\
&\times \cos \left(\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(x) \right) \cos \left(\sqrt{4\pi(1+d_{\bar{\psi}\psi}(g,\alpha_J))} \Theta(y) \right). \tag{7.11}
\end{aligned}$$

Integrating over the Θ -field we get

$$\langle 0 | T(\bar{\psi}(x)\psi(x)\bar{\psi}(y)\psi(y)) | 0 \rangle = \frac{\Lambda^2}{2\pi^2} [-\Lambda^2(x-y)^2 + i0]^{-1-d_{\bar{\psi}\psi}(g,\alpha_J)}. \tag{7.12}$$

The same result for the two-point correlation function (7.12) can be derived from (6.24).

Setting $\sigma(z) = -m$, where m has the meaning of the mass of Thirring fermions, the generating functional (7.10) reduces to the partition function of the sine–Gordon (SG) model

$$\begin{aligned} Z_{\text{Th}}[\sigma] \Big|_{\sigma=-m} &= Z_{\text{SG}} = \\ &= \int \mathcal{D}\Theta \exp i \int d^2z \left\{ \frac{1}{2} \partial_\mu \Theta(z) \partial^\mu \Theta(z) - m \frac{\Lambda}{\pi} \cos \left(\sqrt{4\pi (1 + d_{\bar{\psi}\psi}(g, \alpha_J))} \Theta(z) \right) \right\}. \end{aligned} \quad (7.13)$$

Recall that the sine–Gordon model is described by the Lagrangian (1.28). A positive sign for the interaction in the r.h.s. of (7.13) one can get by the shift

$$\Theta(z) \rightarrow \Theta(z) + \pi \sqrt{\frac{1}{4\pi} \frac{1}{1 + d_{\bar{\psi}\psi}(g, \alpha_J)}}. \quad (7.14)$$

Matching the Lagrangian of the sine–Gordon model (1.29) with the effective Lagrangian in the exponent of the r.h.s. of the generating functional (7.13) we obtain the relations between the sine–Gordon parameters $\bar{\alpha}$ and $\bar{\beta}$ and the Thirring model parameters

$$\begin{aligned} \frac{\bar{\beta}^2}{4\pi} &= 1 + d_{\bar{\psi}\psi}(g, \alpha_J), \\ \bar{\alpha} &= 4m\Lambda \frac{\bar{\beta}^2}{4\pi} = 4m\Lambda (1 + d_{\bar{\psi}\psi}(g, \alpha_J)). \end{aligned} \quad (7.15)$$

Substituting the dynamical dimension $d_{\bar{\psi}\psi}(g, \alpha_J)$ given by (6.22) we get

$$\begin{aligned} \frac{\bar{\beta}^2}{4\pi} &= \frac{1 + (\alpha_J - 1)g/\pi}{1 + \alpha_J g/\pi} \Big|_{\alpha_J=1} = \frac{1}{1 + g/\pi}, \\ \bar{\alpha} &= 4m\Lambda \frac{1 + (\alpha_J - 1)g/\pi}{1 + \alpha_J g/\pi} \Big|_{\alpha_J=1} = \frac{4m\Lambda}{1 + g/\pi}. \end{aligned} \quad (7.16)$$

Our results obtained in this section agree with Naón’s calculations [8] up to the uncertainties defined by the parameter α_J and related to regularizations of chiral Jacobians.

8 Two–point Green function of the massless Thirring fermion field. The path–integral approach

In this section we discuss the evaluation of the two–point Green function of the Thirring fermionic field $S_{\text{F}}(x, y) = i \langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle$ in the path–integral approach suggested in Refs.[7,8].

In terms of the generating functional of Green functions given by (1.11) the two–point Green function $S_{\text{F}}(x, y)$ is determined by

$$S_{\text{F}}(x, y) = i \langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle = i \frac{\delta}{\delta \bar{J}(x)} \frac{\delta}{\delta J(y)} Z_{\text{Th}}[J, \bar{J}] \Big|_{J = \bar{J} = 0}. \quad (8.1)$$

Using the path-integral representation of the generating functional (1.23) the two-point Green function $S_F(x, y)$ is given by

$$\begin{aligned}
S_F(x, y) &= i \langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle = i \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi \\
&\times \exp i \int d^2 z \left[\bar{\Psi}(z) i \gamma^\mu \partial_\mu \Psi(z) + \frac{1}{2} \partial_\mu \eta(z) \partial^\mu \eta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right] \\
&\left[\left(\frac{1 + \gamma^5}{2} \right) \Psi(x) e^{i \sqrt{g} \eta(x) + i \lambda(\alpha_J) \xi(x)} + \left(\frac{1 - \gamma^5}{2} \right) \Psi(x) e^{i \sqrt{g} \eta(x) - i \lambda(\alpha_J) \xi(x)} \right] \\
&\times \left[\bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) e^{-i \sqrt{g} \eta(y) + i \lambda(\alpha_J) \xi(y)} + \bar{\Psi}(y) \left(\frac{1 - \gamma^5}{2} \right) e^{-i \sqrt{g} \eta(y) - i \lambda(\alpha_J) \xi(y)} \right].
\end{aligned} \tag{8.2}$$

In a more convenient form the r.h.s. of (8.2) can be written as

$$\begin{aligned}
S_F(x, y) &= \int \mathcal{D}\eta \mathcal{D}\xi \exp \left\{ i \int d^2 z \left[\frac{1}{2} \partial_\mu \eta(z) \partial^\mu \eta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right] \right\} \\
&\times \left\{ i \langle 0 | T \left(\left(\frac{1 + \gamma^5}{2} \right) \Psi(x) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \right) | 0 \rangle \right. \\
&\quad \times e^{+i \sqrt{g} \eta(x) + i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) + i \lambda(\alpha_J) \xi(y)} \\
&\quad + i \langle 0 | T \left(\left(\frac{1 - \gamma^5}{2} \right) \Psi(x) \bar{\Psi}(y) \left(\frac{1 - \gamma^5}{2} \right) \right) | 0 \rangle \\
&\quad \times e^{+i \sqrt{g} \eta(x) - i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) - i \lambda(\alpha_J) \xi(y)} \\
&\quad + i \langle 0 | T \left(\left(\frac{1 + \gamma^5}{2} \right) \Psi(x) \bar{\Psi}(y) \left(\frac{1 - \gamma^5}{2} \right) \right) | 0 \rangle \\
&\quad \times e^{+i \sqrt{g} \eta(x) + i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) - i \lambda(\alpha_J) \xi(y)} \\
&\quad \left. + i \langle 0 | T \left(\left(\frac{1 - \gamma^5}{2} \right) \Psi(x) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \right) | 0 \rangle \right. \\
&\quad \left. \times e^{+i \sqrt{g} \eta(x) - i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) + i \lambda(\alpha_J) \xi(y)} \right\}.
\end{aligned} \tag{8.3}$$

The Green function of the free massless fermionic field $\Psi(x)$ is defined by (1.10). By using this expression the r.h.s. of (8.3) can be recast into the form

$$\begin{aligned}
S_F(x, y) &= \int \mathcal{D}\eta \mathcal{D}\xi \exp \left\{ i \int d^2 z \left[\frac{1}{2} \partial_\mu \eta(z) \partial^\mu \eta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right] \right\} \\
&\times \left\{ \left(\frac{1 + \gamma^5}{2} \right) \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x - y)^2 - i 0} e^{+i \sqrt{g} \eta(x) + i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) - i \lambda(\alpha_J) \xi(y)} \right. \\
&\quad \left. + \left(\frac{1 - \gamma^5}{2} \right) \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x - y)^2 - i 0} e^{+i \sqrt{g} \eta(x) - i \lambda(\alpha_J) \xi(x) - i \sqrt{g} \eta(y) + i \lambda(\alpha_J) \xi(y)} \right\}.
\end{aligned} \tag{8.4}$$

Since the integrands are identical up to the transformation $\xi \rightarrow -\xi$, the two-point Green function is defined by

$$S_F(x, y) = i \langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x - y)^2 - i 0}$$

$$\begin{aligned}
& \times \int \mathcal{D}\eta \mathcal{D}\xi \exp \left\{ i \int d^2z \left[\frac{1}{2} \partial_\mu \eta(z) \partial^\mu \eta(z) - \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) \right] \right. \\
& \left. + i\sqrt{g} \eta(x) + i\lambda(\alpha_J) \xi(x) - i\sqrt{g} \eta(y) - i\lambda(\alpha_J) \xi(y) \right\}
\end{aligned} \tag{8.5}$$

The integration over η and ξ fields leads to

$$\begin{aligned}
& \int \mathcal{D}\eta \exp \left\{ i \int d^2z \frac{1}{2} \partial_\mu \eta(z) \partial^\mu \eta(z) + i\sqrt{g} \eta(x) - i\sqrt{g} \eta(y) \right\} \\
& \rightarrow \int \mathcal{D}\eta \exp \left\{ - i \int d^2z \frac{1}{2} \eta(z) (\square + \mu^2) \eta(z) + i\sqrt{g} \eta(x) - i\sqrt{g} \eta(y) \right\} = \\
& = \exp \left\{ g i \Delta(0) - g i \Delta(x-y) \right\} = \exp \left\{ - \frac{g}{4\pi} \ell n \frac{\Lambda^2}{\mu^2} - \frac{g}{4\pi} \ell n [-\mu^2(x-y)^2 + i0] \right\} = \\
& = \exp \left\{ - \frac{g}{4\pi} \ell n [-\Lambda^2(x-y)^2 + i0] \right\}, \\
& \int \mathcal{D}\xi \exp \left\{ - i \int d^2z \frac{1}{2} \partial_\mu \xi(z) \partial^\mu \xi(z) + i\lambda(\alpha_J) \xi(x) - i\lambda(\alpha_J) \xi(y) \right\} \\
& \rightarrow \int \mathcal{D}\xi \exp \left\{ i \int d^2z \frac{1}{2} \xi(z) (\square + \mu^2) \xi(z) + i\lambda(\alpha_J) \xi(x) - i\lambda(\alpha_J) \xi(y) \right\} = \\
& = \exp \left\{ - \lambda^2(\alpha_J) i \Delta(0) + \lambda^2(\alpha_J) i \Delta(x-y) \right\} = \\
& = \exp \left\{ \frac{\lambda^2(\alpha_J)}{4\pi} \ell n [-\Lambda^2(x-y)^2 + i0] \right\}.
\end{aligned} \tag{8.6}$$

Thus, the two-point Green function of the Thirring fermionic field is equal to

$$\begin{aligned}
S_F(x, y) &= i \langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0} \\
&\times \exp \left\{ \frac{\lambda^2(\alpha_J)}{4\pi} \ell n [-\Lambda^2(x-y)^2 + i0] - \frac{g}{4\pi} \ell n [-\Lambda^2(x-y)^2 + i0] \right\} = \\
&= \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0} [-\Lambda^2(x-y)^2 + i0]^{-(g - \lambda^2(\alpha_J))/4\pi} = \\
&= \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0} [-\Lambda^2(x-y)^2 + i0]^{-d_\psi(g, \alpha_J)},
\end{aligned} \tag{8.7}$$

where $d_\psi(g, \alpha_J)$, the dynamical dimension of the massless Thirring fermionic field [21]

$$d_\psi(g, \alpha_J) = \frac{g - \lambda^2(\alpha_J)}{4\pi} = \frac{g^2}{4\pi^2} \frac{\alpha_J}{1 + \alpha_J \frac{g}{\pi}}. \tag{8.8}$$

It depends on α_J as well as the dynamical dimension of the left(right)handed fermion densities (6.23).

The two-point Green function (8.7) describes the Green function of massless fermion fields quantized in the chiral symmetric phase. We emphasize that the expression (8.7) does not depend on the infrared cut-off.

9 $2n$ -point Green functions of fermion fields. Path-integral approach and Klaiber's operator formalism

In the massless Thirring model there are two irreducible $2n$ -point Green functions. They read

$$\begin{aligned} G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_1(x_1) \dots \psi_1(x_n) \psi_1^\dagger(y_1) \dots \psi_1^\dagger(y_n)) | 0 \rangle, \\ G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_2(x_1) \dots \psi_2(x_n) \psi_2^\dagger(y_1) \dots \psi_2^\dagger(y_n)) | 0 \rangle. \end{aligned} \quad (9.1)$$

For the evaluation of these Green functions we apply the generating functional (1.11) given by

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\psi_1 \mathcal{D}\psi_1^\dagger \mathcal{D}\psi_2 \mathcal{D}\psi_2^\dagger \\ &\times \exp i \int d^2x \left[\bar{\psi}(x) i \gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \right. \\ &\left. + \psi_1^\dagger(x) J_2(x) + \psi_2^\dagger(x) J_1(x) + J_2^\dagger(x) \psi_1(x) + J_1^\dagger(x) \psi_2(x) \right], \end{aligned} \quad (9.2)$$

After the set of transformations given by (1.12) – (1.20) the generating functional (9.2) acquires the form

$$\begin{aligned} Z_{\text{Th}}[J, \bar{J}] &= \int \mathcal{D}\Psi_1 \mathcal{D}\Psi_1^\dagger \mathcal{D}\Psi_2 \mathcal{D}\Psi_2^\dagger \mathcal{D}\eta \mathcal{D}\xi \\ &\times \exp i \int d^2x \left\{ \bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\ &+ \Psi_1^\dagger(x) J_2(x) e^{-i\sqrt{g}\eta(x) - i\lambda(\alpha_J)\xi(x)} + \Psi_2^\dagger(x) J_1(x) e^{-i\sqrt{g}\eta(x) + i\lambda(\alpha_J)\xi(x)} \\ &\left. + J_2^\dagger(x) \Psi_1(x) e^{i\sqrt{g}\eta(x) + i\lambda(\alpha_J)\xi(x)} + J_1^\dagger(x) \Psi_2(x) e^{i\sqrt{g}\eta(x) - i\lambda(\alpha_J)\xi(x)} \right\}. \end{aligned} \quad (9.3)$$

In terms of the generating functional $Z_{\text{Th}}[J, \bar{J}]$ the Green functions (9.1) are defined by

$$\begin{aligned} G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_1(x_1) \dots \psi_1(x_n) \psi_1^\dagger(y_1) \dots \psi_1^\dagger(y_n)) | 0 \rangle = \\ &= \frac{\delta}{\delta J_2^\dagger(x_1)} \dots \frac{\delta}{\delta J_2^\dagger(x_n)} \frac{\delta}{\delta J_2(y_1)} \dots \frac{\delta}{\delta J_2(y_n)} Z_{\text{Th}}[J, \bar{J}] \Big|_{J_1 = J_2 = J_1^\dagger = J_2^\dagger = 0}, \\ G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_2(x_1) \dots \psi_2(x_n) \psi_2^\dagger(y_1) \dots \psi_2^\dagger(y_n)) | 0 \rangle = \\ &= \frac{\delta}{\delta J_1^\dagger(x_1)} \dots \frac{\delta}{\delta J_1^\dagger(x_n)} \frac{\delta}{\delta J_1(y_1)} \dots \frac{\delta}{\delta J_1(y_n)} Z_{\text{Th}}[J, \bar{J}] \Big|_{J_1 = J_2 = J_1^\dagger = J_2^\dagger = 0}. \end{aligned} \quad (9.4)$$

Substituting (9.3) in (9.4) we get

$$\begin{aligned} G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle \\ &\times \int \mathcal{D}\eta \mathcal{D}\xi \exp i \int d^2x \left\{ \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right\} \\ &\times \prod_{j=1}^n e^{i\sqrt{g}\eta(x_j) + i\lambda(\alpha_J)\xi(x_j)} e^{-i\sqrt{g}\eta(y_j) - i\lambda(\alpha_J)\xi(y_j)}, \end{aligned}$$

$$\begin{aligned}
G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle \\
&\times \int \mathcal{D}\eta \mathcal{D}\xi \exp i \int d^2x \left\{ \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right\} \\
&\times \prod_{j=1}^n e^{i\sqrt{g} \eta(x_j) - i\lambda(\alpha_J) \xi(x_j)} e^{-i\sqrt{g} \eta(y_j) + i\lambda(\alpha_J) \xi(y_j)}
\end{aligned} \tag{9.5}$$

Integration over η and ξ fields gives

$$\begin{aligned}
G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle \\
&\times \frac{\prod_{j < k}^n [-\Lambda^2(x_j - x_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi} \prod_{j < k}^n [-\Lambda^2(y_j - y_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi}}{\prod_{j=1}^n \prod_{k=1}^n [-\Lambda^2(x_j - y_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi}}, \\
G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle \\
&\times \frac{\prod_{j < k}^n [-\Lambda^2(x_j - x_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi} \prod_{j < k}^n [-\Lambda^2(y_j - y_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi}}{\prod_{j=1}^n \prod_{k=1}^n [-\Lambda^2(x_j - y_k)^2 + i0]^{(g - \lambda^2(\alpha_J))/4\pi}}.
\end{aligned} \tag{9.6}$$

In order to compare the expressions (9.8) with Klaiber's result [4] we suggest to represent them in the form

$$\begin{aligned}
G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle e^{iF(x, y)}, \\
G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle e^{iF(x, y)},
\end{aligned} \tag{9.7}$$

where the exponential $e^{iF(x, y)}$ is determined by

$$\begin{aligned}
e^{iF(x, y)} &= e^{-in \frac{g - \lambda^2(\alpha_J)}{4}} \left(\frac{1}{\Lambda} \right)^n \frac{g - \lambda^2(\alpha_J)}{2\pi} \\
&\times \prod_{j < k}^n (v_j - v_k - i0 \cdot \varepsilon(x_j^0 - x_k^0)) \frac{g - \lambda^2(\alpha_J)}{4\pi} (u_j - u_k - i0 \cdot \varepsilon(x_j^0 - x_k^0)) \frac{g}{4\pi} \\
&\times \prod_{j < k}^n (V_j - V_k - i0 \cdot \varepsilon(y_j^0 - y_k^0)) \frac{g - \lambda^2(\alpha_J)}{4\pi} (U_j - U_k - i0 \cdot \varepsilon(y_j^0 - y_k^0)) \frac{g - \lambda^2(\alpha_J)}{4\pi} \\
&\times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{v_j - V_k - i0 \cdot \varepsilon(x_j^0 - y_k^0)} \right) \frac{g - \lambda^2(\alpha_J)}{4\pi} \left(\frac{1}{u_j - U_k - i0 \cdot \varepsilon(x_j^0 - y_k^0)} \right) \frac{g - \lambda^2(\alpha_J)}{4\pi}.
\end{aligned} \tag{9.8}$$

In Klaiber's approach this contribution reads (see Eq.(VII.2) of Ref.[4])

$$e^{iF(x, y)_K} = e^{-in \frac{a+b}{4}} \left(\frac{1}{\mu} \right)^n \frac{a+b}{2\pi}$$

$$\begin{aligned}
& \times \prod_{j < k}^n (v_j - v_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))^{\frac{a+b-2\lambda}{4\pi}} (u_j - u_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))^{\frac{a+b+2\lambda}{4\pi}} \\
& \times \prod_{j < k}^n (V_j - V_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))^{\frac{a+b-2\lambda}{4\pi}} (U_j - U_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))^{\frac{a+b+2\lambda}{4\pi}} \\
& \times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{v_j - V_k - i 0 \cdot \varepsilon(x_j^0 - y_k^0)} \right)^{\frac{a+b-2\lambda}{4\pi}} \left(\frac{1}{u_j - U_k - i 0 \cdot \varepsilon(x_j^0 - y_k^0)} \right)^{\frac{a+b+2\lambda}{4\pi}},
\end{aligned} \tag{9.9}$$

where μ is an infrared cut-off and the parameters a , b and λ are defined by (see Eq.(IV.25) of Ref.[4])

$$\begin{aligned}
a &= \alpha^2 + 2\sqrt{\pi}\alpha, \\
b &= \beta^2 + 2\sqrt{\pi}\beta, \\
\lambda &= \alpha\beta + \sqrt{\pi}\alpha + \sqrt{\pi}\beta.
\end{aligned} \tag{9.10}$$

Invariance of the correlation function (9.10) under Lorentz transformations imposes the constraint $\lambda = 0$ [4] (see also the discussion in section 5). This reduces the correlation function (9.9) to the form

$$\begin{aligned}
e^{iF(x,y)_K} &= e^{-i n \frac{a+b}{4}} \left(\frac{1}{\mu} \right)^n \frac{a+b}{2\pi} \\
& \times \prod_{j < k}^n (v_j - v_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))^{\frac{a+b}{4\pi}} (u_j - u_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))^{\frac{a+b}{4\pi}} \\
& \times \prod_{j < k}^n (V_j - V_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))^{\frac{a+b}{4\pi}} (U_j - U_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))^{\frac{a+b}{4\pi}} \\
& \times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{v_j - V_k - i 0 \cdot \varepsilon(x_j^0 - y_k^0)} \right)^{\frac{a+b}{4\pi}} \left(\frac{1}{u_j - U_k - i 0 \cdot \varepsilon(x_j^0 - y_k^0)} \right)^{\frac{a+b}{4\pi}},
\end{aligned} \tag{9.11}$$

where $a+b = (\alpha - \beta)^2$ valid at $\lambda = 0$. For further discussion it is convenient to define the exponential (9.11) relative to the ultra-violate cut-off

$$\begin{aligned}
e^{iF(x,y)_K} &= e^{-i n \frac{a+b}{4}} \left(\frac{\Lambda}{\mu} \right)^n \frac{a+b}{2\pi} \\
& \times \prod_{j < k}^n [\Lambda(v_j - v_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))]^{\frac{a+b}{4\pi}} [\Lambda(u_j - u_k - i 0 \cdot \varepsilon(x_j^0 - x_k^0))]^{\frac{a+b}{4\pi}} \\
& \times \prod_{j < k}^n [\Lambda(V_j - V_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))]^{\frac{a+b}{4\pi}} [\Lambda(U_j - U_k - i 0 \cdot \varepsilon(y_j^0 - y_k^0))]^{\frac{a+b}{4\pi}}
\end{aligned}$$

$$\times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{[\Lambda(v_j - V_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{\frac{a+b}{4\pi}} \left(\frac{1}{[\Lambda(u_j - U_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{\frac{a+b}{4\pi}}. \quad (9.12)$$

It is seen that Klaiber's correlation function is singular in the infrared limit $\mu \rightarrow 0$.

The dynamical dimension of the massless fermionic field in terms of Klaiber's parameters is determined by

$$d_\psi(g, a, b) = \frac{a+b}{4\pi}. \quad (9.13)$$

Vacuum expectation values of the time-ordered products of the fields Ψ_1 and Ψ_2 can be evaluated by means of the generating functional $Z_F[J, \bar{J}]$ (6.6). The results read

$$\begin{aligned} \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle &= \\ &= \frac{1}{(2\pi i)^n} \frac{\prod_{j < k}^n (u_j - u_k - i0 \cdot \varepsilon(x_j^0 - x_k^0)) \prod_{j < k}^n (U_j - U_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))}{\prod_{j=1}^n \prod_{k=1}^n (u_j - U_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))}, \\ \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle &= \\ &= \frac{1}{(2\pi i)^n} \frac{\prod_{j < k}^n (v_j - v_k - i0 \cdot \varepsilon(x_j^0 - x_k^0)) \prod_{j < k}^n (V_j - V_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))}{\prod_{j=1}^n \prod_{k=1}^n (v_j - V_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))}. \end{aligned} \quad (9.14)$$

Recall that $u = x^0 + x^1$, $v = x^0 - x^1$, $U = y^0 + y^1$ and $V = y^0 - y^1$ (6.7).

Thus the $2n$ -point Green functions (9.1) are equal to

$$\begin{aligned} G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_1(x_1) \dots \psi_1(x_n) \psi_1^\dagger(y_1) \dots \psi_1^\dagger(y_n)) | 0 \rangle = \\ &\times (-1)^n d_\psi(g, \alpha_J) \left(\frac{\Lambda}{2\pi i} \right)^n \prod_{j < k}^n [\Lambda(v_j - v_k - i0 \cdot \varepsilon(x_j^0 - x_k^0))]^{d_\psi(g, \alpha_J)} \\ &\times [\Lambda(u_j - u_k - i0 \cdot \varepsilon(x_j^0 - x_k^0))]^{1 + d_\psi(g, \alpha_J)} \\ &\times \prod_{j < k}^n [\Lambda(V_j - V_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))]^{d_\psi(g, \alpha_J)} [\Lambda(U_j - U_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))]^{1 + d_\psi(g, \alpha_J)} \\ &\times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{[\Lambda(v_j - V_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{d_\psi(g, \alpha_J)} \\ &\times \left(\frac{1}{[\Lambda(u_j - U_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{1 + d_\psi(g, \alpha_J)}, \\ G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\psi_2(x_1) \dots \psi_2(x_n) \psi_2^\dagger(y_1) \dots \psi_2^\dagger(y_n)) | 0 \rangle = \\ &\times (-1)^n d_\psi(g, \alpha_J) \left(\frac{\Lambda}{2\pi i} \right)^n \prod_{j < k}^n [\Lambda(v_j - v_k - i0 \cdot \varepsilon(x_j^0 - x_k^0))]^{1 + d_\psi(g, \alpha_J)} \end{aligned}$$

$$\begin{aligned}
& \times [\Lambda(u_j - u_k - i0 \cdot \varepsilon(x_j^0 - x_k^0))]^{d_\psi(g, \alpha_J)} \\
& \times \prod_{j < k}^n [\Lambda(V_j - V_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))]^{1 + d_\psi(g, \alpha_J)} [\Lambda(U_j - U_k - i0 \cdot \varepsilon(y_j^0 - y_k^0))]^{d_\psi(g, \alpha_J)} \\
& \times \prod_{j=1}^n \prod_{k=1}^n \left(\frac{1}{[\Lambda(v_j - V_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{1 + d_\psi(g, \alpha_J)} \\
& \times \left(\frac{1}{[\Lambda(u_j - U_k - i0 \cdot \varepsilon(x_j^0 - y_k^0))]} \right)^{d_\psi(g, \alpha_J)}, \tag{9.15}
\end{aligned}$$

where $d_\psi(g, \alpha_J)$ is the dynamical dimension of the massless Thirring fermionic field defined by (8.8).

We can reduce the number of bosonic degrees of freedom defining $2n$ -point Green function. Making $O(1, 1)$ rotations in the (η, ξ) functional space

$$\begin{aligned}
\Theta(x) &= \frac{\sqrt{g} \eta(x) + \lambda(\alpha_J) \xi(x)}{\sqrt{g - \lambda^2(\alpha_J)}}, \quad \Phi(x) = \frac{\lambda(\alpha_J) \eta(x) + \sqrt{g} \xi(x)}{\sqrt{g - \lambda^2(\alpha_J)}}, \\
\Theta(x) &= \frac{\sqrt{g} \eta(x) - \lambda(\alpha_J) \xi(x)}{\sqrt{g - \lambda^2(\alpha_J)}}, \quad \Phi(x) = \frac{\lambda(\alpha_J) \eta(x) - \sqrt{g} \xi(x)}{\sqrt{g - \lambda^2(\alpha_J)}} \tag{9.16}
\end{aligned}$$

for $G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n)$ and $G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n)$, respectively, we arrive at the expressions

$$\begin{aligned}
G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle \\
&\times \int \mathcal{D}\Theta \mathcal{D}\Phi e^{i \int d^2x \left[\frac{1}{2} \partial_\mu \Theta(x) \partial^\mu \Theta(x) - \frac{1}{2} \partial_\mu \Phi(x) \partial^\mu \Phi(x) \right]} \\
&\times \prod_{j=1}^n e^{i \sqrt{4\pi d_\psi(g, \alpha_J)} \Theta(x_j)} \prod_{j=1}^n e^{-i \sqrt{4\pi d_\psi(g, \alpha_J)} \Theta(y_j)}, \\
G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle \\
&\times \int \mathcal{D}\Theta \mathcal{D}\Phi e^{i \int d^2x \left[\frac{1}{2} \partial_\mu \Theta(x) \partial^\mu \Theta(x) - \frac{1}{2} \partial_\mu \Phi(x) \partial^\mu \Phi(x) \right]} \\
&\times \prod_{j=1}^n e^{i \sqrt{4\pi d_\psi(g, \alpha_J)} \Theta(x_j)} \prod_{j=1}^n e^{-i \sqrt{4\pi d_\psi(g, \alpha_J)} \Theta(y_j)}. \tag{9.17}
\end{aligned}$$

The Φ -field is decoupled from the system and can be integrated out. This yields

$$\begin{aligned}
G^{(1)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_1(x_1) \dots \Psi_1(x_n) \Psi_1^\dagger(y_1) \dots \Psi_1^\dagger(y_n)) | 0 \rangle \\
&\times \int \mathcal{D}\Theta e^{i \int d^2x \frac{1}{2} \partial_\mu \Theta(x) \partial^\mu \Theta(x)} \prod_{j=1}^n e^{i \sqrt{4\pi d_\psi(g, \alpha_J)} [\Theta(x_j) - \Theta(y_j)]}, \\
G^{(2)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \langle 0 | T(\Psi_2(x_1) \dots \Psi_2(x_n) \Psi_2^\dagger(y_1) \dots \Psi_2^\dagger(y_n)) | 0 \rangle \\
&\times \int \mathcal{D}\Theta e^{i \int d^2x \frac{1}{2} \partial_\mu \Theta(x) \partial^\mu \Theta(x)} \prod_{j=1}^n e^{i \sqrt{4\pi d_\psi(g, \alpha_J)} [\Theta(x_j) - \Theta(y_j)]}. \tag{9.18}
\end{aligned}$$

The integration over the Θ -field returns us to the expressions (9.6).

Thus, the $2n$ -point Green function evaluated within the path-integral approach agree with Klaiber's expressions up to an infrared renormalization removing infrared singularities of Klaiber's expression in the limit $\mu \rightarrow 0$. However, as we will show in section 11, this agreement is only superficial. The correlation functions evaluated in Klaiber's approach do not admit non-perturbative ultra-violet renormalization, whereas the correlation functions obtained within the path-integral approach do.

10 Correlation functions of left(right)handed fermion densities in Klaiber's operator formalism

In this section we analyse correlation functions of left(right)handed fermion densities within Klaiber's operator approach. In order to understand the peculiarities of correlation functions of left(right)handed fermion densities of Klaiber's approach it is sufficient to deal with the two-point correlation function

$$\left\langle 0 \left| T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y) \right) \right| 0 \right\rangle. \quad (10.1)$$

Then, it is convenient to treat the correlation function (10.1) as a limit of four-point correlation function

$$\left\langle 0 \left| T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y) \right) \right| 0 \right\rangle \quad (10.2)$$

at $x_1 \rightarrow x$ and $y_1 \rightarrow y$ [17].

Following Klaiber's prescription and using equations ((IV.12) and (IV.13) of Ref.[4]) for the four-point correlation function in the r.h.s. of (10.2) we obtain

$$\begin{aligned} & \left\langle 0 \left| T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_1) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_1) \right) \right| 0 \right\rangle = \\ & = \left\langle 0 \left| T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) \right| 0 \right\rangle \\ & \times \exp \{ -(a - b) i \Delta(x - x_1) - (a - b) i \Delta(y - y_1) + (a - b) i \Delta(x - y) \\ & + (a - b) i \Delta(x_1 - y_1) - (a + b) i \Delta(x - y_1) - (a + b) i \Delta(x_1 - y) \}. \end{aligned} \quad (10.3)$$

The parameters a and b are defined by (9.10) or (9.13) in terms of Klaiber's parameters α and β or σ , respectively.

Using (6.19) for the definition of the Green function $i \Delta(x - y)$ we get

$$\begin{aligned} & \left\langle 0 \left| T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_1) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_1) \right) \right| 0 \right\rangle = \\ & = \left\langle 0 \left| T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) \right| 0 \right\rangle \\ & \times \frac{[-\mu^2(x - y)^2 + i 0]^{\frac{a - b}{4\pi}} [-\mu^2(x_1 - y_1)^2 + i 0]^{\frac{a - b}{4\pi}}}{[-\mu^2(x - x_1)^2 + i 0]^{\frac{a - b}{4\pi}} [-\mu^2(y - y_1)^2 + i 0]^{\frac{a - b}{4\pi}}} \\ & \times \frac{1}{[-\mu^2(x - y_1)^2 + i 0]^{\frac{a + b}{4\pi}} [-\mu^2(x_1 - y)^2 + i 0]^{\frac{a + b}{4\pi}}}, \end{aligned} \quad (10.4)$$

where μ is infrared cut-off. Finally it should be taken in the limit $\mu \rightarrow 0$. The correlation function (10.4) agrees completely with Klaiber's expression (VII.5).

Setting then $x_1 = x$ and $y_1 = y$ in (10.3) we obtain the two-point correlation function

$$\begin{aligned} & \langle 0 | T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y) \right) | 0 \rangle = \\ & = \langle 0 | T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y) \right) | 0 \rangle \\ & \times \exp \{ -2(a - b) i \Delta(0) - 4b i \Delta(x - y) \} = \\ & = \left(\frac{\Lambda}{\mu} \right)^4 \frac{a + b}{4\pi} \frac{\Lambda^2}{4\pi^2} [-\Lambda^2(x - y)^2 + i0]^{-1 - b/\pi}. \end{aligned} \quad (10.5)$$

From (10.5) we conclude that

(i) the dynamical dimension of the left(right)handed fermion densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ is equal to

$$d_{\bar{\psi}\psi}(g, a, b) = \frac{b}{\pi}, \quad (10.6)$$

(ii) in the limit $\mu \rightarrow 0$ due to the factor $(\Lambda/\mu)^{(a+b)/\pi}$ Klaiber's expressions for correlation functions of $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ are either singular or vanish in accordance with the sign of $(a + b)$,

(iii) for the n -point correlation function of the left(right)handed fermion densities the transition to the ultra-violet cut-off would be accompanied by the appearance of the factor

$$\left(\frac{\Lambda}{\mu} \right)^{4n \frac{a+b}{4\pi}} = \left(\frac{\Lambda}{\mu} \right)^{4n d_{\bar{\psi}\psi}(g, a, b)}, \quad (10.7)$$

where $d_{\bar{\psi}\psi}(g, a, b)$ is defined by (9.13).

Now we suggest to evaluate the four-point correlation function (10.3) within the path-integral approach. For this aim it is convenient to introduce the generating functional

$$\begin{aligned} & Z_{\text{Th}}[J_-, J_+, \bar{J}_-, \bar{J}_+] = \\ & = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp i \int d^2x \left\{ \bar{\psi}(x) i \gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \right. \\ & + \bar{\psi}(x) \left(\frac{1 + \gamma^5}{2} \right) J_-(x) + \bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) J_+(x) \\ & \left. + \bar{J}_-(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x) + \bar{J}_+(x) \left(\frac{1 + \gamma^5}{2} \right) \psi(x) \right\}, \end{aligned} \quad (10.8)$$

where $J_\pm(x)$ and $\bar{J}_\pm(x)$ are external sources of the corresponding fermion densities.

After the set of transformations (1.12) – (1.22) the generating functional (10.8) acquires the form

$$Z_{\text{Th}}[J_-, J_+, \bar{J}_-, \bar{J}_+] = \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}\eta \mathcal{D}\xi$$

$$\begin{aligned}
& \times \exp i \int d^2x \left\{ \bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\
& + \bar{\Psi}(x) \left(\frac{1 + \gamma^5}{2} \right) J_-(x) e^{-i\sqrt{g} \eta(x) + i\lambda(\alpha_J) \xi(x)} \\
& + \bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) J_+(x) e^{-i\sqrt{g} \eta(x) - i\lambda(\alpha_J) \xi(x)} \\
& + \bar{J}_-(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x) e^{i\sqrt{g} \eta(x) - i\lambda(\alpha_J) \xi(x)} \\
& \left. + \bar{J}_+(x) \left(\frac{1 + \gamma^5}{2} \right) \Psi(x) e^{i\sqrt{g} \eta(x) + i\lambda(\alpha_J) \xi(x)} \right\}. \tag{10.9}
\end{aligned}$$

In terms of $Z_{\text{Th}}[J_-, J_+, \bar{J}_-, \bar{J}_+]$ the correlation function (10.3) is defined by

$$\begin{aligned}
& \langle 0 | T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_1) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_1) \right) | 0 \rangle = \\
& = \frac{\delta}{\delta J_+(x)} \frac{\delta}{\delta \bar{J}_-(x_1)} \frac{\delta}{\delta J_-(y)} \frac{\delta}{\delta \bar{J}_+(y_1)} Z_{\text{Th}}[J_-, J_+, \bar{J}_-, \bar{J}_+] \Big|_{J_- = J_+ = \bar{J}_- = \bar{J}_+ = 0} = \\
& = \langle 0 | T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) | 0 \rangle \\
& \times \int \mathcal{D}\eta \mathcal{D}\xi \exp i \int d^2x \left\{ \bar{\Psi}(x) i \gamma^\mu \partial_\mu \Psi(x) + \frac{1}{2} \partial_\mu \eta(x) \partial^\mu \eta(x) - \frac{1}{2} \partial_\mu \xi(x) \partial^\mu \xi(x) \right. \\
& \left. \times e^{-i\sqrt{g} [\eta(x) - \eta(x_1) + \eta(y) - \eta(y_1)]} e^{-i\lambda(\alpha_J) [\xi(x) + \xi(x_1) - \xi(y) - \xi(y_1)]} \right\}. \tag{10.10}
\end{aligned}$$

Integrating over η and ξ fields we obtain

$$\begin{aligned}
& \langle 0 | T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_1) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_1) \right) | 0 \rangle = \\
& = \langle 0 | T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) | 0 \rangle \\
& \times \exp \{ g [2i\Delta(0) - i\Delta(x - x_1) + i\Delta(x - y) - i\Delta(x - y_1) \\
& - i\Delta(x_1 - y) + i\Delta(x_1 - y_1) - i\Delta(y - y_1)] \} \\
& \times \exp \{ -\lambda^2(\alpha_J) [2i\Delta(0) + i\Delta(x - x_1) - i\Delta(x - y) - i\Delta(x - y_1) \\
& - i\Delta(x_1 - y) - i\Delta(x_1 - y_1) + i\Delta(y - y_1)] \} = \\
& = \langle 0 | T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) | 0 \rangle \\
& \times \exp \{ 2(g - \lambda^2(\alpha_J)) i\Delta(0) - (g + \lambda^2(\alpha_J)) i\Delta(x - x_1) + (g + \lambda^2(\alpha_J)) i\Delta(x - y) \\
& - (g - \lambda^2(\alpha_J)) i\Delta(x - y_1) - (g - \lambda^2(\alpha_J)) i\Delta(x_1 - y) + (g + \lambda^2(\alpha_J)) i\Delta(x_1 - y_1) \\
& - (g + \lambda^2(\alpha_J)) i\Delta(y - y_1) \}. \tag{10.11}
\end{aligned}$$

Using (6.19) and (6.20) we get

$$\begin{aligned}
& \langle 0 | T \left(\bar{\psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \psi(x_1) \bar{\psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \psi(y_1) \right) | 0 \rangle = \\
& = \langle 0 | T \left(\bar{\Psi}(x) \left(\frac{1 - \gamma^5}{2} \right) \Psi(x_1) \bar{\Psi}(y) \left(\frac{1 + \gamma^5}{2} \right) \Psi(y_1) \right) | 0 \rangle
\end{aligned}$$

$$\begin{aligned}
& \times \frac{[-\Lambda^2(x-y)^2 + i0] \frac{g + \lambda^2(\alpha_J)}{4\pi} [-\Lambda^2(x_1 - y_1)^2 + i0] \frac{g + \lambda^2(\alpha_J)}{4\pi}}{[-\Lambda^2(x - x_1)^2 + i0] \frac{g + \lambda^2(\alpha_J)}{4\pi} [-\Lambda^2(y - y_1)^2 + i0] \frac{g + \lambda^2(\alpha_J)}{4\pi}} \\
& \times \frac{1}{[-\Lambda^2(x - y_1)^2 + i0] \frac{g - \lambda^2(\alpha_J)}{4\pi} [-\Lambda^2(x_1 - y)^2 + i0] \frac{g - \lambda^2(\alpha_J)}{4\pi}}. \quad (10.12)
\end{aligned}$$

Matching (10.12) with (10.4) up to a redefinition of powers

$$\frac{g + \lambda^2(\alpha_J)}{4\pi} \rightarrow \frac{a - b}{4\pi}, \quad \frac{g - \lambda^2(\alpha_J)}{4\pi} \rightarrow \frac{a + b}{4\pi} \quad (10.13)$$

the expression (10.12) agrees with (10.4). The most important distinction is the dependence of Klaiber's expression (10.4) on the infrared cut-off μ , whereas the correlation function (10.12) evaluated within the path-integral approach is infrared stable and depends only on the ultra-violet cut-off. This confirms our statement that Klaiber's correlation functions demand infrared regularization. After infrared regularization they should be compatible to the correlations functions evaluated within the path-integral approach. The problem of infrared regularization of Klaiber's correlation functions we solve in the next section. But it is important to emphasize that Klaiber's correlation functions are not subject to non-perturbative ultra-violet renormalization.

11 Non-perturbative renormalization of the massless Thirring model in the chiral symmetric phase

As has been shown above a comparison of the correlation functions evaluated within Klaiber's operator formalism with the results obtained in the path-integral technique testifies the necessity of infrared renormalization of Klaiber's expressions making them infrared convergent. This infrared renormalization is related to full extent to the non-perturbative renormalization of the massless Thirring model in the chiral symmetric phase. According to the standard procedure of renormalization in quantum field theory [29] we would understand the renormalizability of the massless Thirring model as a possibility to remove all infrared and ultra-violet divergences by renormalization of the wave function of the massless Thirring fermion field $\psi(x)$ and the coupling constant g .

Let us rewrite the Lagrangian (1.1) in terms of *bare* quantities

$$\mathcal{L}_{\text{Th}}(x) = \bar{\psi}_0(x) i\gamma^\mu \partial_\mu \psi_0(x) - \frac{1}{2} g_0 \bar{\psi}_0(x) \gamma^\mu \psi_0(x) \bar{\psi}_0(x) \gamma_\mu \psi_0(x), \quad (11.1)$$

where $\psi_0(x)$, $\bar{\psi}_0(x)$ are *bare* fermionic field operators and g_0 is a *bare* coupling constant.

The renormalized Lagrangian $\mathcal{L}(x)$ of the massless Thirring model should then read [29]

$$\begin{aligned}
\mathcal{L}_{\text{Th}}(x) &= \bar{\psi}(x) i\gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) \\
&+ (Z_2 - 1) \bar{\psi}(x) i\gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g (Z_1 - 1) \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x) = \\
&= Z_2 \bar{\psi}(x) i\gamma^\mu \partial_\mu \psi(x) - \frac{1}{2} g Z_1 \bar{\psi}(x) \gamma^\mu \psi(x) \bar{\psi}(x) \gamma_\mu \psi(x), \quad (11.2)
\end{aligned}$$

where Z_1 and Z_2 are the renormalization constants of the coupling constant and the wave function of the fermion field.

The renormalized fermionic field operator $\psi(x)$ and the coupling constant g are related to *bare* quantities by the relations [29]

$$\begin{aligned}\psi_0(x) &= Z_2^{1/2} \psi(x), \\ g_0 &= Z_1 Z_2^{-2} g.\end{aligned}\tag{11.3}$$

For the correlation functions of massless Thirring fermions the renormalizability of the massless Thirring model means the possibility to replace the infrared cut-off μ or the ultra-violet cut-off Λ by another finite scale M by means of the renormalization constants Z_1 and Z_2 .

According to the general theory of renormalizations [29] the renormalization constants Z_1 and Z_2 depend on the renormalized quantities g , the infrared scale μ , the ultra-violet scale Λ and the finite scale M :

$$\begin{aligned}Z_1 &= Z_1(g, M; \mu, \Lambda), \\ Z_2 &= Z_2(g, M; \mu, \Lambda).\end{aligned}\tag{11.4}$$

Since the correlation functions evaluated within the path-integral approach do not depend on the infrared cut-off μ , let us, first, consider the renormalization of them. In this a little simpler case the renormalization constants should be written as

$$\begin{aligned}Z_1 &= Z_1(g, M; \Lambda), \\ Z_2 &= Z_2(g, M; \Lambda).\end{aligned}\tag{11.5}$$

For the analysis of the feasibility of the replacement $\Lambda \rightarrow M$ it is convenient to introduce the following notations

$$\begin{aligned}G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \Lambda^n G_0^{(i)}(d_\psi(g_0, \alpha_J); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n), \\ G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \Lambda^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, \alpha_J); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n),\end{aligned}\tag{11.6}$$

where $G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n)$ and $G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)$ are the $2n$ -point Green function (8.6) and the n -point correlation function (6.22).

The transition to a finite scale M changes the Green functions (11.6) as follows

$$\begin{aligned}G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \\ &= \left(\frac{\Lambda}{M}\right)^{-2nd_\psi(g_0, \alpha_J)} M^n G_0^{(i)}(d_\psi(g_0, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n), \\ G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \\ &= \left(\frac{\Lambda}{M}\right)^{-2nd_{\bar{\psi}\psi}(g_0, \alpha_J)} M^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n).\end{aligned}\tag{11.7}$$

The renormalized correlation functions are related to the *bare* ones by the relations [29]:

$$G^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) = Z_2^{-n} G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) =$$

$$\begin{aligned}
&= Z_2^{-n} \left(\frac{\Lambda}{M} \right)^{-2nd_\psi(g_0, \alpha_J)} M^n G_0^{(i)}(d_\psi(g_0, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n), \\
G^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n) &= Z_2^{-2n} G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n) = \\
&= Z_2^{-2n} \left(\frac{\Lambda}{M} \right)^{-2nd_{\bar{\psi}\psi}(g_0, \alpha_J)} M^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n).
\end{aligned} \tag{11.8}$$

Renormalizability demands the relations

$$\begin{aligned}
G^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) &= M^n G^{(i)}(d_\psi(g, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n), \\
G^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n) &= M^{2n} G^{(\pm)}(d_{\bar{\psi}\psi}(g, \alpha_J); Mx_1, \dots, Mx_n; My_1, \dots, My_n),
\end{aligned} \tag{11.9}$$

which impose constraints on the dynamical dimensions and renormalization constants

$$\begin{aligned}
d_\psi(g, \alpha_J) &= d_\psi(Z_1 Z_2^{-2} g, \alpha_J), \\
d_{\bar{\psi}\psi}(g, \alpha_J) &= d_{\bar{\psi}\psi}(Z_1 Z_2^{-2} g, \alpha_J), \\
Z_2^{-1} \left(\frac{\Lambda}{M} \right)^{-2d_\psi(g, \alpha_J)} &= Z_2^{-1} \left(\frac{\Lambda}{M} \right)^{-d_{\bar{\psi}\psi}(g, \alpha_J)} = 1.
\end{aligned} \tag{11.10}$$

Using the exact expressions for the dynamical dimensions (6.22) and (8.8) one can show that the first two constraints on the dynamical dimensions are fulfilled only if the renormalization constants are related by

$$Z_1 = Z_2^2. \tag{11.11}$$

The important consequence of this relation is that the coupling constant g of the massless Thirring model is unrenormalized, i.e.

$$g_0 = g. \tag{11.12}$$

This also implies that the Gell–Mann–Low β -function, defined by [29]

$$M \frac{dg}{dM} = \beta(g, M), \tag{11.13}$$

should vanish, since g is equal to g_0 which does not depend on M , i.e. $\beta(g, M) = 0$ in the massless Thirring model. Our observation concerning the unrenormalizability of the coupling constant, $g_0 = g$, is supported by the analysis of renormalizability of the massive Thirring model carried out by Mueller and Trueman [19] and Gomes and Lowenstein [20].

The fulfillment of the third constraint in (11.10) demands a relation between the dynamical dimensions

$$d_{\bar{\psi}\psi}(g, \alpha_J) = 2 d_\psi(g, \alpha_J). \tag{11.14}$$

In order to satisfy this relation we can use the arbitrariness of the parameter α_J . Solving the equation (11.14) with respect to α_J we find

$$\alpha_J = -\frac{2\pi}{g}. \quad (11.15)$$

The dynamical dimensions are equal to

$$d_{\bar{\psi}\psi}(g, \alpha_J) = 2 d_{\psi}(g, \alpha_J) \Big|_{\alpha_J = -2\pi/g} = \frac{g}{\pi}. \quad (11.16)$$

The renormalization constants Z_1 and Z_2 read then

$$Z_1(g, M; \Lambda) = \left(\frac{M}{\Lambda}\right)^{2g/\pi}, \quad Z_2(g, M; \Lambda) = \left(\frac{M}{\Lambda}\right)^{g/\pi}. \quad (11.17)$$

Hence, for example, the *bare* two-point Green function and the renormalized one are related by

$$\begin{aligned} S_F^{(0)}(x-y) &= i\langle 0 | T(\psi_0(x) \bar{\psi}_0(y)) | 0 \rangle = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0} [-\Lambda^2(x-y)^2 + i0]^{-g_0/2\pi} \rightarrow \\ S_F(x-y) &= i\langle 0 | T(\psi(x) \bar{\psi}(y)) | 0 \rangle = \frac{1}{2\pi} \frac{\hat{x} - \hat{y}}{(x-y)^2 - i0} [-M^2(x-y)^2 + i0]^{-g/2\pi}. \end{aligned} \quad (11.18)$$

The Fourier transform of the two-point Green function reads (see Appendix E):

$$S_F(p) = -\frac{g e^{-ig/2}}{2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \frac{\hat{p}}{p^2 + i0} \left(\frac{p^2 + i0}{\bar{M}^2}\right)^{\frac{g}{2\pi}}, \quad (11.19)$$

where $\bar{M} = 2M$. The Fourier transform of the two-point Green function possesses singularities in the g -plane at $g = \pm 2\pi$ in agreement with Klaiber's analysis [27]. The Green function (11.19) agrees well with Glaser's analysis of the massless Thirring model [30,31] and the spectral analysis of two-point correlation functions developed by Schroer [32].

Now let us show that infrared singularities of Klaiber's correlation functions can be removed by renormalization of the wave function of the massless fermionic field in the case of the unrenormalized coupling constant $g_0 = g$.

In Klaiber's formalism the correlation functions are defined by

$$\begin{aligned} G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= \mu^n G_0^{(i)}(d_{\psi}(g_0, a, b); \mu x_1, \dots, \mu x_n; \mu y_1, \dots, \mu y_n)_K, \\ G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= \mu^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, a, b); \mu x_1, \dots, \mu x_n; \mu y_1, \dots, \mu y_n)_K, \end{aligned} \quad (11.20)$$

where μ is an infrared cut-off. The dynamical dimensions $d_{\psi}(g_0, a, b)$ and $d_{\bar{\psi}\psi}(g_0, a, b)$ are defined by (9.13) and (10.6), respectively.

In order to compare Klaiber's expressions with those evaluated within the path integral approach we should consider the transition of the infrared into an ultra-violet scale, $\mu \rightarrow \Lambda$. This changes the Green functions (11.20) as follows

$$\begin{aligned}
G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= \\
&= \left(\frac{\mu}{\Lambda}\right)^{-2nd_\psi(g_0, a, b)} \Lambda^n G_0^{(i)}(d_\psi(g_0, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K, \\
G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= \\
&= \left(\frac{\mu}{\Lambda}\right)^{-4nd_{\bar{\psi}\psi}(g_0, a, b)} \Lambda^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K.
\end{aligned} \tag{11.21}$$

Here we have taken into account Eq.(10.5) testifying the appearance of the factor (10.7) for the rescaling $\mu \rightarrow \Lambda$.

The infrared-renormalized correlation functions are related to the *bare* ones by the relations [29]:

$$\begin{aligned}
G^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= z_2^{-n} G_0^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n)_K = \\
&= z_2^{-n} \left(\frac{\mu}{\Lambda}\right)^{-2nd_\psi(g_0, a, b)} M^n G_0^{(i)}(d_\psi(g_0, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K, \\
G^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= z_2^{-2n} G_0^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)_K = \\
&= z_2^{-2n} \left(\frac{\mu}{\Lambda}\right)^{-4nd_{\bar{\psi}\psi}(g_0, a, b)} \Lambda^{2n} G_0^{(\pm)}(d_{\bar{\psi}\psi}(g_0, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K.
\end{aligned} \tag{11.22}$$

We denote the renormalization constants providing the infrared renormalization of correlation functions with small letters z_1 and z_2 , respectively. They depend on g , Λ and μ

$$\begin{aligned}
z_1 &= z_1(g, \Lambda; \mu), \\
z_2 &= z_2(g, \Lambda; \mu).
\end{aligned} \tag{11.23}$$

Infrared renormalizability, i.e. independence on the infrared cut-off μ , assumes the relations

$$\begin{aligned}
G^{(i)}(x_1, \dots, x_n; y_1, \dots, y_n) &= \Lambda^n G^{(i)}(d_\psi(g, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K, \\
G^{(\pm)}(x_1, \dots, x_n; y_1, \dots, y_n)_K &= \Lambda^{2n} G^{(\pm)}(d_{\bar{\psi}\psi}(g, a, b); \Lambda x_1, \dots, \Lambda x_n; \Lambda y_1, \dots, \Lambda y_n)_K,
\end{aligned} \tag{11.24}$$

which impose the relations

$$\begin{aligned}
d_\psi(g, a, b) &= d_\psi(z_1 z_2^{-2} g, a, b), \\
d_{\bar{\psi}\psi}(g, a, b) &= d_{\bar{\psi}\psi}(z_1 z_2^{-2} g, a, b), \\
z_2^{-1} \left(\frac{\mu}{\Lambda}\right)^{-2d_\psi(g, a, b)} &= 1.
\end{aligned} \tag{11.25}$$

This gives the following relations. $g_0 = g$, $z_1 = z_2^2$ and

$$z_2 = \left(\frac{\mu}{\Lambda}\right)^{2d_\psi(g, a, b)}. \quad (11.26)$$

Hence, we have shown that the infrared singularities of Klaiber's correlation functions can be removed by regularization of the wave functions of the massless fermionic field at the unrenormalized coupling constant $g_0 = g$.

The ultra-violet renormalizability of the massless Thirring model in Klaiber's formalism imposes the constraint

$$a = b \quad (11.27)$$

which leads to the dynamical dimension of the massless fermionic field equal to

$$d_\psi(g_0, a, b) = \frac{b}{2\pi}. \quad (11.28)$$

The dynamical dimension $d_{2n\psi}$ of any $2n$ -fermion correlation function is merely the product $d_{2n\psi} = 2nd_\psi(g_0, a, b) = nb/\pi$.

In order to show that the constraint (11.27) leads to a quantum field theory of a free massless fermionic field we suggest to use the parameterization of the parameters a and b in terms of ρ . This yield

$$a = \pi(\rho^2 - 1) \quad , \quad b = \pi\left(\frac{1}{\rho^2} - 1\right). \quad (11.29)$$

Thereby, the constraint (11.27) entails that $\rho = \pm 1$. This gives

$$a = b = 0 \quad (11.30)$$

and corresponds to a quantum field theory of a free massless fermionic field. In fact, in our parameterization (5.66) we have $\rho = 1$ that provides $\alpha = \beta = 0$ (see (5.25)). Setting $\alpha = \beta = 0$ in (4.18) and (5.1) we get $\psi(x) = \Psi(x)$.

It is interesting to notice that in Klaiber's parameterization there are two free massless fermionic fields $\psi(x)$ defined for $\alpha = \beta = 0$ at $\rho = 1$ and for $\alpha = \beta = -2\sqrt{\pi}$ at $\rho = -1$.

12 Fermion condensate and instability of chiral symmetric phase under spontaneous breaking of chiral symmetry

As has been shown in Ref.[6] in the massless Thirring model the chirally broken phase is energetically preferable with respect to the chiral symmetric one. In this section we show that the solution of the massless Thirring model in the chiral symmetric phase is unstable under spontaneous breaking of chiral symmetry. For this aim we treat a causal two-point Green function (11.19) and show the existence of a non-vanishing value of the fermion condensate.

The two-point Green function $S_F(x) = i\langle 0 | T(\psi(x)\bar{\psi}(0)) | 0 \rangle$ can be represented in the form [33]

$$S_F(x) = i\langle 0 | T(\psi(x)\bar{\psi}(0)) | 0 \rangle = \frac{g^2 e^{-ig/2}}{\bar{M}^2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \times \int \frac{d^2 p}{(2\pi)^2} e^{-ip \cdot x} \int \frac{d^2 k}{(2\pi)^2 i} \frac{\hat{k}}{k^2 + i0} \left(\frac{(p-k)^2 + i0}{\bar{M}^2} \right)^{-1 + g/2\pi}. \quad (12.1)$$

The Fourier transform is defined by

$$S_F(p) = \frac{g^2 e^{-ig/2}}{\bar{M}^2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \int \frac{d^2 k}{(2\pi)^2 i} \frac{\hat{k}}{k^2 + i0} \left(\frac{(p-k)^2 + i0}{\bar{M}^2} \right)^{-1 + g/2\pi}. \quad (12.2)$$

Now let us proceed to Euclidean momentum space [16]

$$S_F(p_E) = -e^{-ig/2} \frac{C(g)}{\bar{M}^2} \int \frac{d^2 k_E}{(2\pi)^2} \frac{\hat{k}_E}{k_E^2 - i0} \left(\frac{-(p_E - k_E)^2 + i0}{\bar{M}^2} \right)^{-1 + g/2\pi}, \quad (12.3)$$

where we have denoted

$$C(g) = \frac{g^2}{\sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)}. \quad (12.4)$$

Using the integral representation

$$\begin{aligned} & \left(\frac{-(p_E - k_E)^2 + i0}{\bar{M}^2} \right)^{-1 + g/2\pi} = \\ & = \frac{-i e^{ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \int_0^\infty dt t^{-g/2\pi} \exp\left\{ -i \left(\frac{(p_E - k_E)^2 - i0}{\bar{M}^2} \right) t \right\}, \end{aligned} \quad (12.5)$$

substituting (12.5) in (12.3) and integrating over directions of 2-Euclidean vector k_E we obtain

$$\begin{aligned} S_F(p_E) &= \frac{C(g)}{2\pi \bar{M}^2} \frac{i e^{-ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \frac{\hat{p}_E}{p_E} \int_0^\infty dt t^{-g/2\pi} \exp\left(-i \frac{p_E^2 t}{\bar{M}^2} \right) \\ &\times \int_0^\infty dk_E J_1\left(\frac{2t}{\bar{M}^2} p_E k_E \right) \exp\left(-i \frac{k_E^2 t}{\bar{M}^2} \right), \end{aligned} \quad (12.6)$$

where $J_1(z)$ is the Bessel function [34]. Integrating over k_E we obtain [35]

$$\int_0^\infty dk_E J_1\left(\frac{2t}{\bar{M}^2} p_E k_E \right) \exp\left(-i \frac{k_E^2 t}{\bar{M}^2} \right) = \frac{\bar{M}^2}{it p_E} \sin\left(\frac{p_E^2 t}{2\bar{M}^2} \right) \exp\left(+i \frac{p_E^2 t}{2\bar{M}^2} \right). \quad (12.7)$$

Substituting (12.7) in (12.6) we arrive at the expression

$$S_F(p_E) = -\frac{C(g)}{4\pi} \frac{e^{-ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \frac{i}{\hat{p}_E} \int_0^\infty \frac{dt}{t^{1+g/2\pi}} \left[1 - \exp\left(-i\frac{p_E^2 t}{\bar{M}^2}\right)\right]. \quad (12.8)$$

In order to regularize the integral over t we introduce the infrared cut-off μ

$$S_F(p_E) = -\frac{C(g)}{4\pi} \frac{e^{-ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \frac{i}{\hat{p}_E} \int_0^\infty \frac{dt}{t^{1+g/2\pi}} \left[\exp\left(-i\frac{\mu^2 t}{\bar{M}^2}\right) - \exp\left(-i\frac{p_E^2 t}{\bar{M}^2}\right)\right]. \quad (12.9)$$

The result of the integration over t reads

$$S_F(p_E) = \frac{g}{2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \frac{i}{\hat{p}_E} \left[\left(\frac{\mu^2}{\bar{M}^2}\right)^{g/2\pi} - \left(\frac{p_E^2}{\bar{M}^2}\right)^{g/2\pi}\right]. \quad (12.10)$$

Taking the limit $\mu \rightarrow 0$ we obtain

$$S_F(p_E) = -\frac{g}{2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \frac{i}{\hat{p}_E} \left(\frac{p_E^2}{\bar{M}^2}\right)^{g/2\pi}. \quad (12.11)$$

This agrees with (11.19). Then, the fermion condensate is defined by

$$\langle 0 | \bar{\psi}(0) \psi(0) | 0 \rangle = - \int \frac{d^2 p_E}{(2\pi)^2} \text{tr}\{S_F(p_E)\}. \quad (12.12)$$

Since the integrand is singular at $\hat{p}_E = 0$, we suggest to use the following integral representation

$$\frac{i}{\hat{p}_E} = \int_0^\infty d\tau e^{i\hat{p}_E \tau} = \int_0^\infty d\tau \left[\cos(p_E \tau) + i \frac{\hat{p}_E}{p_E} \sin(p_E \tau) \right] \quad (12.13)$$

valid for $p_E \neq 0$. In fact, the direct calculation of the integral over τ gives

$$\int_0^\infty d\tau \left[\cos(p_E \tau) + i \frac{\hat{p}_E}{p_E} \sin(p_E \tau) \right] = \pi \delta(p_E) + \frac{i}{\hat{p}_E}. \quad (12.14)$$

Since the l.h.s. of (12.13) is well-defined for $p_E \neq 0$, the δ -function vanishes and we obtain $i/\hat{p}_E$.

For the fermion condensate we obtain

$$\langle 0 | \bar{\psi}(0) \psi(0) | 0 \rangle = \frac{g}{\sin\left(\frac{g}{2}\right)} \frac{\bar{M}}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \int_0^\infty \frac{d\tau}{2\pi} \int_0^\infty dp_E \cos(p_E \tau) \left(\frac{p_E}{\bar{M}}\right)^{1+g/\pi}. \quad (12.15)$$

The regularization procedure implies the interchangeability of the integrations over τ and p_E . The r.h.s. of (12.15) does not vanish and gives a non-trivial value for the fermion

condensate. The appearance of a non-vanishing fermion condensate testifies instability of the chiral symmetric phase under spontaneous breaking of chiral symmetry.

The calculation of the integral over p_E should be carried out in the meaning of generalized functions [38]. This yields

$$\int_0^\infty dp_E \cos(p_E \tau) \left(\frac{p_E}{\bar{M}} \right)^{1+g/\pi} = -\cos\left(\frac{g}{2}\right) \Gamma\left(2 + \frac{g}{\pi}\right) \tau^{-2-g/\pi}. \quad (12.16)$$

Substituting (12.16) in (12.15), regularizing the integral over τ by the ultra-violet parameter Λ and renormalizing the fermion fields $\psi(x) \rightarrow Z_2^{1/2}(g, \Lambda, \bar{M}) \psi(x)$ we obtain

$$\langle 0 | \bar{\psi}(0) \psi(0) | 0 \rangle = -\frac{M(g)}{g}, \quad (12.17)$$

where $M(g)$, a dynamical mass of Thirring fermions in the chirally broken phase, we define following [6]

$$M(g) = \frac{\Lambda}{\sqrt{e^{2\pi/g} - 1}} \quad (12.18)$$

and $Z_2(g, \Lambda, \bar{M})$, the renormalization constant of the wave function of Thirring fermion fields, is equal to

$$Z_2(g, \Lambda, \bar{M}) = \left(e^{2\pi/g} - 1 \right)^{1/2} \frac{g^2}{2\pi} \frac{\text{ctg}\left(\frac{g}{2}\right)}{\Gamma\left(1 + \frac{g}{2\pi}\right)} \left(\frac{\Lambda}{\bar{M}} \right)^{g/\pi}. \quad (12.19)$$

Introducing the normalization scale $\bar{\mu}$ defined by

$$\bar{\mu} = \left(\frac{\Lambda^2}{\bar{M}} \right) \left[\left(e^{2\pi/g} - 1 \right)^{1/2} \frac{g^2}{2\pi} \frac{\text{ctg}\left(\frac{g}{2}\right)}{\Gamma\left(1 + \frac{g}{2\pi}\right)} \right]^{\pi/g}. \quad (12.20)$$

we transform the renormalization constant $Z_2(g, \Lambda, \bar{M})$ given by (12.19) to the form

$$Z_2(g, \Lambda, \bar{\mu}) = \left(\frac{\bar{\mu}}{\Lambda} \right)^{g/\pi} \quad (12.21)$$

in agreement with the definition (11.17).

The result (12.17) confirms qualitatively our conclusion concerning the instability of the chiral symmetric phase of the massless Thirring model under spontaneous breaking of chiral symmetry pointed out in Ref.[6].

Conclusion

We have analysed the solution of the massless Thirring model with fermion fields quantized in the chiral symmetric phase. In the literature the solution of the massless Thirring model in the form of n -point correlation functions has been carried out within the operator formalism developed by Klaiber [4] and by the path-integral technique initiated by Furuya, Gamboa Saravi and Schaposnik [7] and Naón [8]. We have discussed these two independent approaches. We have found that Klaiber's operator formalism suffers from the following problems: (i) unusual properties of composite fermionic field operators caused by the use of a solution of the Dirac equation contradicting to standard parity properties of fermionic fields, (ii) the definition of quantum vector and axial-vector fermion currents within the point-splitting technique leading to explicit breaking of chiral symmetry and (iii) the absence of a one-parameter family of solutions of quantum equations of motion of the massless Thirring model. In fact, a simultaneous evaluation of quantum vector and axial-vector fermion currents within Klaiber's approach supplemented by the requirement of the fulfillment of the standard relation $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$, caused by the properties of Dirac matrices in 1+1-dimensional space-time, fixes the parameter σ to $\sigma = 0$. Therefore, unlike Klaiber's claim none relation between Klaiber's [4] and Johnson's solutions of the massless Thirring model [36, 37] exists. Taking into account that at intermediate steps Klaiber's definition of quantum vector and axial-vector currents leads to an explicit breakdown of chiral symmetry (5.54), it is obvious that Klaiber's parameterization (5.43) is not applicable at all. On this way Klaiber's parameterization should be replaced by ours given by (5.66). In fact, this parameterization has been introduced within a correct definition of quantum vector and axial-vector current without violation of chiral symmetry and the standard relation $j_5^\mu(x) = -\varepsilon^{\mu\nu} j_\nu(x)$.

Unlike Klaiber's operator formalism the existence of a one-parameter family of solutions of the massless Thirring model is supported within the path-integral approach to the solution of the massless Thirring model. We have evaluated n -point correlation functions of left(right)handed fermion densities and $2n$ -point Green functions of fermion fields and found dynamical dimensions of these quantities depending on an arbitrary parameter α_J caused by ambiguities of the evaluation of chiral Jacobians quoted by Christos [11].

Such a free parameter in the solution of the massless Thirring model has turned out to be of use for the non-perturbative renormalization of the model. We have analysed the non-perturbative renormalization of the massless Thirring model in the standard meaning of renormalization as a possibility to replace the dependence of correlation functions and Green functions on the ultra-violet cut-off Λ by any finite scale M , $\Lambda \rightarrow M$, by means of the renormalization of the wave function of fermion fields and the coupling constant. We have shown that the coupling constant of the model g is unrenormalized that corresponds to the vanishing Gell-Mann-Low function, $\beta(g, M) = 0$. Our result on the unrenormalized coupling constant g is supported by similar conclusions which have been obtained by Mueller and Trueman [19] and Gomes and Lowenstein [20] who investigated the renormalizability of the massive Thirring model. The requirement of the complete replacement $\Lambda \rightarrow M$ in the expressions for n -point correlation functions of left(right)handed fermion field densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ and $2n$ -point Green function has allowed to fix the parameter α_J to the value $\alpha_J = -2\pi/g$. According to this result the dynamical dimensions of any n -fermion correlation function $d_{n\psi}$ are equal to $d_{n\psi} = nd_\psi$,

where $d_\psi = g/2\pi$.

We would like to emphasize that for $\alpha_J = -2\pi/g$ the kinetic term of the pseudoscalar field $\xi(x)$ acquires a correct sign. Due to this the symmetry group of the massless scalar and pseudoscalar fields $(\eta(x), \xi(x))$ becomes equal to the compact $O(2)$ group instead of the former non-compact $O(1, 1)$ group.

We have noted that n -point correlation functions evaluated with Klaiber's operator formalism contain infrared divergences. This is unlike the path-integral approach giving correlations functions and Green functions regular in the infrared limit. We have shown that by means of the renormalization of the wave functions of fermion fields the infrared cut-off can be replaced by an ultra-violet cut-off. This confirms the infrared stability of the massless Thirring model. Unfortunately, unlike the path-integral approach Klaiber's operator formalism has turned out to be inconsistent with a non-perturbative ultra-violet renormalization of the massless Thirring model. We have shown that Klaiber's definition of dynamical dimensions of n -point correlation functions of left(right)handed fermion densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ and $2n$ -point fermion Green functions are not invariant under the replacement of the ultra-violet cut-off Λ by any finite scale M . After a change $\Lambda \rightarrow M$ multiplicative factors depending on Λ are left which cannot be removed by a renormalization of wave functions of fermion fields.

One has to confess that Klaiber's operator formalism is not flexible enough for the non-perturbative ultra-violet renormalizability of the massless Thirring model. The dependence of the scalar and pseudoscalar fermion densities $S(x)$ and $P(x)$, related to vector and axial vector currents $J_\mu(x) = \partial_\mu S(x)/\sqrt{\pi}$ and $J_{5\mu}(x) = \partial_\mu P(x)/\sqrt{\pi}$, on the operators of creation $C^\dagger(k^1)$ and annihilation $C(k^1)$ leads to the appearance of Lorentz non-covariant contributions to correlation functions of fermion fields (9.9) which can be removed only by means of the constraint (5.26). Due to these constraint Klaiber's parameters α and β become expressed in terms of only one parameter ρ (5.42). This rules out the possibility for these parameters to be equal to each other. In fact, the requirement of an ultra-violet non-perturbative renormalizability of the massless Thirring model (11.27) leads to equality $\rho = \pm 1$.

Taking into account that the derivation of the quantum equation of motion of the massless Thirring model fixes all parameters of the canonical transformation suggested by Klaiber, (4.18) and (5.1), the possibility of the realization of a one-parameter family of solutions should be attributed to non-canonical contributions like chiral Jacobians in the path-integral approach. Hence, we argue that Klaiber's operator approach is not complete and should be supplemented by non-canonical contributions which are determined by chiral Jacobians, for example. We are planning to carry out this program in our forthcoming publications.

The transition from Klaiber's parameterization to ours changes Coleman's coupling constant relation (1.30) between the coupling constants of the Thirring model and the sine-Gordon model. Now according to our parameterization carried out within Klaiber's operator formalism it should read

$$\frac{4\pi}{\bar{\beta}^2} = \frac{1 + \frac{g}{2\pi}}{1 - \frac{g}{2\pi}}. \quad (\text{II.1})$$

In turn, within the path-integral definition of the dynamical dimension of the fermion field

densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$, $d_{\bar{\psi}\psi} = g/\pi$, Coleman's relation between coupling constants g and $\bar{\beta}$ should read

$$\frac{\bar{\beta}^2}{4\pi} = 1 + \frac{g}{\pi}. \quad (\text{II.2})$$

This relation is preferable with respect to (II.1), since it is obtained in the massless Thirring model subject to all requirements of non-perturbative renormalizability. The non-perturbative renormalizability of the sine-Gordon model we demonstrate in Appendix F by deriving a renormalized generating functional of Green functions.

We would like to emphasize that in the strong coupling limit (II.2) agrees with the relation obtained in Ref.[6] for the massive Thirring model quantized in the chirally broken phase. Indeed, as has been found in Ref.[6] in the chirally broken phase the relation between the coupling constants g and β reads

$$\frac{8\pi}{\bar{\beta}^2} = 1 - e^{-2\pi/g}. \quad (\text{II.3})$$

In the strong coupling limit $g \gg \pi$ the relations (II.2) and (II.3) have the same asymptotic behaviour

$$\frac{8\pi}{\bar{\beta}^2} = \frac{2\pi}{g} + O(g^{-2}). \quad (\text{II.4})$$

This implies the inequality $\bar{\beta}^2 \gg 8\pi$ that leads to the population of the 1+1-dimensional world with solitons having quantum numbers of Thirring fermions, see also Ref.[6]. It is important to underscore that according to Coleman's relation (1.30) the coupling constant $\bar{\beta}$ tends to zero, $\bar{\beta} \rightarrow 0$, at $g \rightarrow \infty$. This suppresses the production of soliton states with respect to the production of single quanta of the sine-Gordon field. The former is hardly comprehensible from a physical point of view, since in the strong coupling limit collective phenomena should play an important role. In Appendix F we have carried out a non-perturbative renormalization of the sine-Gordon model. We have shown that the sine-Gordon model is not asymptotically free. This is in complete agreement with the equivalence between the Thirring model and the sine-Gordon model. The unrenormalizability of the coupling constant $\bar{\beta}$ leading to the vanishing Gell-Mann-Low function can be easily understood following the similarity between $\bar{\beta}$ and $\hbar$ which has been drawn in Ref.[6]. As has been shown in Ref.[6] the limits $\bar{\beta} \rightarrow 0$ and $\bar{\beta} \rightarrow \infty$ distinguish *classical* and *quantum* regimes of the sine-Gordon model. Within such an understanding of the coupling constant $\bar{\beta}$ its unrenormalizability is justified by the unrenormalizability of $\hbar$.

Now we would like to make some remarks concerning Coleman's perturbation theory developed with respect to the mass m of the Thirring fermion field which was used for the derivation of equivalence between the massive Thirring model and the sine-Gordon model. An expansion of the generating functional of correlation functions in powers of m assumes the integration over fermionic degrees of freedom described by a free massless fermion field. However, as we have discussed in section 2 due to, say, the non-standard properties under parity transformations breaking the relation $v(p^0, p^1) = -\gamma^0 v(p^0, -p^1)$ (see (2.18)), Klaiber's solution for a free massless fermion field cannot be obtained as the massless limit of a free massive fermion field. In fact, a free massive fermion field with mass m possesses standard properties under parity transformations obeying the relation $v(p^0, p^1) = -\gamma^0 v(p^0, -p^1)$ (see Appendix A (A.15)). Therefore, the quantum field theory

of a free massless fermion field obtained in the limit $m \rightarrow 0$ differs from the quantum field theory of a free massless fermion field in Klaiber's approach. Thus, we think that Coleman's perturbation theory developed with respect to m does not exist from the point of view of constructive quantum field theory.

Finally, we would like to note that we have shown that the chiral symmetric phase of the massless Thirring model is not stable, and fermion condensation can occur even if the fermion system evolves from the chiral symmetric phase. Taking the Fourier transform of the two-point Green function of a massless Thirring fermion field evaluated in the chiral symmetric phase we have shown that fermion condensation can occur dynamically. The former confirms our statement in Ref.[6] concerning the energetic advantage of the chirally broken phase relative to the chiral symmetric one.

Acknowledgement

This work was supported in part by Fonds zur Förderung der Wissenschaftlichen Forschung P13997-TPH.

Appendix A. Parity properties of fermion fields

In this Appendix we recall the properties of fermion fields under parity transformation.

First, let us discuss fermion fields in 3+1-dimensional space-time. A free massive Dirac field of mass m obeys the equation

$$(i \gamma^\mu \partial_\mu - m) \psi(x) = 0, \quad (\text{A.1})$$

where $\gamma^\mu = (\gamma^0, \vec{\gamma})$ are Dirac matrices in the Dirac representation and $\psi(x) = \psi(t, \vec{x})$. The solution of (A.1) can be represented in the standard form of the expansion into plane waves

$$\begin{aligned} \psi(t, \vec{x}) = & \int \frac{d^3 p}{(2\pi)^{3/2}} \frac{1}{\sqrt{2p^0}} \\ & \times \sum_{\sigma=\pm 1/2} \left[u(p^0, \vec{p}, \sigma) a(\vec{p}, \sigma) e^{-ip^0 t + i\vec{p} \cdot \vec{x}} + v(p^0, \vec{p}, \sigma) b^\dagger(\vec{p}, \sigma) e^{ip^0 t - i\vec{p} \cdot \vec{x}} \right], \end{aligned} \quad (\text{A.2})$$

where $a(\vec{p}, \sigma)$ and $b(\vec{p}, \sigma)$ are annihilation operators of fermions and antifermions with momentum $\vec{p}$, energy $p^0 = \sqrt{\vec{p}^2 + m^2}$ and polarization σ . The Dirac bispinorial functions $u(p^0, \vec{p}, \sigma)$ and $v(p^0, \vec{p}, \sigma)$ have the standard form

$$u(p^0, \vec{p}, \sigma) = \sqrt{p^0 + m} \begin{pmatrix} \varphi_\sigma \\ \frac{\vec{\sigma} \cdot \vec{p}}{p^0 + m} \varphi_\sigma \end{pmatrix}, \quad v(p^0, \vec{p}, \sigma) = \sqrt{p^0 + m} \begin{pmatrix} \frac{\vec{\sigma} \cdot \vec{p}}{p^0 + m} \chi_\sigma \\ \chi_\sigma \end{pmatrix}, \quad (\text{A.3})$$

where $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are Pauli matrices, φ_σ and χ_σ are two-component spinors. The Dirac bispinorial functions (A.3) are normalized to

$$\begin{aligned} u^\dagger(p^0, \vec{p}, \sigma) u(p^0, \vec{p}, \sigma') &= v^\dagger(p^0, \vec{p}, \sigma) v(p^0, \vec{p}, \sigma') = 2p^0 \delta_{\sigma\sigma'}, \\ \bar{u}(p^0, \vec{p}, \sigma) u(p^0, \vec{p}, \sigma') &= -\bar{v}(p^0, \vec{p}, \sigma) v(p^0, \vec{p}, \sigma') = 2m \delta_{\sigma\sigma'}, \\ u^\dagger(p^0, \vec{p}, \sigma) v(p^0, \vec{p}, \sigma') &= v^\dagger(p^0, \vec{p}, \sigma) u(p^0, \vec{p}, \sigma') = 0 \end{aligned} \quad (\text{A.4})$$

and satisfy the matrix relations

$$\begin{aligned} \sum_{\sigma=\pm 1/2} u(p^0, \vec{p}, \sigma) \bar{u}(p^0, \vec{p}, \sigma) &= \begin{pmatrix} p^0 + m & -\vec{\sigma} \cdot \vec{p} \\ \vec{\sigma} \cdot \vec{p} & p^0 + m \end{pmatrix} = \hat{p} + m, \\ \sum_{\sigma=\pm 1/2} v(p^0, \vec{p}, \sigma) \bar{v}(p^0, \vec{p}, \sigma) &= \begin{pmatrix} p^0 - m & -\vec{\sigma} \cdot \vec{p} \\ \vec{\sigma} \cdot \vec{p} & -p^0 - m \end{pmatrix} = \hat{p} - m. \end{aligned} \quad (\text{A.5})$$

The Dirac bispinorial functions (A.3) are normalized by the conditions (A.4) and obey the identities

$$u(p^0, -\vec{p}, \sigma) = \gamma^0 u(p^0, \vec{p}, \sigma) \quad , \quad v(p^0, -\vec{p}, \sigma) = -\gamma^0 v(p^0, \vec{p}, \sigma). \quad (\text{A.6})$$

Under parity transformation the fermionic field (A.1) transforms in the standard way [2,3]

$$\mathcal{P} \psi(t, \vec{x}) \mathcal{P}^\dagger = \gamma^0 \psi(t, -\vec{x}), \quad (\text{A.7})$$

where we have dropped an insignificant phase factor. This relation follows directly from the scalar properties of the Lagrangian [3]

$$\mathcal{P} \mathcal{L}(t, \vec{x}) \mathcal{P}^\dagger = \mathcal{L}(t, -\vec{x}). \quad (\text{A.8})$$

Using the properties of the Dirac bispinorial functions (A.6) one can show that the operators $a(\vec{p}, \sigma)$, $a^\dagger(\vec{p}, \sigma)$, $b(\vec{p}, \sigma)$ and $b^\dagger(\vec{p}, \sigma)$ transform as follows

$$\begin{aligned} \mathcal{P} a(\vec{p}, \sigma) \mathcal{P}^\dagger &= a(-\vec{p}, \sigma) \quad , \quad \mathcal{P} b(\vec{p}, \sigma) \mathcal{P}^\dagger = -b(-\vec{p}, \sigma), \\ \mathcal{P} a^\dagger(\vec{p}, \sigma) \mathcal{P}^\dagger &= a^\dagger(-\vec{p}, \sigma) \quad , \quad \mathcal{P} b^\dagger(\vec{p}, \sigma) \mathcal{P}^\dagger = -b^\dagger(-\vec{p}, \sigma). \end{aligned} \quad (\text{A.9})$$

The properties of the fermionic field discussed above do not depend on the dimension of space-time and should be retained in 1+1-dimensional space-time too. We should only recall that in 1+1-dimensional space-time fermion fields have no spin and the dependence on σ , the projection of the spin onto the fermion momentum, vanishes.

In 1+1-dimensional space-time a solution of the Dirac equation for a free massive fermionic field expanded into plane waves reads [5,16]

$$\begin{aligned} \psi(x) &= \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} \left[u(p^0, p^1) a(p^1) e^{-ip \cdot x} + v(p^0, p^1) b^\dagger(p^1) e^{ip \cdot x} \right], \\ \bar{\psi}(x) &= \psi^\dagger(x) \gamma^0 = \int_{-\infty}^{\infty} \frac{dp^1}{\sqrt{2\pi}} \frac{1}{\sqrt{2p^0}} \left[\bar{u}(p^0, p^1) a^\dagger(p^1) e^{ip \cdot x} + \bar{v}(p^0, p^1) b(p^1) e^{-ip \cdot x} \right], \end{aligned} \quad (\text{A.10})$$

where $p \cdot x = p^0 x^0 - p^1 x^1$. The creation $a^\dagger(p^1)$ ($b^\dagger(p^1)$) and annihilation $a(p^1)$ ($b(p^1)$) operators of fermions (antifermions) with momentum p^1 and energy $p^0 = \sqrt{(p^1)^2 + m^2}$ obey the anticommutation relations

$$\begin{aligned} \{a(p^1), a^\dagger(q^1)\} &= \{b(p^1), b^\dagger(q^1)\} = \delta(p^1 - q^1), \\ \{a(p^1), a(q^1)\} &= \{a^\dagger(p^1), a^\dagger(q^1)\} = \{b(p^1), b(q^1)\} = \{b^\dagger(p^1), b^\dagger(q^1)\} = 0. \end{aligned} \quad (\text{A.11})$$

The wave functions $u(p^0, p^1)$ and $v(p^0, p^1)$ are the solutions of the Dirac equation in the momentum representation for positive and negative energies, respectively. They are defined by

$$\begin{aligned} u(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ \sqrt{p^0 - p^1} \end{pmatrix}, \quad \bar{u}(p^0, p^1) = (\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) \\ v(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ -\sqrt{p^0 - p^1} \end{pmatrix}, \quad \bar{v}(p^0, p^1) = (-\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) \end{aligned} \quad (\text{A.12})$$

at $p^0 = \sqrt{(p^1)^2 + m^2}$ and normalized to

$$\begin{aligned} u^\dagger(p^0, p^1) u(p^0, p^1) &= v^\dagger(p^0, p^1) v(p^0, p^1) = 2p^0, \\ \bar{u}(p^0, p^1) u(p^0, p^1) &= -\bar{v}(p^0, p^1) v(p^0, p^1) = 2m, \\ \bar{u}(p^0, p^1) v(p^0, p^1) &= \bar{v}(p^0, p^1) u(p^0, p^1) = 0. \end{aligned} \quad (\text{A.13})$$

The functions $u(p^0, p^1)$ and $v(p^0, p^1)$ satisfy the following matrix relations

$$\begin{aligned}
u(p^0, p^1) \bar{u}(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ \sqrt{p^0 - p^1} \end{pmatrix} (\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) = \\
&= \begin{pmatrix} \sqrt{(p^0)^2 - (p^1)^2} & p^0 + p^1 \\ p^0 - p^1 & \sqrt{(p^0)^2 - (p^1)^2} \end{pmatrix} = \begin{pmatrix} m & p^0 + p^1 \\ p^0 - p^1 & m \end{pmatrix} = \\
&= \gamma^0 p^0 - \gamma^1 p^1 + m = \hat{p} + m, \\
v(p^0, p^1) \bar{v}(p^0, p^1) &= \begin{pmatrix} \sqrt{p^0 + p^1} \\ -\sqrt{p^0 - p^1} \end{pmatrix} (-\sqrt{p^0 - p^1}, \sqrt{p^0 + p^1}) = \\
&= \begin{pmatrix} -\sqrt{(p^0)^2 - (p^1)^2} & p^0 + p^1 \\ p^0 - p^1 & -\sqrt{(p^0)^2 - (p^1)^2} \end{pmatrix} = \begin{pmatrix} -m & p^0 + p^1 \\ p^0 - p^1 & -m \end{pmatrix} = \\
&= \gamma^0 p^0 - \gamma^1 p^1 - m = \hat{p} - m
\end{aligned} \tag{A.14}$$

and the identities

$$u(p^0, -p^1) = \gamma^0 u(p^0, p^1) \quad , \quad v(p^0, -p^1) = -\gamma^0 v(p^0, p^1). \tag{A.15}$$

Under parity transformations a fermionic field in 1+1-dimensional space-time transforms in the usual way

$$\mathcal{P} \psi(t, x) \mathcal{P}^\dagger = \gamma^0 \psi(t, -x). \tag{A.15}$$

By virtue of the identities (A.15) the operators of creation and annihilation of fermions and antifermions behave under parity transformation in the standard way

$$\begin{aligned}
\mathcal{P} a(p^1) \mathcal{P}^\dagger &= a(-p^1) \quad , \quad \mathcal{P} b(p^1) \mathcal{P}^\dagger = -b(-p^1), \\
\mathcal{P} a^\dagger(p^1) \mathcal{P}^\dagger &= a^\dagger(-p^1) \quad , \quad \mathcal{P} b^\dagger(p^1) \mathcal{P}^\dagger = -b^\dagger(-p^1).
\end{aligned} \tag{A.16}$$

In the massless limit the wave functions $u(p^0, p^1)$ and $v(p^0, p^1)$ can be written in terms of Heaviside functions and read

$$\begin{aligned}
u(p^0, p^1) &= \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ \theta(-p^1) \end{pmatrix} \quad , \quad \bar{u}(p^0, p^1) = \sqrt{2p^0} (\theta(-p^1), \theta(+p^1)) \\
v(p^0, p^1) &= \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ -\theta(-p^1) \end{pmatrix} \quad , \quad \bar{v}(p^0, p^1) = \sqrt{2p^0} (-\theta(-p^1), \theta(+p^1)),
\end{aligned} \tag{A.17}$$

where $\theta(\pm p^1)$ are Heaviside functions.

In his seminal paper [1] Thirring and then Klaiber [4] used the solution for a free massless fermionic field in 1+1-dimensional space-time with identical spinorial functions $u(p^0, p^1)$ and $v(p^0, p^1)$

$$u(p^0, p^1) = v(p^0, p^1) = \sqrt{2p^0} \begin{pmatrix} \theta(+p^1) \\ \theta(-p^1) \end{pmatrix}. \tag{A.18}$$

Since the wave function $v(p^0, p^1)$ taken in the form (A.17) breaks the identity (A.14) that entails, in turn, a contradiction with the standard properties of fermion fields under parity transformations

$$\begin{aligned}
\mathcal{P} a(p^1) \mathcal{P}^\dagger &= a(-p^1) \quad , \quad \mathcal{P} b(p^1) \mathcal{P}^\dagger = b(-p^1), \\
\mathcal{P} a^\dagger(p^1) \mathcal{P}^\dagger &= a^\dagger(-p^1) \quad , \quad \mathcal{P} b^\dagger(p^1) \mathcal{P}^\dagger = b^\dagger(-p^1).
\end{aligned} \tag{A.19}$$

This violates the smooth transition from massive to massless fermionic fields in the zero-mass limit.

Appendix B. Chiral Jacobian

In this Appendix we adduce the calculation of the Jacobian induced by chiral rotations. We follow the procedure formulated in Refs.[7–14]. For the calculation of the chiral Jacobian we start with the Lagrangian defined by

$$\mathcal{L}_\psi(x) = \bar{\psi}(x) i\gamma^\mu (\partial_\mu - i\partial_\mu \eta(x) - i\varepsilon_{\mu\nu} \partial^\nu \xi(x)) \psi(x) = \bar{\psi}(x) D(x; 0) \psi(x), \quad (\text{B.1})$$

where $D(x; 0)$ is the Dirac operator given by

$$D(x; 0) = i\gamma^\mu (\partial_\mu - i\partial_\mu \eta(x) - i\varepsilon_{\mu\nu} \partial^\nu \xi(x)). \quad (\text{B.2})$$

By a chiral rotation

$$\begin{aligned} \psi(x) &= e^{i\eta(x) + i\alpha \gamma^5 \xi(x)} \chi(x), \\ \bar{\psi}(x) &= \bar{\chi}(x) e^{-i\eta(x) + i\alpha \gamma^5 \xi(x)}, \end{aligned} \quad (\text{B.3})$$

where $0 \leq \alpha \leq 1$, we reduce the Lagrangian (B.1) to the form

$$\mathcal{L}_\chi(x) = \bar{\chi}(x) D(x, \alpha) \chi(x). \quad (\text{B.4})$$

The Dirac operator $D(x; \alpha)$ reads

$$D(x; \alpha) = i\gamma^\mu \partial_\mu + (1 - \alpha) \gamma^\mu \gamma^5 \partial_\mu \xi(x). \quad (\text{B.5})$$

At $\alpha = 1$ we obtain the Lagrangian

$$\mathcal{L}_\chi(x) = \bar{\chi}(x) D(x, 1) \chi(x) = \bar{\chi}(x) i\gamma^\mu \partial_\mu \chi(x). \quad (\text{B.6})$$

Due to the chiral rotation (B.3) the fermionic measure changes as follows

$$\mathcal{D}\psi \mathcal{D}\bar{\psi} = J[\xi] \mathcal{D}\chi \mathcal{D}\bar{\chi}, \quad (\text{B.7})$$

For the calculation $J[\vartheta]$ we follow Fujikawa's procedure [7–14] and introduce eigenfunctions $\varphi_n(x; \alpha)$ and eigenvalues $\lambda_n(\alpha)$ of the Dirac operator $D(x; \alpha)$:

$$D(x; \alpha) \varphi_n(x; \alpha) = \lambda_n(\alpha) \varphi_n(x; \alpha). \quad (\text{B.8})$$

The operator $D^2(x; \alpha)$ reads then

$$D^2(x; \alpha) = -\square - 2i(1 - \alpha) \gamma^\mu \gamma^\nu \gamma^5 \partial_\mu \xi(x) \partial_\nu + i(1 - \alpha) \gamma^5 \square \xi - (1 - \alpha)^2 \partial_\mu \xi(x) \partial^\mu \xi(x). \quad (\text{B.9})$$

In terms of the eigenfunctions and eigenvalues of the Dirac operator $D(x; \alpha)$ the Jacobian $J[\xi]$ is defined by [9,10]

$$J[\xi] = \exp 2i \int_0^1 d\alpha w[\xi; \alpha], \quad (\text{B.10})$$

where the functional $w[\xi; \alpha]$ is given by [9,10]

$$w[\xi; \alpha] = - \lim_{\Lambda_F \rightarrow \infty} \sum_n \varphi_n^\dagger(x; \alpha) \gamma^5 \xi(x) e^{i\lambda^2(\alpha)_n / \Lambda_F^2} \varphi_n(x; \alpha) =$$

$$= - \lim_{\Lambda_F \rightarrow \infty} \int d^2x \xi(x) \int \frac{d^2k}{(2\pi)^2} \text{tr} \left\{ \gamma^5 \langle x | e^{iD^2(x; \alpha)/\Lambda_F^2} | x \rangle \right\}. \quad (\text{B.10})$$

For the calculation of the matrix element $\langle x | \dots | x \rangle$ we use plane waves [7–14] and get

$$\begin{aligned} & \langle x | e^{iD^2(x; \alpha)/\Lambda_F^2} | x \rangle = \\ & = \exp \left\{ \frac{i}{\Lambda_F^2} [k^2 + 2(1 - \alpha) \gamma^\mu \gamma^\nu \gamma^5 \partial_\mu \xi(x) k_\nu + i(1 - \alpha) \gamma^5 \square \xi - (1 - \alpha)^2 \partial_\mu \xi(x) \partial^\mu \xi(x)] \right\}. \end{aligned} \quad (\text{B.11})$$

Substituting (B.11) in (B.10) we obtain

$$\lim_{\Lambda_F \rightarrow \infty} \int \frac{d^2k}{(2\pi)^2} \text{tr} \left\{ \gamma^5 \langle x | e^{iD^2(x; \alpha)/\Lambda_F^2} | x \rangle \right\} = - \frac{1}{2\pi} (1 - \alpha) \square \xi(x). \quad (\text{B.12})$$

The functional $w[\xi, \alpha]$ is then given by

$$w[\xi, \alpha] = \frac{1}{2\pi} (1 - \alpha) \int d^2x \xi(x) \square \xi(x). \quad (\text{B.13})$$

Inserting $w[\xi, \alpha]$ into (B.9) and integrating over α we get the Jacobian

$$J[\xi] = \exp \left\{ - i \int d^2x \frac{1}{2\pi} \partial^\mu \xi(x) \partial_\mu \xi(x) \right\}. \quad (\text{B.14})$$

For the derivation of the exponent we have integrated by parts and dropped the surface contributions.

According to Refs.[10–12] the exponent in (B.14) is ill-defined and depends on the regularization procedure. Therefore, the chiral Jacobian can be written as

$$J[\xi] = \exp \left\{ - i \int d^2x \frac{\alpha_J}{2\pi} \partial^\mu \xi(x) \partial_\mu \xi(x) \right\}, \quad (\text{B.15})$$

where α_J is an arbitrary parameter.

In fact, following Christos [10] the functional $w[\xi; \alpha]$ can be represented in the form

$$\begin{aligned} & w[\xi; \alpha] = \\ & = - \lim_{\Lambda_F \rightarrow \infty} \sum_n \varphi_n^\dagger(x; \alpha) \gamma^5 \xi(x) \left[\alpha_J e^{i\lambda^2(\alpha_J)_n/\Lambda_F^2} + (1 - \alpha_J) e^{i(i\hat{\partial})^2/\Lambda_F^2} \right] \varphi_n(x; \alpha) = \\ & = - \lim_{\Lambda_F \rightarrow \infty} \int d^2x \xi(x) \int \frac{d^2k}{(2\pi)^2} \text{tr} \left\{ \gamma^5 \langle x | \alpha_J e^{iD^2(x; \alpha)/\Lambda_F^2} + (1 - \alpha_J) e^{i(i\hat{\partial})^2/\Lambda_F^2} | x \rangle \right\}. \end{aligned} \quad (\text{B.16})$$

Since the contribution of the last term vanishes, the functional $w[\xi; \alpha]$ becomes equal to

$$w[\xi, \alpha] = \frac{\alpha_J}{2\pi} (1 - \alpha) \int d^2x \xi(x) \square \xi(x). \quad (\text{B.17})$$

This leads to the chiral Jacobian given by (B.15).

In turn, in the massless Thirring model with fermionic fields quantized in the chirally broken phase the chiral Jacobian is well-defined and is equal to unity. In fact, as has been shown in [6] following Fujikawa's procedure chiral Jacobian can be obtained in the form

$$J[\vartheta] = \exp i \int d^2x \left[\frac{1}{8\pi} \partial^\mu \vartheta(x) \partial_\mu \vartheta(x) + \frac{M^2}{4\pi} (\cos 2\vartheta(x) - 1) \right], \quad (\text{B.18})$$

where $\vartheta(x)$ is a pseudoscalar field. According to Christos' remark the exponent in the r.h.s. of (11.19) should enter with an arbitrary parameter α_J , i.e.

$$J[\vartheta] = \exp i \frac{\alpha_J}{4\pi} \int d^2x \left[\frac{1}{2} \partial^\mu \vartheta(x) \partial_\mu \vartheta(x) + M^2 (\cos 2\vartheta(x) - 1) \right]. \quad (\text{B.19})$$

The term proportional to M^2 gives a contribution to the effective potential breaking the rotational symmetry of the effective potential under global rotations $\vartheta(x) \rightarrow \vartheta(x) + \theta$, where θ is an arbitrary constant. The effective potential has been evaluated explicitly [6] without chiral rotations and found invariant under global rotations $\vartheta(x) \rightarrow \vartheta(x) + \theta$. Therefore, the requirement of rotational symmetry of all contributions to the effective potential gives $\alpha_J = 0$. This yields $J[\vartheta] = 1$ [6].

Appendix C. Two-point correlation functions of scalar and pseudoscalar fermionic field densities $S^{(\pm)}(x)$ and $P^{(\pm)}(y)$

In this Appendix we calculate the correlation functions $D^{(\pm)}(x)$ and $D_5^{(\pm)}(x)$ defining the commutators of scalar and pseudoscalar fermionic field densities $S^{(\pm)}(x)$ and $P^{(\pm)}(y)$ given by (5.5) and (5.6).

Since the momentum integrals defining $D^{(\pm)}(x)$ and $D_5^{(\pm)}(x)$ are divergent in the infrared region, we have to regularize them. The simplest regularization is related to the inclusion of a non-zero mass μ which finally should be taken in the limit $\mu \rightarrow 0$. The correlation functions $D^{(\pm)}(x)$ and $D_5^{(\pm)}(x)$ read

$$\begin{aligned} D^{(\pm)}(x) &\rightarrow D^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{(k^1)^2 + \mu^2}} e^{\mp i \sqrt{(k^1)^2 + \mu^2} x^0 \pm i k^1 x^1}, \\ D_5^{(\pm)}(x) &\rightarrow D_5^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} \frac{dk^1}{\sqrt{(k^1)^2 + \mu^2}} \varepsilon(k^1) e^{\mp i \sqrt{(k^1)^2 + \mu^2} x^0 \pm i k^1 x^1}. \end{aligned} \quad (\text{C.1})$$

By a change of variables $k^1 = \mu \sinh \varphi$ we recast the integral over k^1 into the form

$$\begin{aligned} D^{(\pm)}(x) &\rightarrow D^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi e^{-i \mu (\pm x^0) \cosh \varphi \pm i \mu x^1 \sinh \varphi}, \\ D_5^{(\pm)}(x) &\rightarrow D_5^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi \varepsilon(\varphi) e^{-i \mu (\pm x^0) \cosh \varphi \pm i \mu x^1 \sinh \varphi}. \end{aligned} \quad (\text{C.2})$$

For the convergence of the integrals over φ we should introduce an infinitesimal imaginary part for x^0 : $\pm x^0 \rightarrow \pm x^0 - i0$. This yields

$$\begin{aligned} D^{(\pm)}(x) &\rightarrow D^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi e^{-i\mu(\pm x^0 - i0) \cosh \varphi \pm i\mu x^1 \sinh \varphi}, \\ D_5^{(\pm)}(x) &\rightarrow D_5^{(\pm)}(x; \mu) = \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi \varepsilon(\varphi) e^{-i\mu(\pm x^0 - i0) \cosh \varphi \pm i\mu x^1 \sinh \varphi}. \end{aligned} \quad (\text{C.3})$$

Then, the calculation of $D^{(\pm)}(x; \mu)$ runs in the way

$$\begin{aligned} D^{(\pm)}(x; \mu) &= \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi e^{-\mu \sqrt{-(x^0)^2 - (x^1)^2 \pm i0 \cdot x^0} \cosh(\varphi - \varphi_0)} = \\ &= \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh \varphi} = \pm \frac{i}{2\pi} K_0(\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)}), \end{aligned} \quad (\text{C.4})$$

where $x^2 = (x^0)^2 - (x^1)^2$, $K_0(z)$ is the McDonald function [34] and we have replaced $i0 x^0 \rightarrow i0 \cdot \varepsilon(x^0)$. For the calculation of $D^{(\pm)}(x; \mu)$ in (C.4) we have denoted

$$\cosh \varphi_0 = \frac{\pm x^0 - i0}{\sqrt{x^2 \mp i0 \cdot \varepsilon(x^0)}} \quad , \quad \sinh \varphi_0 = \frac{\pm x^1}{\sqrt{x^2 \mp i0 \cdot \varepsilon(x^0)}}. \quad (\text{C.5})$$

and made a shift $\varphi - \varphi_0 \rightarrow \varphi$.

In the limit $\mu \rightarrow 0$ the correlation function $D^{(\pm)}(x; \mu)$ is defined by

$$D^{(\pm)}(x; \mu) = \mp \frac{i}{4\pi} \ell n[-\mu^2 x^2 \pm i0 \cdot \varepsilon(x^0)], \quad (\text{C.6})$$

where we have used the asymptotic behaviour of the McDonald function $K_0(z)$ at $z \rightarrow 0$ [34]. Our result (C.6) agrees with that obtained in Ref.[16] (see Eqs.(2.6) and (2.11) of Ref.[16]).

For the correlation function $D_5^{(\pm)}(x; \mu)$ we get

$$\begin{aligned} D_5^{(\pm)}(x; \mu) &= \pm \frac{i}{4\pi} \int_{-\infty}^{\infty} d\varphi \varepsilon(\varphi) e^{-i\mu(\pm x^0 - i0) \cosh \varphi \pm i\mu x^1 \sinh \varphi} = \\ &= \pm \frac{i}{4\pi} \left[\int_{-\infty}^0 d\varphi \varepsilon(\varphi) e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh(\varphi - \varphi_0)} \right. \\ &\quad \left. + \int_0^{\infty} d\varphi \varepsilon(\varphi) e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh(\varphi - \varphi_0)} \right] = \\ &= \pm \frac{i}{4\pi} \left[- \int_0^{\infty} d\varphi e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh(\varphi + \varphi_0)} \right. \\ &\quad \left. + \int_0^{\infty} d\varphi e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh(\varphi - \varphi_0)} \right] = \\ &= \mp \frac{i}{4\pi} \int_{-\varphi_0}^{\varphi_0} d\varphi e^{-\mu \sqrt{-x^2 \pm i0 \cdot \varepsilon(x^0)} \cosh \varphi} = \mp \frac{i}{2\pi} \varphi_0 + O(\mu). \end{aligned} \quad (\text{C.7})$$

Taking into account that

$$\varphi_0 = \frac{1}{2} \ell n \left(\frac{x^0 + x^1 \mp i 0}{x^0 - x^1 \mp i 0} \right) \quad (\text{C.8})$$

we obtain

$$D_5^{(\pm)}(x) = \lim_{\mu \rightarrow 0} D_5^{(\pm)}(x; \mu) = \mp \frac{i}{4\pi} \ell n \left(\frac{x^0 + x^1 \mp i 0}{x^0 - x^1 \mp i 0} \right). \quad (\text{C.9})$$

Within the operator approach we apply these correlation functions to the derivation of quantum equations of motion of the massless Thirring model and the evaluation of n -point correlation functions of left(right)handed fermionic densities $\bar{\psi}(x)(1 \pm \gamma^5)\psi(x)$ of Coleman's proof of equivalence between the massive Thirring model and the sine-Gordon model [18].

The causal Green function $\Delta(x) = i\langle 0 | T(S(x)S(0)) | 0 \rangle = i\langle 0 | T(P(x)P(0)) | 0 \rangle$ is equal to

$$\begin{aligned} \Delta(x) &= \theta(x^0) D^{(+)}(x) - \theta(-x^0) D^{(-)}(x) = \\ &= -\frac{i}{4\pi} \theta(+x^0) \ell n[-\mu^2 x^2 + i 0 \cdot \varepsilon(x^0)] - \frac{i}{4\pi} \theta(-x^0) \ell n[-\mu^2 x^2 - i 0 \cdot \varepsilon(x^0)] = \\ &= -\frac{i}{4\pi} \ell n(-\mu^2 x^2 + i 0). \end{aligned} \quad (\text{C.10})$$

The same result can be obtained by using the momentum representation of $\Delta(x)$ [6,16]

$$\begin{aligned} \Delta(x) &= i \int \frac{d^2 p}{(2\pi)^2 i} \frac{e^{-ip \cdot x}}{\mu^2 - p^2 - i 0} = i \int_0^\infty \frac{dp_E}{(2\pi)^2} \frac{p_E}{\mu^2 + p_E^2} \int_0^{2\pi} d\varphi e^{ip_E \sqrt{-x^2 + i 0} \cos \varphi} = \\ &= i \int_0^\infty \frac{dp_E}{2\pi} \frac{p_E J_0(p_E \sqrt{-x^2 + i 0})}{\mu^2 + p_E^2} = \frac{i}{2\pi} K_0(\mu \sqrt{-x^2 + i 0}) = -\frac{i}{4\pi} \ell n(-\mu^2 x^2 + i 0). \end{aligned} \quad (\text{C.11})$$

By the Wick rotation $p^0 \rightarrow i p_2$ we have passed to Euclidean momentum space and used the integral representations for the Bessel and McDonald functions $J_0(z)$ and $K_0(z)$ (see p. 360, eq.(9.1.18) and p.488, eq.(11.4.44) of Ref.[34]).

Appendix D. On Klaiber's ansatz for the definition of the quantum vector fermionic current $K^\mu(x)$

In section 5 we have discussed the definition of the fermion vector current $K^\mu(x)$ determined by (5.37) and the analogous definition of the axial-vector current $K_5^\mu(x)$ given by (5.50). In this Appendix we would like to give some hints for our interpretation of Klaiber's ansatz (5.37). For this aim let us transcribe the Lagrangian (1.1) as follows

$$\mathcal{L}_{\text{Th}}(x) = \bar{\psi}(x) i \gamma^\mu (\partial_\mu + i g K_\mu(x)) \psi(x) + \frac{1}{2} g K_\mu(x) K^\mu(x). \quad (\text{D.1})$$

The equations of motion for this Lagrangian are equal to

$$\frac{\delta \mathcal{L}_{\text{Th}}(x)}{\delta \bar{\psi}(x)} = i \gamma^\mu (\partial_\mu + i g K_\mu(x)) \psi(x) = 0,$$

$$\frac{\delta \mathcal{L}_{\text{Th}}(x)}{\delta K^\mu(x)} = -g \bar{\psi}(x) \gamma^\mu \psi(x) + g K_\mu(x) = 0. \quad (\text{D.2})$$

Thus, the equations of motion read

$$i\gamma^\mu (\partial_\mu + i g K_\mu(x)) \psi(x) = 0 \quad (\text{D.3})$$

and

$$K_\mu(x) = \bar{\psi}(x) \gamma^\mu \psi(x). \quad (\text{D.4})$$

Substituting (D.4) in (D.3) we arrive at the equation of motion of the massless Thirring model

$$i\gamma^\mu \partial_\mu \psi(x) = g \gamma_\mu \psi(x) \bar{\psi}(x) \gamma^\mu \psi(x). \quad (\text{D.5})$$

The general form of the solution of the equation of motion (D.3) can be given in the form

$$\psi(x) = \exp \left\{ i g \int_{-\infty}^x dz^\lambda K_\lambda(z) \right\} \Psi(x), \quad (\text{D.6})$$

where the fermion field $\Psi(x)$ obeys a free Dirac equation

$$i\gamma^\mu \partial_\mu \Psi(x) = 0. \quad (\text{D.7})$$

The quantum fermionic vector current $K^\mu(x)$ should be defined in the normal-ordered form

$$K^\mu(x) = :\bar{\psi}(x) \gamma^\mu \psi(x):, \quad (\text{D.7})$$

which could be understood within the point-splitting technique as follows

$$K^\mu(x) = \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} [\bar{\psi}_a(x + \epsilon) \psi_b(x) - \psi_b(x + \epsilon) \bar{\psi}_a(x)]. \quad (\text{D.8})$$

Using the solution (D.6) of the equation of motion (D.3) and substituting it into (D.8) we obtain the quantum vector current $K^\mu(x)$ in the form

$$\begin{aligned} K^\mu(x) = \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} [\bar{\Psi}_a(x + \epsilon) e^{-i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \Psi_b(x) \\ - \Psi_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\Psi}_a(x)]. \end{aligned} \quad (\text{D.9})$$

Expanding the r.h.s. of (D.9) in powers of ϵ we get

$$\begin{aligned} K^\mu(x) = \\ = \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} \{ \bar{\Psi}_a(x + \epsilon) \Psi_b(x) [1 - i g \epsilon^\lambda K_\lambda(x)] - \Psi_b(x + \epsilon) \bar{\Psi}_a(x) [1 + i g \epsilon^\lambda K_\lambda(x)] \} = \\ = \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} [\bar{\Psi}_a(x + \epsilon) \Psi_b(x) - \Psi_b(x + \epsilon) \bar{\Psi}_a(x)] \\ + \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} \{ \bar{\Psi}_a(x + \epsilon) \Psi_b(x) + \Psi_b(x + \epsilon) \bar{\Psi}_a(x) \} (-i g) \epsilon^\lambda K_\lambda(x). \end{aligned} \quad (\text{D.10})$$

Using (5.28) and (5.34) we recast the r.h.s. of (D.10) into the form

$$K^\mu(x) = :\bar{\Psi}(x)\gamma^\mu\Psi(x): - \frac{g}{\pi} \frac{\epsilon^\mu \epsilon^\lambda}{\epsilon^2} K_\lambda(x). \quad (\text{D.11})$$

Averaging over the directions of the 2-vector ϵ and solving the obtained relation with respect to $K^\mu(x)$ we get

$$K^\mu(x) = \frac{1}{1 + \frac{g}{2\pi}} : \bar{\Psi}(x)\gamma^\mu\Psi(x) :. \quad (\text{D.12})$$

The quantum axial-vector current $K_5^\mu(x)$ can be defined in an analogous way

$$\begin{aligned} K_5^\mu(x) &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} [\bar{\psi}_a(x+\epsilon) \psi_b(x) - \psi_b(x+\epsilon) \bar{\psi}_a(x)] = \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} [\bar{\Psi}_a(x+\epsilon) e^{-ig \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \Psi_b(x) \\ &\quad - \Psi_b(x+\epsilon) e^{+ig \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\Psi}_a(x)] = \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} \{ \bar{\Psi}_a(x+\epsilon) \Psi_b(x) [1 - ig \epsilon^\lambda K_\lambda(x)] - \Psi_b(x+\epsilon) \bar{\Psi}_a(x) [1 + ig \epsilon^\lambda K_\lambda(x)] \} = \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} [\bar{\Psi}_a(x+\epsilon) \Psi_b(x) - \Psi_b(x+\epsilon) \bar{\Psi}_a(x)] \\ &+ \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} \{ \bar{\Psi}_a(x+\epsilon) \Psi_b(x) + \Psi_b(x+\epsilon) \bar{\Psi}_a(x) \} (-ig) \epsilon^\lambda K_\lambda(x) = \\ &=: \bar{\Psi}(x) \gamma^\mu \gamma^5 \Psi(x) : + \varepsilon^{\mu\nu} \frac{g}{\pi} \frac{\epsilon_\nu \epsilon^\lambda}{\epsilon^2} K_\lambda(x). \end{aligned} \quad (\text{D.13})$$

Averaging again over the directions of the 2-vector ϵ , using the identity $\varepsilon^{\mu\nu} \gamma_\nu = -\gamma^\mu \gamma^5$ and solving the result with respect to $K_5^\mu(x)$ we obtain

$$K_5^\mu(x) = \frac{1}{1 + \frac{g}{2\pi}} : \bar{\Psi}(x) \gamma^\mu \gamma^5 \Psi(x) :. \quad (\text{D.14})$$

One can see that the quantum vector and axial-vector currents $K^\mu(x)$ and $K_5^\mu(x)$ defined by (D.12) and (D.14), respectively, are related by the standard relation $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$.

Since according to the solution (D.6)) we are able to define $\Psi(x)$ in terms of $\psi(x)$

$$\Psi(x) = \exp \left\{ -ig \int_{-\infty}^x dz^\lambda K_\lambda(z) \right\} \psi(x), \quad (\text{D.15})$$

the quantum vector and axial-vector fermionic currents $K^\mu(x)$ and $K_5^\mu(x)$ can be determined in terms of the self-coupled field $\psi(x)$

$$K^\mu(x) = \frac{1}{1 + \frac{g}{2\pi}} : \bar{\psi}(x) \gamma^\mu \psi(x) : \quad , \quad K_5^\mu(x) = \frac{1}{1 + \frac{g}{2\pi}} : \bar{\psi}(x) \gamma^\mu \gamma^5 \psi(x) :. \quad (\text{D.16})$$

Making a finite renormalization of the wave function of the ψ -field, $\psi(x) \rightarrow Z_2^{1/2} \psi(x)$ we can remove the common factor in the definition of quantum vector and axial-vector current and obtain

$$K^\mu(x) =: \bar{\psi}(x) \gamma^\mu \psi(x) : \quad , \quad K_5^\mu(x) =: \bar{\psi}(x) \gamma^\mu \gamma^5 \psi(x) : . \quad (\text{D.17})$$

This testifies the self-consistency of the approach.

In turn, the inclusion of gauge dependent terms violates the self-consistency. Indeed, with such terms the definition of the vector current $K^\mu(x)$ given by (D.9) should read (see (5.45))

$$\begin{aligned} K^\mu(x) &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu)_{ab} [\bar{\Psi}_a(x + \epsilon) e^{-i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \Psi_b(x) \\ &\quad - \Psi_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\Psi}_a(x)] \\ &- i \sigma \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\hat{\epsilon})_{ab} [\bar{\Psi}_a(x + \epsilon) \Psi_b(x) + \Psi_b(x + \epsilon) \bar{\Psi}_a(x)] K^\mu(x) = \\ &=: \bar{\Psi}(x) \gamma^\mu \Psi(x) : - \frac{g}{\pi} \frac{\epsilon^\mu \epsilon^\lambda}{\epsilon^2} K_\lambda(x) - \frac{\sigma}{\pi} K^\mu(x). \end{aligned} \quad (\text{D.18})$$

Thus, unlike Klaiber stated, the contribution of the terms violating gauge invariance does not remove non-covariant term and an average over directions of the 2-vector ϵ is demanded. This yields

$$K^\mu(x) =: \bar{\Psi}(x) \gamma^\mu \Psi(x) : - \left(\frac{g}{2\pi} + \frac{\sigma}{\pi} \right) K^\mu(x). \quad (\text{D.19})$$

Solving this relation with respect to $K^\mu(x)$ we get a new expression for the quantum vector current $K^\mu(x)$

$$K^\mu(x) = \frac{1}{1 + \frac{g}{2\pi} + \frac{\sigma}{\pi}} : \bar{\Psi}(x) \gamma^\mu \Psi(x) : . \quad (\text{D.20})$$

In order to underscore the inconsistency of this extended definition of a quantum vector fermionic current we have to define a quantum axial-vector current in analogues way

$$\begin{aligned} K_5^\mu(x) &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\gamma^\mu \gamma^5)_{ab} [\bar{\Psi}_a(x + \epsilon) e^{-i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \Psi_b(x) \\ &\quad - \Psi_b(x + \epsilon) e^{+i g \int_x^{x+\epsilon} dz^\lambda K_\lambda(z)} \bar{\Psi}_a(x)] \\ &- i \sigma \lim_{\epsilon \rightarrow 0} \frac{1}{2} (\hat{\epsilon} \gamma^5)_{ab} [\bar{\Psi}_a(x + \epsilon) \Psi_b(x) + \Psi_b(x + \epsilon) \bar{\Psi}_a(x)] K^\mu(x). \end{aligned} \quad (\text{D.21})$$

The presence of the γ^5 -matrix thwarts the appearance of contributions proportional to σ and restores fully the former result (D.13). After averaging over directions of the 2-vector ϵ we get again (D.14). Due to the necessity to satisfy the standard relation $K_5^\mu(x) = -\varepsilon^{\mu\nu} K_\nu(x)$ the only solution for the parameter σ is $\sigma = 0$. This confirms our statement below (5.59).

Appendix E. Fourier transform of the two-point Green function of the massless Thirring fermion field

In this Appendix we calculate the Fourier transform of the two-point renormalized Green function of the massless Thirring fermion field defined by (11.18)

$$S_F(p) = \int d^2x \frac{M^2}{2\pi} \frac{-\hat{x} e^{+i p \cdot x}}{(-M^2 x^2 + i 0)^\lambda}, \quad (\text{E.1})$$

where we have denoted $\lambda = 1 + g/2\pi$.

The r.h.s. of (E.1) we transcribe in equivalent form

$$S_F(p) = -\frac{M^2}{2\pi i} \gamma^\mu \frac{\partial}{\partial p^\mu} \int d^2x \frac{e^{+i p \cdot x}}{(-M^2 x^2 + i 0)^\lambda} = \gamma^\mu \frac{\partial}{\partial p^\mu} \Delta(p), \quad (\text{E.2})$$

where $\Delta(p^0, p^1)$ is defined by

$$\Delta(p) = -\frac{M^2}{2\pi i} \int d^2x \frac{e^{+i p \cdot x}}{(-M^2 x^2 + i 0)^\lambda}. \quad (\text{E.4})$$

For the denominator we suggest to use the following integral representation

$$\frac{1}{(-M^2 x^2 + i 0)^\lambda} = \frac{(-i)^\lambda}{\Gamma(\lambda)} \int_0^\infty dt t^{\lambda-1} e^{+i(-M^2 x^2 + i 0)t}. \quad (\text{E.5})$$

Substituting (E.5) in (E.4) and integrating over x with the help of

$$\int d^2x e^{-i M^2 x^2 t + i p \cdot x} = \frac{\pi}{M^2 t} e^{+i(p^2 + i 0)/4M^2 t}. \quad (\text{E.6})$$

Thus, we define the function $\Delta(p)$ in terms of the integral over t

$$\Delta(p) = -\frac{1}{2i} \frac{(-i)^\lambda}{\Gamma(\lambda)} \int_0^\infty dt t^{\lambda-2} e^{+i(p^2 + i 0)/4M^2 t}. \quad (\text{E.7})$$

Integrating over t we obtain

$$\Delta(p) = -\frac{1}{2} e^{-i\pi\lambda} \frac{\Gamma(1-\lambda)}{\Gamma(\lambda)} \left(\frac{\bar{M}^2}{p^2 + i 0} \right)^{1-\lambda}, \quad (\text{E.8})$$

where $\bar{M} = 2M$.

Substituting (E.8) in (E.2) we derive the Fourier transform of the Green function (E.1)

$$S_F(p) = e^{-i\pi\lambda} \frac{\hat{p}}{p^2 + i 0} \frac{\Gamma(2-\lambda)}{\Gamma(\lambda)} \left(\frac{\bar{M}^2}{p^2 + i 0} \right)^{1-\lambda}, \quad (\text{E.9})$$

Setting $\lambda = 1 + g/2\pi$ we obtain the Fourier transform of the causal two-point Green function of massless Thirring fermion field

$$S_F(p) = -e^{-ig/2} \frac{\hat{p}}{p^2 + i 0} \frac{\Gamma(1-g/2\pi)}{\Gamma(1+g/2\pi)} \left(\frac{p^2 + i 0}{\bar{M}^2} \right)^{g/2\pi}. \quad (\text{E.10})$$

Using the properties of Γ -functions [34] we reduce $S_F(p)$ to the form

$$S_F(p) = -\frac{g e^{-ig/2}}{2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \frac{\hat{p}}{p^2 + i0} \left(\frac{p^2 + i0}{M^2}\right)^{\frac{g}{2\pi}}. \quad (\text{E.11})$$

In the limit $g \rightarrow 0$ we get the Green function of a free massless fermion field: $S_F(p) = \hat{p}/(-p^2 - i0)$.

The obtained result agrees well with Glaser's analysis of the massless Thirring model [30,31] and spectral analysis of two-point correlation functions in 1+1-dimensional quantum field theories developed by Schroer [32].

For the calculation of the fermion condensate it is convenient to use the double-integral representation for the two-point Green function $S_F(x)$ [33]:

$$\begin{aligned} S_F(x) &= i\langle 0 | T(\psi(x) \bar{\psi}(0)) | 0 \rangle = \frac{g^2 e^{-ig/2}}{\bar{M}^2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \\ &\times \int \frac{d^2 p}{(2\pi)^2} e^{-ip \cdot x} \int \frac{d^2 k}{(2\pi)^2 i} \frac{\hat{k}}{k^2 + i0} \left(\frac{(p-k)^2 + i0}{\bar{M}^2}\right)^{-1 + g/2\pi}. \end{aligned} \quad (\text{E.12})$$

Let us prove this representation. Making a shift of variables $p \rightarrow k + q$ we bring up the r.h.s. of (E.12) to the form

$$\begin{aligned} S_F(x) &= \frac{g^2 e^{-ig/2}}{\bar{M}^2 \sin\left(\frac{g}{2}\right)} \frac{1}{\Gamma^2\left(1 + \frac{g}{2\pi}\right)} \int \frac{d^2 k}{(2\pi)^2 i} \frac{\hat{k}}{k^2 + i0} e^{-ik \cdot x} \\ &\times \int \frac{d^2 q}{(2\pi)^2} \left(\frac{q^2 + i0}{\bar{M}^2}\right)^{-1 + g/2\pi} e^{-iq \cdot x}. \end{aligned} \quad (\text{E.13})$$

The integral over k is equal to

$$\int \frac{d^2 k}{(2\pi)^2 i} \frac{\hat{k}}{k^2 + i0} e^{-ik \cdot x} = -\gamma^\mu \frac{\partial}{\partial x^\mu} \Delta(x) = \frac{i}{2\pi} \frac{\hat{x}}{x^2 - i0}. \quad (\text{E.14})$$

For the integration over q we would use the representation analogous to (E.5) and change the order of integrations. This yields

$$\begin{aligned} &\int \frac{d^2 q}{(2\pi)^2} \left(\frac{q^2 + i0}{\bar{M}^2}\right)^{-1 + g/2\pi} e^{-iq \cdot x} = \frac{-i e^{ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \int_0^\infty \frac{dt}{t^{g/2\pi}} \\ &\times \int \frac{d^2 q}{(2\pi)^2} e^{it(q^2 + i0)/\bar{M}^2} e^{-iq \cdot x} = \frac{-i e^{ig/4}}{\Gamma\left(1 - \frac{g}{2\pi}\right)} \int_0^\infty \frac{dt}{t^{g/2\pi}} \frac{M^2}{\pi t} e^{i(-M^2 x^2 + i0)/t} = \end{aligned}$$

$$= -i e^{ig/2} \frac{M^2}{\pi} \frac{\Gamma\left(\frac{g}{2\pi}\right)}{\Gamma\left(1 - \frac{g}{2\pi}\right)} (-M^2 x^2 + i0)^{-g/2\pi}. \quad (\text{E.15})$$

Substituting (E.15) and (E.14) in (E.12) and using the properties of Γ -functions [34] we obtain

$$S_F(x) = \frac{1}{2\pi} \frac{\hat{x}}{x^2 - i0} (-M^2 x^2 + i0)^{-g/2\pi}. \quad (\text{E.16})$$

By the change $x \rightarrow x - y$ we bring up the r.h.s. of the expression (E.16) to the form of the Green function (11.18). We would like to accentuate that this agreement has been obtained due to shifts of variables – the procedure which is rather sensitive to convergence of integrals. For the derivation of the final expression (E.16) we have tacitly assumed that all integrals we have dealt with are convergent even if in the meaning of generalized functions [38].

Appendix F. Non-perturbative renormalizability of the sine-Gordon model

As has been shown in section 7 the massless Thirring model is equivalent to the sine-Gordon (SG) model, when the mass of Thirring fermion fields m is considered as an external source $\sigma(x) = -m$ of the scalar fermion density $\bar{\psi}(x)\psi(x)$. Therefore, the properties of non-perturbative renormalizability of the massless Thirring model investigated in section 11 should be fully extended to the SG model.

The generating functional of Green functions in the SG model we define as

$$Z_{\text{SG}}[J] = \int \mathcal{D}\vartheta \exp i \int d^2x \left\{ \frac{1}{2} \partial_\mu \vartheta(x) \partial^\mu \vartheta(x) + \frac{\bar{\alpha}}{\beta^2} (\cos \bar{\beta} \vartheta(x) - 1) + \vartheta(x) J(x) \right\}, \quad (\text{F.1})$$

where $J(x)$ is an external source of the $\vartheta(x)$ -field.

The Lagrangian of the SG model is invariant under the transformations

$$\vartheta(x) \rightarrow \vartheta'(x) = \vartheta(x) + \frac{2\pi n}{\bar{\beta}}, \quad (\text{F.2})$$

where n is an integer number running over $n = 0, \pm 1, \pm 2, \dots$. In order to get the generating functional $Z_{\text{SG}}[J]$ invariant under the transformations (F.2) it is sufficient to restrict the class of functions describing the external source of the ϑ -field and impose the constraint

$$\int d^2x J(x) = 0. \quad (\text{F.3})$$

Formally, for the pseudoscalar SG field $\vartheta(x)$, that we really have [6,18], the constraint (F.3) is fulfilled automatically. Indeed, due to the conservation of parity the external source of the SG field $\vartheta(x)$ should be a pseudoscalar quantity $J(x^0, x^1) = -J(x^0, -x^1)$. In the case of the scalar SG field $\vartheta(x)$ the situation with the fulfillment of the constraint (F.3) is a little bit more complicated but not crucial for the validity of it. Below we show that the

constraint (F.3) will be a great deal of importance for the infrared renormalizability of the SG model.

Non-perturbative renormalizability of the SG model we understand as a possibility to remove all divergences by renormalizing the coupling constant $\bar{\alpha}$. Indeed, since the coupling constant $\bar{\beta}$ is related to the coupling constant of the Thirring model g which is unrenormalized $g_0 = g$, so the coupling constant $\bar{\beta}$ should possess the same property, i.e. $\bar{\beta}_0 = \bar{\beta}$. Hence, only the coupling constant α should undergo renormalization.

The Lagrangian of the SG model written in terms of *bare* quantities reads

$$\mathcal{L}_{\text{SG}}(x) = \frac{1}{2} \partial_\mu \vartheta_0(x) \partial^\mu \vartheta_0(x) + \frac{\bar{\alpha}_0}{\bar{\beta}_0^2} (\cos \bar{\beta}_0 \vartheta_0(x) - 1). \quad (\text{F.4})$$

Since $\bar{\beta}$ is the unrenormalized coupling constant, the field $\vartheta(x)$ should be also unrenormalized, $\vartheta_0(x) = \vartheta(x)$. This means that there is no renormalization of the wave function of the ϑ -field. As a result the Lagrangian $\mathcal{L}_{\text{SG}}(x)$ of the SG model in terms of renormalized quantities can be written by

$$\begin{aligned} \mathcal{L}_{\text{SG}}(x) &= \frac{1}{2} \partial_\mu \vartheta(x) \partial^\mu \vartheta(x) + \frac{\bar{\alpha}}{\bar{\beta}^2} (\cos \bar{\beta} \vartheta(x) - 1) + (Z_1 - 1) \frac{\bar{\alpha}}{\bar{\beta}^2} (\cos \bar{\beta} \vartheta(x) - 1) = \\ &= \frac{1}{2} \partial_\mu \vartheta(x) \partial^\mu \vartheta(x) + Z_1 \frac{\bar{\alpha}}{\bar{\beta}^2} (\cos \bar{\beta} \vartheta(x) - 1), \end{aligned} \quad (\text{F.5})$$

where Z_1 is the renormalization constant of the coupling constant $\bar{\alpha}$. The renormalized coupling constant $\bar{\alpha}$ is related to the *bare* one by the relation [29]

$$\bar{\alpha} = Z_1^{-1} \bar{\alpha}_0. \quad (\text{F.6})$$

Renormalizability of the SG model as well as the Thirring model we understand as the possibility to replace the infrared cut-off μ and the ultra-violet cut-off Λ by another finite scale M by means of the renormalization constant Z_1 . According to the general theory of renormalizations [29] Z_1 should be a function of the coupling constants $\bar{\beta}$, $\bar{\alpha}$, the infrared cut-off μ , the ultra-violet cut-off Λ and a finite scale M :

$$Z_1 = Z_1(\bar{\beta}, \bar{\alpha}, M; \mu, \Lambda). \quad (\text{F.7})$$

Now let us proceed to the evaluation of the generating functional (F.1). For this aim, we first replace $\bar{\alpha} \rightarrow \bar{\alpha}_0$ and expand the integrand in powers of $\bar{\alpha}_0$. This gives

$$\begin{aligned} Z_{\text{SG}}[J] &= \lim_{\mu \rightarrow 0} e^{-i \int d^2 x \frac{\bar{\alpha}_0}{\bar{\beta}^2} \sum_{n=0}^{\infty} \frac{i^n}{n!} \left(\frac{\bar{\alpha}_0}{\bar{\beta}^2} \right)^n \prod_{i=1}^n \int d^2 x_i} \\ &\times \int \mathcal{D}\vartheta \prod_{i=1}^n \cos \bar{\beta} \vartheta(x_i) \exp i \int d^2 x \left\{ \frac{1}{2} \partial_\mu \vartheta(x) \partial^\mu \vartheta(x) - \frac{1}{2} \mu^2 \vartheta^2(x) + \vartheta(x) J(x) \right\}. \end{aligned} \quad (\text{F.8})$$

The integration over the ϑ -field can be carried out explicitly and we get

$$Z_{\text{SG}}[J] = \lim_{\mu \rightarrow 0} \sum_{n=0}^{\infty} \frac{i^n}{n!} \left(\frac{\bar{\alpha}_0}{2\bar{\beta}^2} \right)^n \sum_{p=0}^n \frac{n!}{(n-p)! p!} \prod_{j=1}^{n-p} \prod_{k=1}^p \int d^2 x_j d^2 y_k$$

$$\begin{aligned}
& \times \exp \left\{ \frac{1}{2} n \bar{\beta}^2 i\Delta(0) + \bar{\beta}^2 \sum_{j < k}^{n-p} i\Delta(x_j - x_k) + \bar{\beta}^2 \sum_{j < k}^p i\Delta(y_j - y_k) - \bar{\beta}^2 \sum_{j=1}^{n-p} \sum_{k=1}^p i\Delta(x_j - y_k) \right\} \\
& \times \exp \left\{ \int d^2x \bar{\beta} \left[\sum_{j=1}^{n-p} i\Delta(x_j - x) - \sum_{k=1}^p i\Delta(y_j - x) \right] J(x) + \iint d^2x d^2y \frac{1}{2} J(x) i\Delta(x-y) J(y) \right\},
\end{aligned} \tag{F.9}$$

where the Green functions $i\Delta(x-y)$ and $i\Delta(0)$ are defined by (6.19) and (6.20).

Taking the limit $\mu \rightarrow 0$ we reduce the r.h.s. of (F.6) to the form

$$\begin{aligned}
Z_{\text{SG}}[J] &= \sum_{p=0}^{\infty} \frac{(-1)^p}{(p!)^2} \left(\frac{\bar{\alpha}_0}{2\bar{\beta}^2} \right)^{2p} \prod_{j=1}^p \iint d^2x_j d^2y_j \left[\left(\frac{M}{\Lambda} \right)^{\bar{\beta}^2/2\pi} \right]^{2p} \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \sum_{j < k}^p \left(\ell n[-M^2(x_j - x_k)^2] + \ell n[-M^2(y_j - y_k)^2] \right) - \frac{\bar{\beta}^2}{4\pi} \sum_{j=1}^p \sum_{k=1}^p \ell n[-M^2(x_j - y_k)^2] \right\} \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \int d^2x \sum_{j=1}^p \ell n \left[\frac{(x_j - x)^2}{(y_j - x)^2} \right] J(x) + \frac{1}{8\pi} \iint d^2x d^2y J(x) \ell n[-M^2(x - y)^2] J(y) \right\} \\
&\times \lim_{\mu \rightarrow 0} \exp \left\{ -\frac{1}{4\pi} \ell n \left(\frac{M}{\mu} \right) \left(\int d^2x J(x) \right)^2 \right\}.
\end{aligned} \tag{F.10}$$

Due to the constraint (F.3) the generating functional $Z_{\text{SG}}[J]$ does not depend on the infrared cut-off. Using (F.3) we get

$$\begin{aligned}
Z_{\text{SG}}[J] &= \sum_{p=0}^{\infty} \frac{(-1)^p}{(p!)^2} \left(\frac{\bar{\alpha}_0}{2\bar{\beta}^2} \right)^{2p} \prod_{j=1}^p \iint d^2x_j d^2y_j \left[\left(\frac{M}{\Lambda} \right)^{\bar{\beta}^2/2\pi} \right]^{2p} \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \sum_{j < k}^p \left(\ell n[-M^2(x_j - x_k)^2] + \ell n[-M^2(y_j - y_k)^2] \right) - \frac{\bar{\beta}^2}{4\pi} \sum_{j=1}^p \sum_{k=1}^p \ell n[-M^2(x_j - y_k)^2] \right\} \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \int d^2x \sum_{j=1}^p \ell n \left[\frac{(x_j - x)^2}{(y_j - x)^2} \right] J(x) + \frac{1}{8\pi} \iint d^2x d^2y J(x) \ell n[-M^2(x - y)^2] J(y) \right\}.
\end{aligned} \tag{F.11}$$

Passing to a renormalized constant $\bar{\alpha}$ we recast the r.h.s. of (F.11) into the form

$$\begin{aligned}
Z_{\text{SG}}[J] &= \sum_{p=0}^{\infty} \left[\frac{1}{Z_1} \left(\frac{M}{\Lambda} \right)^{\bar{\beta}^2/2\pi} \right]^{2p} \frac{(-1)^p}{(p!)^2} \left(\frac{\bar{\alpha}}{2\bar{\beta}^2} \right)^{2p} \prod_{j=1}^p \iint d^2x_j d^2y_j \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \sum_{j < k}^p \left(\ell n[-M^2(x_j - x_k)^2] + \ell n[-M^2(y_j - y_k)^2] \right) - \frac{\bar{\beta}^2}{4\pi} \sum_{j=1}^p \sum_{k=1}^p \ell n[-M^2(x_j - y_k)^2] \right\} \\
&\times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \int d^2x \sum_{j=1}^p \ell n \left[\frac{(x_j - x)^2}{(y_j - x)^2} \right] J(x) + \frac{1}{8\pi} \iint d^2x d^2y J(x) \ell n[-M^2(x - y)^2] J(y) \right\}.
\end{aligned} \tag{F.12}$$

Setting

$$Z_1 = \left(\frac{M}{\Lambda} \right)^{\bar{\beta}^2/2\pi} \quad (\text{F.13})$$

we remove the dependence of the generating functional $Z_{\text{SG}}[J]$ on the ultra-violet cut-off Λ

$$\begin{aligned} Z_{\text{SG}}[J] = & \sum_{p=0}^{\infty} \frac{(-1)^p}{(p!)^2} \left(\frac{\bar{\alpha}}{2\bar{\beta}^2} \right)^{2p} \prod_{j=1}^p \iint d^2x_j d^2y_j \\ & \times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \sum_{j < k}^p \left(\ell n[-M^2(x_j - x_k)^2] + \ell n[-M^2(y_j - y_k)^2] \right) - \frac{\bar{\beta}^2}{4\pi} \sum_{j=1}^p \sum_{k=1}^p \ell n[-M^2(x_j - y_k)^2] \right\} \\ & \times \exp \left\{ \frac{\bar{\beta}^2}{4\pi} \int d^2x \sum_{j=1}^p \ell n \left[\frac{(x_j - x)^2}{(y_j - x)^2} \right] J(x) + \frac{1}{8\pi} \iint d^2x d^2y J(x) \ell n[-M^2(x - y)^2] J(y) \right\}. \end{aligned} \quad (\text{F.14})$$

However, for the evaluation of correlation functions there appears the causal Green function $\Delta(x - y; M)$ taken at $x = y$. For the definition of $\Delta(0; M)$ one can use dimensional or analytical regularization procedures allowing to set $\Delta(0; M) = 0$ [6]. As a result no divergences appear for the evaluation of any correlation function of the SG model.

The generating functional (F.14) is expressed in terms of the renormalized constant $\bar{\alpha}$, the constant $\bar{\beta}$ and the finite scale M . Now it can be applied to the evaluation of any renormalized correlation function of the SG model. This testifies the complete non-perturbative renormalizability of the SG model.

We would like to emphasize that due to unrenormalizability of the coupling constant, $\bar{\beta}_0 = \bar{\beta}$, the Gell-Mann-Low β -function vanishes, $\beta(\bar{\beta}, M) = 0$. This means that the SG model is not an asymptotically free theory that would require non-zero negative value for the Gell-Mann-Low function, $\beta(\bar{\beta}, M) < 0$. Our result disagrees with that obtained within perturbation theory [39].

The unrenormalizability of the coupling constant $\bar{\beta}$ leading to the vanishing Gell-Mann-Low function can be easily understood following the similarity between $\bar{\beta}$ and $\hbar$ which has been drawn in Ref.[6]. As has been shown in Ref.[6] the limits $\bar{\beta} \rightarrow 0$ and $\bar{\beta} \rightarrow \infty$ distinguish *classical* and *quantum* regimes of the sine-Gordon model. Within such an understanding of the coupling constant $\bar{\beta}$ its unrenormalizability is justified by the unrenormalizability of $\hbar$.

The same procedure which we have applied to the renormalization of the SG model can be implemented to the formulation of a quantum field theory of a free massless (pseudo)scalar field in 1+1-dimensional space-time free of the *infrared* problem pointed out by Klaiber [4] and Coleman [40]. This program we are planning to realize in forthcoming publications.

References

- [1] W. Thirring, *Ann. Phys. (N.Y.)* **3**, 91 (1958).
- [2] C. Itzykson and J.-B. Zuber, in *QUANTUM FIELD THEORY*, McGraw-Hill Book Company, New York 1980.
- [3] J. D. Bjorken and S. D. Drell, in *RELATIVISTIC QUANTUM FIELDS*, McGraw-Hill Book Company, New York, 1965.
- [4] B. Klaiber, in *LECTURES IN THEORETICAL PHYSICS*, Lectures delivered at the Summer Institute for Theoretical Physics, University of Colorado, Boulder, 1967, edited by A. Barut and W. Brittin, Gordon and Breach, New York, 1968, Vol. X, part A, pp.141–176.
- [5] J. H. Lowenstein, *Commun. Math. Phys.* **16**, 265 (1970).
- [6] M. Faber and A. N. Ivanov, *Eur. Phys. J. C* **20**, 723 (2001), hep-th/0105057.
- [7] K. Furuya, Re. E. Gamboa Saravi and F. A. Schaposnik, *Nucl. Phys. B* **208**, 159 (1982).
- [8] C. M. Naón, *Phys. Rev. D* **31**, 2035 (1985).
- [9] K. Fujikawa, *Phys. Rev. Lett.* **44**, 1195 (1979); *Phys. Rev. D* **21**, 2848 (1982); *Phys. Rev. D* **22**, 1499(E) (1980).
- [10] R. Roskies and F. A. Schaposnik, *Phys. Rev. D* **23**, 558 (1981); R. E. Gamboa Saravi, F. A. Schaposnik, and J. E. Solomin, *Nucl. Phys. B* **153**, 112 (1979); R. E. Gamboa Savari, M. A. Muschetti, F. A. Schaposnik, and J. E. Solomin, *Ann. of Phys. (N.Y.)* **157**, 360 (1984); O. Alvarez, *Nucl. Phys. B* **238**, 61 (1984); H. Dorn, *Phys. Lett. B* **167**, 86 (1986).
- [11] G. V. Christos, *Z. Phys. C* **18**, 155 (1983).
- [12] R. Banerjee, *Z. Phys. C* **25**, 251 (1984); T. Ikehashi, *Phys. Lett. B* **313**, 103 (1993).
- [13] R. Jackiw and R. Rajaraman, *Phys. Rev. Lett.* **54**, 1219 (1985); K. Harada, H. Kubota, and I. Tsutsui, *Phys. Lett. B* **173**, 77 (1986).
- [14] R. A. Bertlmann, in *ANOMALIES IN QUANTUM FIELD THEORY*, Oxford Science Publications, Clarendon Press • Oxford, 1996, pp.248–286.
- [15] P. H. Damgaard, H. B. Nielsen, and R. Sollacher, *Nucl. Phys. B* **385**, 227 (1992).
- [16] E. Abdala, M. C. B. Abdala, and K. D. Rothe, in *Non-perturbative methods in 2 DIMENSIONAL QUANTUM FIELD THEORY*, World Scientific, Singapore, 1991.
- [17] J. Zinn-Justin, in *QUANTUM FIELD THEORY AND CRITICAL PHENOMENA*, Clarendon Press•Oxford, 1996.
- [18] S. Coleman, *Phys. Rev. D* **11**, 2088 (1975).

- [19] A. H. Mueller and T. L. Trueman, Phys. Rev. D **4**, 1635 (1971).
- [20] M. Gomes and J. H. Lowenstein, Nucl. Phys. B **45**, 252 (1972).
- [21] C. G. Callan, R. F. Dashen, and D. H. Sharp, Phys. Rev. **165**, 1883 (1965).
- [22] J. Schwinger, Phys. Rev. Lett. **3**, 296 (1959).
- [23] C. M. Sommerfield, Ann. of Phys. (N.Y.) **26**, 1 (1963).
- [24] G. C. Wick, Phys. Rev. **80**, 268 (1950).
- [25] Y. Takahashi, Nuovo Cimento A **47**, 392 (1978).
- [26] M. Faber, A. N. Ivanov, N. I. Troitskaya, and M. Zach, Phys. Rev. D **62** (2000) 025019; M. Faber, A. N. Ivanov, and N. I. Troitskaya, hep-th/0012083, hep-th/0012147v3.
- [27] B. Klaiber, Helv. Phys. Acta, **37**, 554 (1964).
- [28] K. G. Wilson, Phys. Rev. **179**, 1499 (1969); Phys. Rev. D **2**, 1473 (1970).
- [29] J. C. Collins, in *RENORMALIZATION, An Introduction to Renormalization, the Renormalization Group, and the Operator-Product Expansion*, Cambridge University Press, Cambridge, 1984.
- [30] V. Glaser, Nuovo Cim. **9**, 990 (1958).
- [31] W. Thirring, Nuovo Cim. **9**, 1007 (1958).
- [32] B. Schroer, Fortschritte der Physik, **11**, 1 (1963).
- [33] A. J. de Silva, M. Gomes, and R. Köberle, Phys. Rev. D **20**, 895 (1979).
- [34] *HANDBOOK OF MATHEMATICAL FUNCTIONS, with Formulas, Graphs, and Mathematical Tables*, ed. by M. Abramowitz and I. E. Stegun, U.S. Department of Commerce, National Bureau of Standards, Applied Mathematics Series • 55, 1972.
- [35] I. S. Gradshteyn and I. M. Ryzik, in *TABLE OF INTEGRALS, SERIES, AND PRODUCTS*, Academic Press, New York & London, 1965.
- [36] K. Johnson, Nuovo Cim. **20**, 773 (1961).
- [37] F. L. Scarf and J. Wess, Nuovo Cim. **26**, 150 (1962)
- [38] I. M. Gel'fand and G. E. Shilov, in *GENERALIZED FUNCTIONS, Properties and Operations*, Vol. I, Academic Press, New York and London, 1964.
- [39] D. J. Amit, Y. Y. Goldschmidt, and G. Grinstein, J. Phys. A **13**, 585 (1980); J. Balog and Á. Hegedűs, J. Phys. A **33**, 6543 (2000), hep-th/0003258.
- [40] S. Coleman, Comm. Math. Phys. **31**, 259 (1973).